

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
29 November 2001 (29.11.2001)

PCT

(10) International Publication Number
WO 01/90152 A2

(51) International Patent Classification⁷: **C07K 14/00**

(21) International Application Number: **PCT/US01/16776**

(22) International Filing Date: **22 May 2001 (22.05.2001)**

(25) Filing Language: **English**

(26) Publication Language: **English**

(30) Priority Data:

09/577,505	24 May 2000 (24.05.2000)	US
09/590,583	8 June 2000 (08.06.2000)	US
09/699,295	26 October 2000 (26.10.2000)	US
09/810,936	16 March 2001 (16.03.2001)	US


(72) Inventors; and

(75) Inventors/Applicants (for US only): **FRUDAKIS, Tony, N.** [US/US]; 7937 Broadmoor Pines Blvd., Sarasota, FL 34243 (US). **REED, Steven, G.** [US/US]; 2843 122nd Place N.E., Bellevue, WA 98005 (US). **SMITH, John, M.** [US/US]; 1101 Pierce Ter N.E., Columbia Heights, MN 55421 (US). **MISHER, Lynda, E.** [US/US]; 5043 16th Avenue N.E., Seattle, WA 98105 (US). **DILLON, Davin, C.** [US/US]; 18112 N.W. Montreux Drive, Issaquah, WA 98027 (US). **RETTTER, Marc, W.** [US/US]; 33402 N.E. 43rd Place, Carnation, WA 98014 (US). **WANG, Aijun** [CN/US]; 3106 213th Place S.E., Issaquah, WA 98029 (US). **SKEIKY, Yasir, A., W.** [LB/US]; 15106 S.E. 47th Place, Bellevue, WA 98006 (US). **HARLOCKER, Susan, L.** [US/US]; 7522 13th Avenue W., Seattle, WA 98117 (US). **DAY, Craig, H.** [US/US]; 11501 Stone Avenue N., C122, Seattle, WA 98133-8317 (US).

(71) Applicant (for all designated States except US): **CORIXA CORPORATION** [US/US]; 1124 Columbia Street, Suite 200, Seattle, WA 98104 (US).

[Continued on next page]

(54) Title: **COMPOSITIONS AND METHODS FOR THE THERAPY AND DIAGNOSIS OF BREAST CANCER**



(57) Abstract: Compositions and methods for the therapy and diagnosis of cancer, particularly breast cancer, are disclosed. Illustrative compositions comprise one or more breast tumor polypeptides, immunogenic portions thereof, polynucleotides that encode such polypeptides, antigen presenting cell that expresses such polypeptides, and T cells that are specific for cells expressing such polypeptides. The disclosed compositions are useful, for example, in the diagnosis, prevention and/or treatment of diseases, particularly breast cancer.

WO 01/90152 A2



(74) **Agents:** POTTER, Jane, E., R.; Seed Intellectual Property Law Group PLLC, Suite 6300, 701 Fifth Avenue, Seattle, WA 98104-7092 et al. (US).

(81) **Designated States (national):** AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) **Designated States (regional):** ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

COMPOSITIONS AND METHODS FOR THE THERAPY AND DIAGNOSIS OF BREAST CANCER

TECHNICAL FIELD OF THE INVENTION

The present invention relates generally to therapy and diagnosis of
5 cancer, such as breast cancer. The invention is more specifically related to
polypeptides, comprising at least a portion of a breast tumor protein, and to
polynucleotides encoding such polypeptides. Such polypeptides and polynucleotides
are useful in pharmaceutical compositions, *e.g.*, vaccines, and other compositions for
the diagnosis and treatment of breast cancer.

10 BACKGROUND OF THE INVENTION

Breast cancer is a significant health problem for women in the United
States and throughout the world. Although advances have been made in detection and
treatment of the disease, breast cancer remains the second leading cause of cancer-
related deaths in women, affecting more than 180,000 women in the United States each
15 year. For women in North America, the life-time odds of getting breast cancer are now
one in eight.

No vaccine or other universally successful method for the prevention or
treatment of breast cancer is currently available. Management of the disease currently
relies on a combination of early diagnosis (through routine breast screening procedures)
20 and aggressive treatment, which may include one or more of a variety of treatments
such as surgery, radiotherapy, chemotherapy and hormone therapy. The course of
treatment for a particular breast cancer is often selected based on a variety of prognostic
parameters, including an analysis of specific tumor markers. *See, e.g.,* Porter-Jordan
and Lippman, *Breast Cancer* 8:73-100 (1994). However, the use of established markers
25 often leads to a result that is difficult to interpret, and the high mortality observed in
breast cancer patients indicates that improvements are needed in the treatment,
diagnosis and prevention of the disease.

Accordingly, there is a need in the art for improved methods for therapy and diagnosis of breast cancer. The present invention fulfills these needs and further provides other related advantages.

SUMMARY OF THE INVENTION

5 In one aspect, the present invention provides polynucleotide compositions comprising a sequence selected from the group consisting of:

- (a) sequences provided in SEQ ID NO: 1, 3-86, 142-298, 301-303, 307, 313, 314, 316, 317, 325 and 327-330;
- (b) complements of the sequences provided in SEQ ID NO: 1, 3-86,
10 142-298, 301-303, 307, 313, 314, 316, 317, 325 and 327-330;
- (c) sequences consisting of at least 20 contiguous residues of a sequence provided in SEQ ID NO: 1, 3-86, 142-298, 301-303, 307, 313, 314, 316, 317, 325 and 327-330;
- (d) sequences that hybridize to a sequence provided in SEQ ID NO:
15 1, 3-86, 142-298, 301-303, 307, 313, 314, 316, 317, 325 and 327-330, under moderately stringent conditions;
- (e) sequences having at least 75% identity to a sequence of SEQ ID NO: 1, 3-86, 142-298, 301-303, 307, 313, 314, 316, 317, 325 and 327-330;
- (f) sequences having at least 90% identity to a sequence of SEQ ID
20 NO: 1, 3-86, 142-298, 301-303, 307, 313, 314, 316, 317, 325 and 327-330; and
- (g) degenerate variants of a sequence provided in SEQ ID NO: 1, 3-86, 142-298, 301-303, 307, 313, 314, 316, 317, 325 and 327-330.

In one preferred embodiment, the polynucleotide compositions of the
25 invention are expressed in at least about 20%, more preferably in at least about 30%, and most preferably in at least about 50% of breast tumors samples tested, at a level that is at least about 2-fold, preferably at least about 5-fold, and most preferably at least about 10-fold higher than that for normal tissues.

The present invention, in another aspect, provides polypeptide compositions comprising an amino acid sequence that is encoded by a polynucleotide sequence described above.

The present invention further provides polypeptide compositions
5 comprising an amino acid sequence selected from the group consisting of sequences recited in SEQ ID NO: 299, 300, 304-306, 308-312, 314, 326 and 331-334.

In certain preferred embodiments, the polypeptides and/or polynucleotides of the present invention are immunogenic, *i.e.*, they are capable of eliciting an immune response, particularly a humoral and/or cellular immune response,
10 as further described herein.

The present invention further provides fragments, variants and/or derivatives of the disclosed polypeptide and/or polynucleotide sequences, wherein the fragments, variants and/or derivatives preferably have a level of immunogenic activity of at least about 50%, preferably at least about 70% and more preferably at least about
15 90% of the level of immunogenic activity of a polypeptide sequence set forth in SEQ ID NOs: 299, 300, 304-306, 308-312, 314, 326 and 331-334 or a polypeptide sequence encoded by a polynucleotide sequence set forth in SEQ ID NOs: 1, 3-86, 142-298, 301-303, 307, 313, 314, 316, 317, 325 and 327-330.

The present invention further provides polynucleotides that encode a
20 polypeptide described above, expression vectors comprising such polynucleotides and host cells transformed or transfected with such expression vectors.

Within other aspects, the present invention provides pharmaceutical compositions comprising a polypeptide or polynucleotide as described above and a physiologically acceptable carrier.

25 Within a related aspect of the present invention, the pharmaceutical compositions, *e.g.*, vaccine compositions, are provided for prophylactic or therapeutic applications. Such compositions generally comprise an immunogenic polypeptide or polynucleotide of the invention and an immunostimulant, such as an adjuvant.

The present invention further provides pharmaceutical compositions that
30 comprise: (a) an antibody or antigen-binding fragment thereof that specifically binds to

a polypeptide of the present invention, or a fragment thereof; and (b) a physiologically acceptable carrier.

Within further aspects, the present invention provides pharmaceutical compositions comprising: (a) an antigen presenting cell that expresses a polypeptide as described above and (b) a pharmaceutically acceptable carrier or excipient. Illustrative antigen presenting cells include dendritic cells, macrophages, monocytes, fibroblasts and B cells.

Within related aspects, pharmaceutical compositions are provided that comprise: (a) an antigen presenting cell that expresses a polypeptide as described above and (b) an immunostimulant.

The present invention further provides, in other aspects, fusion proteins that comprise at least one polypeptide as described above, as well as polynucleotides encoding such fusion proteins, typically in the form of pharmaceutical compositions, *e.g.*, vaccine compositions, comprising a physiologically acceptable carrier and/or an immunostimulant. The fusions proteins may comprise multiple immunogenic polypeptides or portions/variants thereof, as described herein, and may further comprise one or more polypeptide segments for facilitating the expression, purification and/or immunogenicity of the polypeptide(s).

Within further aspects, the present invention provides methods for stimulating an immune response in a patient, preferably a T cell response in a human patient, comprising administering a pharmaceutical composition described herein. The patient may be afflicted with breast cancer, in which case the methods provide treatment for the disease, or patient considered at risk for such a disease may be treated prophylactically.

Within further aspects, the present invention provides methods for inhibiting the development of a cancer in a patient, comprising administering to a patient a pharmaceutical composition as recited above. The patient may be afflicted with breast cancer, in which case the methods provide treatment for the disease, or patient considered at risk for such a disease may be treated prophylactically.

The present invention further provides, within other aspects, methods for removing tumor cells from a biological sample, comprising contacting a biological sample with T cells that specifically react with a polypeptide of the present invention, wherein the step of contacting is performed under conditions and for a time sufficient to
5 permit the removal of cells expressing the protein from the sample.

Within related aspects, methods are provided for inhibiting the development of a cancer in a patient, comprising administering to a patient a biological sample treated as described above.

Methods are further provided, within other aspects, for stimulating
10 and/or expanding T cells specific for a polypeptide of the present invention, comprising contacting T cells with one or more of: (i) a polypeptide as described above; (ii) a polynucleotide encoding such a polypeptide; and/or (iii) an antigen presenting cell that expresses such a polypeptide; under conditions and for a time sufficient to permit the stimulation and/or expansion of T cells. Isolated T cell populations comprising T cells
15 prepared as described above are also provided.

Within further aspects, the present invention provides methods for inhibiting the development of a cancer in a patient, comprising administering to a patient an effective amount of a T cell population as described above.

The present invention further provides methods for inhibiting the
20 development of a cancer in a patient, comprising the steps of: (a) incubating CD4⁺ and/or CD8⁺ T cells isolated from a patient with one or more of: (i) a polypeptide comprising at least an immunogenic portion of polypeptide disclosed herein; (ii) a polynucleotide encoding such a polypeptide; and (iii) an antigen-presenting cell that expressed such a polypeptide; and (b) administering to the patient an effective amount
25 of the proliferated T cells, and thereby inhibiting the development of a cancer in the patient. Proliferated cells may, but need not, be cloned prior to administration to the patient.

Within further aspects, the present invention provides methods for determining the presence or absence of a cancer, preferably a breast cancer, in a patient
30 comprising: (a) contacting a biological sample obtained from a patient with a binding

agent that binds to a polypeptide as recited above; (b) detecting in the sample an amount of polypeptide that binds to the binding agent; and (c) comparing the amount of polypeptide with a predetermined cut-off value, and therefrom determining the presence or absence of a cancer in the patient. Within preferred embodiments, the binding agent
5 is an antibody, more preferably a monoclonal antibody.

The present invention also provides, within other aspects, methods for monitoring the progression of a cancer in a patient. Such methods comprise the steps of: (a) contacting a biological sample obtained from a patient at a first point in time with a binding agent that binds to a polypeptide as recited above; (b) detecting in the
10 sample an amount of polypeptide that binds to the binding agent; (c) repeating steps (a) and (b) using a biological sample obtained from the patient at a subsequent point in time; and (d) comparing the amount of polypeptide detected in step (c) with the amount detected in step (b) and therefrom monitoring the progression of the cancer in the patient.

15 The present invention further provides, within other aspects, methods for determining the presence or absence of a cancer in a patient, comprising the steps of: (a) contacting a biological sample obtained from a patient with an oligonucleotide that hybridizes to a polynucleotide that encodes a polypeptide of the present invention; (b) detecting in the sample a level of a polynucleotide, preferably mRNA, that hybridizes to
20 the oligonucleotide; and (c) comparing the level of polynucleotide that hybridizes to the oligonucleotide with a predetermined cut-off value, and therefrom determining the presence or absence of a cancer in the patient. Within certain embodiments, the amount of mRNA is detected via polymerase chain reaction using, for example, at least one oligonucleotide primer that hybridizes to a polynucleotide encoding a polypeptide as
25 recited above, or a complement of such a polynucleotide. Within other embodiments, the amount of mRNA is detected using a hybridization technique, employing an oligonucleotide probe that hybridizes to a polynucleotide that encodes a polypeptide as recited above, or a complement of such a polynucleotide.

In related aspects, methods are provided for monitoring the progression
30 of a cancer in a patient, comprising the steps of: (a) contacting a biological sample

obtained from a patient with an oligonucleotide that hybridizes to a polynucleotide that encodes a polypeptide of the present invention; (b) detecting in the sample an amount of a polynucleotide that hybridizes to the oligonucleotide; (c) repeating steps (a) and (b) using a biological sample obtained from the patient at a subsequent point in time; and
5 (d) comparing the amount of polynucleotide detected in step (c) with the amount detected in step (b) and therefrom monitoring the progression of the cancer in the patient.

Within further aspects, the present invention provides antibodies, such as monoclonal antibodies, that bind to a polypeptide as described above, as well as
10 diagnostic kits comprising such antibodies. Diagnostic kits comprising one or more oligonucleotide probes or primers as described above are also provided.

These and other aspects of the present invention will become apparent upon reference to the following detailed description and attached drawings. All references disclosed herein are hereby incorporated by reference in their entirety as if
15 each was incorporated individually.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows the differential display PCR products, separated by gel electrophoresis, obtained from cDNA prepared from normal breast tissue (lanes 1 and 2) and from cDNA prepared from breast tumor tissue from the same patient (lanes 3 and
20 4). The arrow indicates the band corresponding to B18Ag1.

Figure 2 is a northern blot comparing the level of B18Ag1 mRNA in breast tumor tissue (lane 1) with the level in normal breast tissue.

Figure 3 shows the level of B18Ag1 mRNA in breast tumor tissue compared to that in various normal and non-breast tumor tissues as determined by
25 RNase protection assays.

Figure 4 is a genomic clone map showing the location of additional retroviral sequences obtained from ends of XbaI restriction digests (provided in SEQ ID NO:3 - SEQ ID NO:10) relative to B18Ag1.

Figures 5A and 5B show the sequencing strategy, genomic organization and predicted open reading frame for the retroviral element containing B18Ag1.

Figure 6 shows the nucleotide sequence of the representative breast tumor-specific cDNA B18Ag1.

5 Figure 7 shows the nucleotide sequence of the representative breast tumor-specific cDNA B17Ag1.

Figure 8 shows the nucleotide sequence of the representative breast tumor-specific cDNA B17Ag2.

Figure 9 shows the nucleotide sequence of the representative breast
10 tumor-specific cDNA B13Ag2a.

Figure 10 shows the nucleotide sequence of the representative breast tumor-specific cDNA B13Ag1b.

Figure 11 shows the nucleotide sequence of the representative breast tumor-specific cDNA B13Ag1a.

15 Figure 12 shows the nucleotide sequence of the representative breast tumor-specific cDNA B11Ag1.

Figure 13 shows the nucleotide sequence of the representative breast tumor-specific cDNA B3CA3c.

Figure 14 shows the nucleotide sequence of the representative breast
20 tumor-specific cDNA B9CG1.

Figure 15 shows the nucleotide sequence of the representative breast tumor-specific cDNA B9CG3.

Figure 16 shows the nucleotide sequence of the representative breast tumor-specific cDNA B2CA2.

25 Figure 17 shows the nucleotide sequence of the representative breast tumor-specific cDNA B3CA1.

Figure 18 shows the nucleotide sequence of the representative breast tumor-specific cDNA B3CA2.

Figure 19 shows the nucleotide sequence of the representative breast
30 tumor-specific cDNA B3CA3.

Figure 20 shows the nucleotide sequence of the representative breast tumor-specific cDNA B4CA1.

Figure 21A depicts RT-PCR analysis of breast tumor genes in breast tumor tissues (lanes 1-8) and normal breast tissues (lanes 9-13) and H₂O (lane 14).

5 Figure 21B depicts RT-PCR analysis of breast tumor genes in prostate tumors (lane 1, 2), colon tumors (lane 3), lung tumor (lane 4), normal prostate (lane 5), normal colon (lane 6), normal kidney (lane 7), normal liver (lane 8), normal lung (lane 9), normal ovary (lanes 10, 18), normal pancreases (lanes 11, 12), normal skeletal muscle (lane 13), normal skin (lane 14), normal stomach (lane 15), normal testes (lane 10 16), normal small intestine (lane 17), HBL-100 (lane 19), MCF-12A (lane 20), breast tumors (lanes 21-23), H₂O (lane 24), and colon tumor (lane 25).

Figure 22 shows the recognition of a B11Ag1 peptide (referred to as B11-8) by an anti-B11-8 CTL line.

Figure 23 shows the recognition of a cell line transduced with the antigen 15 B11Ag1 by the B11-8 specific clone A1.

Figure 24 shows recognition of a lung adenocarcinoma line (LT-140-22) and a breast adenocarcinoma line (CAMA-1) by the B11-8 specific clone A1.

DETAILED DESCRIPTION OF THE INVENTION

The present invention is directed generally to compositions and their use 20 in the therapy and diagnosis of cancer, particularly breast cancer. As described further below, illustrative compositions of the present invention include, but are not restricted to, polypeptides, particularly immunogenic polypeptides, polynucleotides encoding such polypeptides, antibodies and other binding agents, antigen presenting cells (APCs) and immune system cells (*e.g.*, T cells).

25 The practice of the present invention will employ, unless indicated specifically to the contrary, conventional methods of virology, immunology, microbiology, molecular biology and recombinant DNA techniques within the skill of the art, many of which are described below for the purpose of illustration. Such techniques are explained fully in the literature. See, *e.g.*, Sambrook, et al. Molecular

Cloning: A Laboratory Manual (2nd Edition, 1989); Maniatis et al. Molecular Cloning: A Laboratory Manual (1982); DNA Cloning: A Practical Approach, vol. I & II (D. Glover, ed.); Oligonucleotide Synthesis (N. Gait, ed., 1984); Nucleic Acid Hybridization (B. Hames & S. Higgins, eds., 1985); Transcription and Translation (B. Hames & S. Higgins, eds., 1984); Animal Cell Culture (R. Freshney, ed., 1986); Perbal, A Practical Guide to Molecular Cloning (1984).

All publications, patents and patent applications cited herein, whether supra or infra, are hereby incorporated by reference in their entirety.

As used in this specification and the appended claims, the singular forms "a," "an" and "the" include plural references unless the content clearly dictates otherwise.

Polypeptide Compositions

As used herein, the term "polypeptide" is used in its conventional meaning, *i.e.*, as a sequence of amino acids. The polypeptides are not limited to a specific length of the product; thus, peptides, oligopeptides, and proteins are included within the definition of polypeptide, and such terms may be used interchangeably herein unless specifically indicated otherwise. This term also does not refer to or exclude post-expression modifications of the polypeptide, for example, glycosylations, acetylations, phosphorylations and the like, as well as other modifications known in the art, both naturally occurring and non-naturally occurring. A polypeptide may be an entire protein, or a subsequence thereof. Particular polypeptides of interest in the context of this invention are amino acid subsequences comprising epitopes, *i.e.*, antigenic determinants substantially responsible for the immunogenic properties of a polypeptide and being capable of evoking an immune response.

Particularly illustrative polypeptides of the present invention comprise those encoded by a polynucleotide sequence set forth in any one of SEQ ID NOs: 1, 3-86, 142-298, 301-303, 307, 313, 314, 316, 317, 325 and 327-330, or a sequence that hybridizes under moderately stringent conditions, or, alternatively, under highly stringent conditions, to a polynucleotide sequence set forth in any one of SEQ ID NOs:

1, 3-86, 142-298, 301-303, 307, 313, 314, 316, 317, 325 and 327-330. Certain other illustrative polypeptides of the invention comprise amino acid sequences as set forth in any one of SEQ ID NOs: 299, 300, 304-306, 308-312, 314, 326 and 331-334.

The polypeptides of the present invention are sometimes herein referred
5 to as breast tumor proteins or breast tumor polypeptides, as an indication that their identification has been based at least in part upon their increased levels of expression in breast tumor samples. Thus, a "breast tumor polypeptide" or "breast tumor protein," refers generally to a polypeptide sequence of the present invention, or a polynucleotide sequence encoding such a polypeptide, that is expressed in a substantial proportion of
10 breast tumor samples, for example preferably greater than about 20%, more preferably greater than about 30%, and most preferably greater than about 50% or more of breast tumor samples tested, at a level that is at least two fold, and preferably at least five fold, greater than the level of expression in normal tissues, as determined using a representative assay provided herein. A breast tumor polypeptide sequence of the
15 invention, based upon its increased level of expression in tumor cells, has particular utility both as a diagnostic marker as well as a therapeutic target, as further described below.

In certain preferred embodiments, the polypeptides of the invention are immunogenic, *i.e.*, they react detectably within an immunoassay (such as an ELISA or
20 T-cell stimulation assay) with antisera and/or T-cells from a patient with breast cancer. Screening for immunogenic activity can be performed using techniques well known to the skilled artisan. For example, such screens can be performed using methods such as those described in Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, 1988. In one illustrative example, a polypeptide may be
25 immobilized on a solid support and contacted with patient sera to allow binding of antibodies within the sera to the immobilized polypeptide. Unbound sera may then be removed and bound antibodies detected using, for example, ¹²⁵I-labeled Protein A.

As would be recognized by the skilled artisan, immunogenic portions of the polypeptides disclosed herein are also encompassed by the present invention. An
30 "immunogenic portion," as used herein, is a fragment of an immunogenic polypeptide

of the invention that itself is immunologically reactive (*i.e.*, specifically binds) with the B-cells and/or T-cell surface antigen receptors that recognize the polypeptide. Immunogenic portions may generally be identified using well known techniques, such as those summarized in Paul, *Fundamental Immunology*, 3rd ed., 243-247 (Raven Press, 5 1993) and references cited therein. Such techniques include screening polypeptides for the ability to react with antigen-specific antibodies, antisera and/or T-cell lines or clones. As used herein, antisera and antibodies are "antigen-specific" if they specifically bind to an antigen (*i.e.*, they react with the protein in an ELISA or other immunoassay, and do not react detectably with unrelated proteins). Such antisera and 10 antibodies may be prepared as described herein, and using well-known techniques.

In one preferred embodiment, an immunogenic portion of a polypeptide of the present invention is a portion that reacts with antisera and/or T-cells at a level that is not substantially less than the reactivity of the full-length polypeptide (*e.g.*, in an ELISA and/or T-cell reactivity assay). Preferably, the level of immunogenic activity of 15 the immunogenic portion is at least about 50%, preferably at least about 70% and most preferably greater than about 90% of the immunogenicity for the full-length polypeptide. In some instances, preferred immunogenic portions will be identified that have a level of immunogenic activity greater than that of the corresponding full-length polypeptide, *e.g.*, having greater than about 100% or 150% or more immunogenic 20 activity.

In certain other embodiments, illustrative immunogenic portions may include peptides in which an N-terminal leader sequence and/or transmembrane domain have been deleted. Other illustrative immunogenic portions will contain a small N- and/or C-terminal deletion (*e.g.*, 1-30 amino acids, preferably 5-15 amino acids), 25 relative to the mature protein.

In another embodiment, a polypeptide composition of the invention may also comprise one or more polypeptides that are immunologically reactive with T cells and/or antibodies generated against a polypeptide of the invention, particularly a polypeptide having an amino acid sequence disclosed herein, or to an immunogenic 30 fragment or variant thereof.

In another embodiment of the invention, polypeptides are provided that comprise one or more polypeptides that are capable of eliciting T cells and/or antibodies that are immunologically reactive with one or more polypeptides described herein, or one or more polypeptides encoded by contiguous nucleic acid sequences contained in the polynucleotide sequences disclosed herein, or immunogenic fragments or variants thereof, or to one or more nucleic acid sequences which hybridize to one or more of these sequences under conditions of moderate to high stringency.

The present invention, in another aspect, provides polypeptide fragments comprising at least about 5, 10, 15, 20, 25, 50, or 100 contiguous amino acids, or more, including all intermediate lengths, of a polypeptide compositions set forth herein, such as those set forth in SEQ ID NOs: 299, 300, 304-306, 308-312, 314, 326 and 331-334, or those encoded by a polynucleotide sequence set forth in a sequence of SEQ ID NOs: 1, 3-86, 142-298, 301-303, 307, 313, 314, 316, 317, 325 and 327-330.

In another aspect, the present invention provides variants of the polypeptide compositions described herein. Polypeptide variants generally encompassed by the present invention will typically exhibit at least about 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% or more identity (determined as described below), along its length, to a polypeptide sequences set forth herein.

In one preferred embodiment, the polypeptide fragments and variants provide by the present invention are immunologically reactive with an antibody and/or T-cell that reacts with a full-length polypeptide specifically set for the herein.

In another preferred embodiment, the polypeptide fragments and variants provided by the present invention exhibit a level of immunogenic activity of at least about 50%, preferably at least about 70%, and most preferably at least about 90% or more of that exhibited by a full-length polypeptide sequence specifically set forth herein.

A polypeptide "variant," as the term is used herein, is a polypeptide that typically differs from a polypeptide specifically disclosed herein in one or more substitutions, deletions, additions and/or insertions. Such variants may be naturally

occurring or may be synthetically generated, for example, by modifying one or more of the above polypeptide sequences of the invention and evaluating their immunogenic activity as described herein and/or using any of a number of techniques well known in the art.

5 For example, certain illustrative variants of the polypeptides of the invention include those in which one or more portions, such as an N-terminal leader sequence or transmembrane domain, have been removed. Other illustrative variants include variants in which a small portion (*e.g.*, 1-30 amino acids, preferably 5-15 amino acids) has been removed from the N- and/or C-terminal of the mature protein.

10 In many instances, a variant will contain conservative substitutions. A "conservative substitution" is one in which an amino acid is substituted for another amino acid that has similar properties, such that one skilled in the art of peptide chemistry would expect the secondary structure and hydrophobic nature of the polypeptide to be substantially unchanged. As described above, modifications may be
15 made in the structure of the polynucleotides and polypeptides of the present invention and still obtain a functional molecule that encodes a variant or derivative polypeptide with desirable characteristics, *e.g.*, with immunogenic characteristics. When it is desired to alter the amino acid sequence of a polypeptide to create an equivalent, or even an improved, immunogenic variant or portion of a polypeptide of the invention,
20 one skilled in the art will typically change one or more of the codons of the encoding DNA sequence according to Table 1.

 For example, certain amino acids may be substituted for other amino acids in a protein structure without appreciable loss of interactive binding capacity with structures such as, for example, antigen-binding regions of antibodies or binding sites
25 on substrate molecules. Since it is the interactive capacity and nature of a protein that defines that protein's biological functional activity, certain amino acid sequence substitutions can be made in a protein sequence, and, of course, its underlying DNA coding sequence, and nevertheless obtain a protein with like properties. It is thus contemplated that various changes may be made in the peptide sequences of the

disclosed compositions, or corresponding DNA sequences which encode said peptides without appreciable loss of their biological utility or activity.

TABLE 1

5

Amino Acids			Codons						
Alanine	Ala	A	GCA	GCC	GCG	GCU			
Cysteine	Cys	C	UGC	UGU					
Aspartic acid	Asp	D	GAC	GAU					
Glutamic acid	Glu	E	GAA	GAG					
Phenylalanine	Phe	F	UUC	UUU					
Glycine	Gly	G	GGA	GGC	GGG	GGU			
Histidine	His	H	CAC	CAU					
Isoleucine	Ile	I	AUA	AUC	AUU				
Lysine	Lys	K	AAA	AAG					
Leucine	Leu	L	UUA	UUG	CUA	CUC	CUG	CUU	
Methionine	Met	M	AUG						
Asparagine	Asn	N	AAC	AAU					
Proline	Pro	P	CCA	CCC	CCG	CCU			
Glutamine	Gln	Q	CAA	CAG					
Arginine	Arg	R	AGA	AGG	CGA	CGC	CGG	CGU	
Serine	Ser	S	AGC	AGU	UCA	UCC	UCG	UCU	
Threonine	Thr	T	ACA	ACC	ACG	ACU			
Valine	Val	V	GUA	GUC	GUG	GUU			
Tryptophan	Trp	W	UGG						
Tyrosine	Tyr	Y	UAC	UAU					

In making such changes, the hydropathic index of amino acids may be considered. The importance of the hydropathic amino acid index in conferring interactive biologic function on a protein is generally understood in the art (Kyte and Doolittle, 1982, incorporated herein by reference). It is accepted that the relative hydropathic character of the amino acid contributes to the secondary structure of the

resultant protein, which in turn defines the interaction of the protein with other molecules, for example, enzymes, substrates, receptors, DNA, antibodies, antigens, and the like. Each amino acid has been assigned a hydropathic index on the basis of its hydrophobicity and charge characteristics (Kyte and Doolittle, 1982). These values are:

5 isoleucine (+4.5); valine (+4.2); leucine (+3.8); phenylalanine (+2.8); cysteine/cystine (+2.5); methionine (+1.9); alanine (+1.8); glycine (−0.4); threonine (−0.7); serine (−0.8); tryptophan (−0.9); tyrosine (−1.3); proline (−1.6); histidine (−3.2); glutamate (−3.5); glutamine (−3.5); aspartate (−3.5); asparagine (−3.5); lysine (−3.9); and arginine (−4.5).

It is known in the art that certain amino acids may be substituted by other

10 amino acids having a similar hydropathic index or score and still result in a protein with similar biological activity, *i.e.* still obtain a biological functionally equivalent protein. In making such changes, the substitution of amino acids whose hydropathic indices are within ± 2 is preferred, those within ± 1 are particularly preferred, and those within ± 0.5 are even more particularly preferred. It is also understood in the art that the substitution

15 of like amino acids can be made effectively on the basis of hydrophilicity. U. S. Patent 4,554,101 (specifically incorporated herein by reference in its entirety), states that the greatest local average hydrophilicity of a protein, as governed by the hydrophilicity of its adjacent amino acids, correlates with a biological property of the protein.

As detailed in U. S. Patent 4,554,101, the following hydrophilicity values

20 have been assigned to amino acid residues: arginine (+3.0); lysine (+3.0); aspartate (+3.0 \pm 1); glutamate (+3.0 \pm 1); serine (+0.3); asparagine (+0.2); glutamine (+0.2); glycine (0); threonine (−0.4); proline (−0.5 \pm 1); alanine (−0.5); histidine (−0.5); cysteine (−1.0); methionine (−1.3); valine (−1.5); leucine (−1.8); isoleucine (−1.8); tyrosine (−2.3); phenylalanine (−2.5); tryptophan (−3.4). It is understood that an amino acid can be

25 substituted for another having a similar hydrophilicity value and still obtain a biologically equivalent, and in particular, an immunologically equivalent protein. In such changes, the substitution of amino acids whose hydrophilicity values are within ± 2 is preferred, those within ± 1 are particularly preferred, and those within ± 0.5 are even more particularly preferred.

As outlined above, amino acid substitutions are generally therefore based on the relative similarity of the amino acid side-chain substituents, for example, their hydrophobicity, hydrophilicity, charge, size, and the like. Exemplary substitutions that take various of the foregoing characteristics into consideration are well known to those
5 of skill in the art and include: arginine and lysine; glutamate and aspartate; serine and threonine; glutamine and asparagine; and valine, leucine and isoleucine.

In addition, any polynucleotide may be further modified to increase stability *in vivo*. Possible modifications include, but are not limited to, the addition of flanking sequences at the 5' and/or 3' ends; the use of phosphorothioate or 2' O-methyl
10 rather than phosphodiesterase linkages in the backbone; and/or the inclusion of nontraditional bases such as inosine, queosine and wybutosine, as well as acetyl-methyl-, thio- and other modified forms of adenine, cytidine, guanine, thymine and uridine.

Amino acid substitutions may further be made on the basis of similarity
15 in polarity, charge, solubility, hydrophobicity, hydrophilicity and/or the amphipathic nature of the residues. For example, negatively charged amino acids include aspartic acid and glutamic acid; positively charged amino acids include lysine and arginine; and amino acids with uncharged polar head groups having similar hydrophilicity values include leucine, isoleucine and valine; glycine and alanine; asparagine and glutamine;
20 and serine, threonine, phenylalanine and tyrosine. Other groups of amino acids that may represent conservative changes include: (1) ala, pro, gly, glu, asp, gln, asn, ser, thr; (2) cys, ser, tyr, thr; (3) val, ile, leu, met, ala, phe; (4) lys, arg, his; and (5) phe, tyr, trp, his. A variant may also, or alternatively, contain nonconservative changes. In a preferred embodiment, variant polypeptides differ from a native sequence by
25 substitution, deletion or addition of five amino acids or fewer. Variants may also (or alternatively) be modified by, for example, the deletion or addition of amino acids that have minimal influence on the immunogenicity, secondary structure and hydrophobic nature of the polypeptide.

As noted above, polypeptides may comprise a signal (or leader) sequence
30 at the N-terminal end of the protein, which co-translationally or post-translationally

directs transfer of the protein. The polypeptide may also be conjugated to a linker or other sequence for ease of synthesis, purification or identification of the polypeptide (e.g., poly-His), or to enhance binding of the polypeptide to a solid support. For example, a polypeptide may be conjugated to an immunoglobulin Fc region.

5 When comparing polypeptide sequences, two sequences are said to be “identical” if the sequence of amino acids in the two sequences is the same when aligned for maximum correspondence, as described below. Comparisons between two sequences are typically performed by comparing the sequences over a comparison window to identify and compare local regions of sequence similarity. A “comparison
10 window” as used herein, refers to a segment of at least about 20 contiguous positions, usually 30 to about 75, 40 to about 50, in which a sequence may be compared to a reference sequence of the same number of contiguous positions after the two sequences are optimally aligned.

Optimal alignment of sequences for comparison may be conducted using
15 the Megalign program in the Lasergene suite of bioinformatics software (DNASTAR, Inc., Madison, WI), using default parameters. This program embodies several alignment schemes described in the following references: Dayhoff, M.O. (1978) A model of evolutionary change in proteins – Matrices for detecting distant relationships. In Dayhoff, M.O. (ed.) *Atlas of Protein Sequence and Structure*, National Biomedical
20 Research Foundation, Washington DC Vol. 5, Suppl. 3, pp. 345-358; Hein J. (1990) Unified Approach to Alignment and Phylogenies pp. 626-645 *Methods in Enzymology* vol. 183, Academic Press, Inc., San Diego, CA; Higgins, D.G. and Sharp, P.M. (1989) *CABIOS* 5:151-153; Myers, E.W. and Muller W. (1988) *CABIOS* 4:11-17; Robinson, E.D. (1971) *Comb. Theor* 11:105; Santou, N. Nes, M. (1987) *Mol. Biol. Evol.* 4:406-
25 425; Sneath, P.H.A. and Sokal, R.R. (1973) *Numerical Taxonomy – the Principles and Practice of Numerical Taxonomy*, Freeman Press, San Francisco, CA; Wilbur, W.J. and Lipman, D.J. (1983) *Proc. Natl. Acad. Sci. USA* 80:726-730.

Alternatively, optimal alignment of sequences for comparison may be conducted by the local identity algorithm of Smith and Waterman (1981) *Add. APL*.
30 *Math* 2:482, by the identity alignment algorithm of Needleman and Wunsch (1970) *J.*

Mol. Biol. 48:443, by the search for similarity methods of Pearson and Lipman (1988) *Proc. Natl. Acad. Sci. USA* 85: 2444, by computerized implementations of these algorithms (GAP, BESTFIT, BLAST, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group (GCG), 575 Science Dr., Madison, WI),
5 or by inspection.

One preferred example of algorithms that are suitable for determining percent sequence identity and sequence similarity are the BLAST and BLAST 2.0 algorithms, which are described in Altschul et al. (1977) *Nucl. Acids Res.* 25:3389-3402 and Altschul et al. (1990) *J. Mol. Biol.* 215:403-410, respectively. BLAST and BLAST
10 2.0 can be used, for example with the parameters described herein, to determine percent sequence identity for the polynucleotides and polypeptides of the invention. Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information. For amino acid sequences, a scoring matrix can be used to calculate the cumulative score. Extension of the word hits in each direction are halted
15 when: the cumulative alignment score falls off by the quantity X from its maximum achieved value; the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm parameters W, T and X determine the sensitivity and speed of the alignment.

20 In one preferred approach, the "percentage of sequence identity" is determined by comparing two optimally aligned sequences over a window of comparison of at least 20 positions, wherein the portion of the polypeptide sequence in the comparison window may comprise additions or deletions (*i.e.*, gaps) of 20 percent or less, usually 5 to 15 percent, or 10 to 12 percent, as compared to the reference
25 sequences (which does not comprise additions or deletions) for optimal alignment of the two sequences. The percentage is calculated by determining the number of positions at which the identical amino acid residue occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the reference sequence (*i.e.*, the window size) and multiplying the results by
30 100 to yield the percentage of sequence identity.

Within other illustrative embodiments, a polypeptide may be a fusion polypeptide that comprises multiple polypeptides as described herein, or that comprises at least one polypeptide as described herein and an unrelated sequence, such as a known tumor protein. A fusion partner may, for example, assist in providing T helper epitopes (an immunological fusion partner), preferably T helper epitopes recognized by humans, or may assist in expressing the protein (an expression enhancer) at higher yields than the native recombinant protein. Certain preferred fusion partners are both immunological and expression enhancing fusion partners. Other fusion partners may be selected so as to increase the solubility of the polypeptide or to enable the polypeptide to be targeted to desired intracellular compartments. Still further fusion partners include affinity tags, which facilitate purification of the polypeptide.

Fusion polypeptides may generally be prepared using standard techniques, including chemical conjugation. Preferably, a fusion polypeptide is expressed as a recombinant polypeptide, allowing the production of increased levels, relative to a non-fused polypeptide, in an expression system. Briefly, DNA sequences encoding the polypeptide components may be assembled separately, and ligated into an appropriate expression vector. The 3' end of the DNA sequence encoding one polypeptide component is ligated, with or without a peptide linker, to the 5' end of a DNA sequence encoding the second polypeptide component so that the reading frames of the sequences are in phase. This permits translation into a single fusion polypeptide that retains the biological activity of both component polypeptides.

A peptide linker sequence may be employed to separate the first and second polypeptide components by a distance sufficient to ensure that each polypeptide folds into its secondary and tertiary structures. Such a peptide linker sequence is incorporated into the fusion polypeptide using standard techniques well known in the art. Suitable peptide linker sequences may be chosen based on the following factors: (1) their ability to adopt a flexible extended conformation; (2) their inability to adopt a secondary structure that could interact with functional epitopes on the first and second polypeptides; and (3) the lack of hydrophobic or charged residues that might react with the polypeptide functional epitopes. Preferred peptide linker sequences contain Gly,

Asn and Ser residues. Other near neutral amino acids, such as Thr and Ala may also be used in the linker sequence. Amino acid sequences which may be usefully employed as linkers include those disclosed in Maratea et al., *Gene* 40:39-46, 1985; Murphy et al., *Proc. Natl. Acad. Sci. USA* 83:8258-8262, 1986; U.S. Patent No. 4,935,233 and U.S. Patent No. 4,751,180. The linker sequence may generally be from 1 to about 50 amino acids in length. Linker sequences are not required when the first and second polypeptides have non-essential N-terminal amino acid regions that can be used to separate the functional domains and prevent steric interference.

The ligated DNA sequences are operably linked to suitable transcriptional or translational regulatory elements. The regulatory elements responsible for expression of DNA are located only 5' to the DNA sequence encoding the first polypeptides. Similarly, stop codons required to end translation and transcription termination signals are only present 3' to the DNA sequence encoding the second polypeptide.

The fusion polypeptide can comprise a polypeptide as described herein together with an unrelated immunogenic protein, such as an immunogenic protein capable of eliciting a recall response. Examples of such proteins include tetanus, tuberculosis and hepatitis proteins (*see*, for example, Stoute et al. *New Engl. J. Med.*, 336:86-91, 1997).

In one preferred embodiment, the immunological fusion partner is derived from a *Mycobacterium* sp., such as a *Mycobacterium tuberculosis*-derived Ra12 fragment. Ra12 compositions and methods for their use in enhancing the expression and/or immunogenicity of heterologous polynucleotide/polypeptide sequences is described in U.S. Patent Application 60/158,585, the disclosure of which is incorporated herein by reference in its entirety. Briefly, Ra12 refers to a polynucleotide region that is a subsequence of a *Mycobacterium tuberculosis* MTB32A nucleic acid. MTB32A is a serine protease of 32 KD molecular weight encoded by a gene in virulent and avirulent strains of *M. tuberculosis*. The nucleotide sequence and amino acid sequence of MTB32A have been described (for example, U.S. Patent Application 60/158,585; *see also*, Skeiky et al., *Infection and Immun.* (1999) 67:3998-4007,

incorporated herein by reference). C-terminal fragments of the MTB32A coding sequence express at high levels and remain as soluble polypeptides throughout the purification process. Moreover, Ra12 may enhance the immunogenicity of heterologous immunogenic polypeptides with which it is fused. One preferred Ra12 fusion polypeptide comprises a 14 KD C-terminal fragment corresponding to amino acid residues 192 to 323 of MTB32A. Other preferred Ra12 polynucleotides generally comprise at least about 15 consecutive nucleotides, at least about 30 nucleotides, at least about 60 nucleotides, at least about 100 nucleotides, at least about 200 nucleotides, or at least about 300 nucleotides that encode a portion of a Ra12 polypeptide. Ra12 polynucleotides may comprise a native sequence (*i.e.*, an endogenous sequence that encodes a Ra12 polypeptide or a portion thereof) or may comprise a variant of such a sequence. Ra12 polynucleotide variants may contain one or more substitutions, additions, deletions and/or insertions such that the biological activity of the encoded fusion polypeptide is not substantially diminished, relative to a fusion polypeptide comprising a native Ra12 polypeptide. Variants preferably exhibit at least about 70% identity, more preferably at least about 80% identity and most preferably at least about 90% identity to a polynucleotide sequence that encodes a native Ra12 polypeptide or a portion thereof.

Within other preferred embodiments, an immunological fusion partner is derived from protein D, a surface protein of the gram-negative bacterium *Haemophilus influenza B* (WO 91/18926). Preferably, a protein D derivative comprises approximately the first third of the protein (*e.g.*, the first N-terminal 100-110 amino acids), and a protein D derivative may be lipidated. Within certain preferred embodiments, the first 109 residues of a Lipoprotein D fusion partner is included on the N-terminus to provide the polypeptide with additional exogenous T-cell epitopes and to increase the expression level in *E. coli* (thus functioning as an expression enhancer). The lipid tail ensures optimal presentation of the antigen to antigen presenting cells. Other fusion partners include the non-structural protein from influenzae virus, NS1 (hemagglutinin). Typically, the N-terminal 81 amino acids are used, although different fragments that include T-helper epitopes may be used.

In another embodiment, the immunological fusion partner is the protein known as LYTA, or a portion thereof (preferably a C-terminal portion). LYTA is derived from *Streptococcus pneumoniae*, which synthesizes an N-acetyl-L-alanine amidase known as amidase LYTA (encoded by the *LytA* gene; *Gene* 43:265-292, 1986).

5 LYTA is an autolysin that specifically degrades certain bonds in the peptidoglycan backbone. The C-terminal domain of the LYTA protein is responsible for the affinity to the choline or to some choline analogues such as DEAE. This property has been exploited for the development of *E. coli* C-LYTA expressing plasmids useful for expression of fusion proteins. Purification of hybrid proteins containing the C-LYTA
10 fragment at the amino terminus has been described (*see Biotechnology* 10:795-798, 1992). Within a preferred embodiment, a repeat portion of LYTA may be incorporated into a fusion polypeptide. A repeat portion is found in the C-terminal region starting at residue 178. A particularly preferred repeat portion incorporates residues 188-305.

Yet another illustrative embodiment involves fusion polypeptides, and
15 the polynucleotides encoding them, wherein the fusion partner comprises a targeting signal capable of directing a polypeptide to the endosomal/lysosomal compartment, as described in U.S. Patent No. 5,633,234. An immunogenic polypeptide of the invention, when fused with this targeting signal, will associate more efficiently with MHC class II molecules and thereby provide enhanced in vivo stimulation of CD4⁺ T-cells specific
20 for the polypeptide.

Polypeptides of the invention are prepared using any of a variety of well known synthetic and/or recombinant techniques, the latter of which are further described below. Polypeptides, portions and other variants generally less than about 150 amino acids can be generated by synthetic means, using techniques well known to
25 those of ordinary skill in the art. In one illustrative example, such polypeptides are synthesized using any of the commercially available solid-phase techniques, such as the Merrifield solid-phase synthesis method, where amino acids are sequentially added to a growing amino acid chain. *See Merrifield, J. Am. Chem. Soc.* 85:2149-2146, 1963. Equipment for automated synthesis of polypeptides is commercially available from

suppliers such as Perkin Elmer/Applied BioSystems Division (Foster City, CA), and may be operated according to the manufacturer's instructions.

In general, polypeptide compositions (including fusion polypeptides) of the invention are isolated. An "isolated" polypeptide is one that is removed from its original environment. For example, a naturally-occurring protein or polypeptide is isolated if it is separated from some or all of the coexisting materials in the natural system. Preferably, such polypeptides are also purified, *e.g.*, are at least about 90% pure, more preferably at least about 95% pure and most preferably at least about 99% pure.

10 Polynucleotide Compositions

The present invention, in other aspects, provides polynucleotide compositions. The terms "DNA" and "polynucleotide" are used essentially interchangeably herein to refer to a DNA molecule that has been isolated free of total genomic DNA of a particular species. "Isolated," as used herein, means that a polynucleotide is substantially away from other coding sequences, and that the DNA molecule does not contain large portions of unrelated coding DNA, such as large chromosomal fragments or other functional genes or polypeptide coding regions. Of course, this refers to the DNA molecule as originally isolated, and does not exclude genes or coding regions later added to the segment by the hand of man.

20 As will be understood by those skilled in the art, the polynucleotide compositions of this invention can include genomic sequences, extra-genomic and plasmid-encoded sequences and smaller engineered gene segments that express, or may be adapted to express, proteins, polypeptides, peptides and the like. Such segments may be naturally isolated, or modified synthetically by the hand of man.

25 As will be also recognized by the skilled artisan, polynucleotides of the invention may be single-stranded (coding or antisense) or double-stranded, and may be DNA (genomic, cDNA or synthetic) or RNA molecules. RNA molecules may include HnRNA molecules, which contain introns and correspond to a DNA molecule in a one-to-one manner, and mRNA molecules, which do not contain introns. Additional coding

or non-coding sequences may, but need not, be present within a polynucleotide of the present invention, and a polynucleotide may, but need not, be linked to other molecules and/or support materials.

Polynucleotides may comprise a native sequence (*i.e.*, an endogenous
5 sequence that encodes a polypeptide/protein of the invention or a portion thereof) or may comprise a sequence that encodes a variant or derivative, preferably and immunogenic variant or derivative, of such a sequence.

Therefore, according to another aspect of the present invention, polynucleotide compositions are provided that comprise some or all of a polynucleotide
10 sequence set forth in any one of SEQ ID NOs: 1, 3-86, 142-298, 301-303, 307, 313, 314, 316, 317, 325 and 327-330, complements of a polynucleotide sequence set forth in any one of SEQ ID NOs: 1, 3-86, 142-298, 301-303, 307, 313, 314, 316, 317, 325 and 327-330, and degenerate variants of a polynucleotide sequence set forth in any one of SEQ ID NOs: 1, 3-86, 142-298, 301-303, 307, 313, 314, 316, 317, 325 and 327-330. In
15 certain preferred embodiments, the polynucleotide sequences set forth herein encode immunogenic polypeptides, as described above.

In other related embodiments, the present invention provides polynucleotide variants having substantial identity to the sequences disclosed herein in SEQ ID NOs: 1, 3-86, 142-298, 301-303, 307, 313, 314, 316, 317, 325 and 327-330, for
20 example those comprising at least 70% sequence identity, preferably at least 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% or higher, sequence identity compared to a polynucleotide sequence of this invention using the methods described herein, (*e.g.*, BLAST analysis using standard parameters, as described below). One skilled in this art will recognize that these values can be appropriately adjusted to determine
25 corresponding identity of proteins encoded by two nucleotide sequences by taking into account codon degeneracy, amino acid similarity, reading frame positioning and the like.

Typically, polynucleotide variants will contain one or more substitutions, additions, deletions and/or insertions, preferably such that the immunogenicity of the
30 polypeptide encoded by the variant polynucleotide is not substantially diminished

relative to a polypeptide encoded by a polynucleotide sequence specifically set forth herein). The term "variants" should also be understood to encompass homologous genes of xenogenic origin.

In additional embodiments, the present invention provides
5 polynucleotide fragments comprising various lengths of contiguous stretches of sequence identical to or complementary to one or more of the sequences disclosed herein. For example, polynucleotides are provided by this invention that comprise at least about 10, 15, 20, 30, 40, 50, 75, 100, 150, 200, 300, 400, 500 or 1000 or more contiguous nucleotides of one or more of the sequences disclosed herein as well as all
10 intermediate lengths there between. It will be readily understood that "intermediate lengths", in this context, means any length between the quoted values, such as 16, 17, 18, 19, *etc.*; 21, 22, 23, *etc.*; 30, 31, 32, *etc.*; 50, 51, 52, 53, *etc.*; 100, 101, 102, 103, *etc.*; 150, 151, 152, 153, *etc.*; including all integers through 200-500; 500-1,000, and the like.

15 In another embodiment of the invention, polynucleotide compositions are provided that are capable of hybridizing under moderate to high stringency conditions to a polynucleotide sequence provided herein, or a fragment thereof, or a complementary sequence thereof. Hybridization techniques are well known in the art of molecular biology. For purposes of illustration, suitable moderately stringent conditions for
20 testing the hybridization of a polynucleotide of this invention with other polynucleotides include prewashing in a solution of 5 X SSC, 0.5% SDS, 1.0 mM EDTA (pH 8.0); hybridizing at 50°C-60°C, 5 X SSC, overnight; followed by washing twice at 65°C for 20 minutes with each of 2X, 0.5X and 0.2X SSC containing 0.1% SDS. One skilled in the art will understand that the stringency of hybridization can be readily manipulated,
25 such as by altering the salt content of the hybridization solution and/or the temperature at which the hybridization is performed. For example, in another embodiment, suitable highly stringent hybridization conditions include those described above, with the exception that the temperature of hybridization is increased, *e.g.*, to 60-65°C or 65-70°C.

In certain preferred embodiments, the polynucleotides described above, *e.g.*, polynucleotide variants, fragments and hybridizing sequences, encode polypeptides that are immunologically cross-reactive with a polypeptide sequence specifically set forth herein. In other preferred embodiments, such polynucleotides encode
5 polypeptides that have a level of immunogenic activity of at least about 50%, preferably at least about 70%, and more preferably at least about 90% of that for a polypeptide sequence specifically set forth herein.

The polynucleotides of the present invention, or fragments thereof, regardless of the length of the coding sequence itself, may be combined with other DNA
10 sequences, such as promoters, polyadenylation signals, additional restriction enzyme sites, multiple cloning sites, other coding segments, and the like, such that their overall length may vary considerably. It is therefore contemplated that a nucleic acid fragment of almost any length may be employed, with the total length preferably being limited by the ease of preparation and use in the intended recombinant DNA protocol. For
15 example, illustrative polynucleotide segments with total lengths of about 10,000, about 5000, about 3000, about 2,000, about 1,000, about 500, about 200, about 100, about 50 base pairs in length, and the like, (including all intermediate lengths) are contemplated to be useful in many implementations of this invention.

When comparing polynucleotide sequences, two sequences are said to be
20 "identical" if the sequence of nucleotides in the two sequences is the same when aligned for maximum correspondence, as described below. Comparisons between two sequences are typically performed by comparing the sequences over a comparison window to identify and compare local regions of sequence similarity. A "comparison window" as used herein, refers to a segment of at least about 20 contiguous positions,
25 usually 30 to about 75, 40 to about 50, in which a sequence may be compared to a reference sequence of the same number of contiguous positions after the two sequences are optimally aligned.

Optimal alignment of sequences for comparison may be conducted using the Megalign program in the Lasergene suite of bioinformatics software (DNASTAR,
30 Inc., Madison, WI), using default parameters. This program embodies several

- alignment schemes described in the following references: Dayhoff, M.O. (1978) A model of evolutionary change in proteins – Matrices for detecting distant relationships. In Dayhoff, M.O. (ed.) *Atlas of Protein Sequence and Structure*, National Biomedical Research Foundation, Washington DC Vol. 5, Suppl. 3, pp. 345-358; Hein J. (1990) Unified Approach to Alignment and Phylogenesis pp. 626-645 *Methods in Enzymology* vol. 183, Academic Press, Inc., San Diego, CA; Higgins, D.G. and Sharp, P.M. (1989) *CABIOS* 5:151-153; Myers, E.W. and Muller W. (1988) *CABIOS* 4:11-17; Robinson, E.D. (1971) *Comb. Theor* 11:105; Santou, N. Nes, M. (1987) *Mol. Biol. Evol.* 4:406-425; Sneath, P.H.A. and Sokal, R.R. (1973) *Numerical Taxonomy – the Principles and Practice of Numerical Taxonomy*, Freeman Press, San Francisco, CA; Wilbur, W.J. and Lipman, D.J. (1983) *Proc. Natl. Acad. Sci. USA* 80:726-730.

Alternatively, optimal alignment of sequences for comparison may be conducted by the local identity algorithm of Smith and Waterman (1981) *Add. APL. Math* 2:482, by the identity alignment algorithm of Needleman and Wunsch (1970) *J. Mol. Biol.* 48:443, by the search for similarity methods of Pearson and Lipman (1988) *Proc. Natl. Acad. Sci. USA* 85: 2444, by computerized implementations of these algorithms (GAP, BESTFIT, BLAST, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group (GCG), 575 Science Dr., Madison, WI), or by inspection.

- One preferred example of algorithms that are suitable for determining percent sequence identity and sequence similarity are the BLAST and BLAST 2.0 algorithms, which are described in Altschul et al. (1977) *Nucl. Acids Res.* 25:3389-3402 and Altschul et al. (1990) *J. Mol. Biol.* 215:403-410, respectively. BLAST and BLAST 2.0 can be used, for example with the parameters described herein, to determine percent sequence identity for the polynucleotides of the invention. Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information. In one illustrative example, cumulative scores can be calculated using, for nucleotide sequences, the parameters M (reward score for a pair of matching residues; always >0) and N (penalty score for mismatching residues; always <0). Extension of the word hits in each direction are halted when: the cumulative alignment score falls off

by the quantity X from its maximum achieved value; the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm parameters W, T and X determine the sensitivity and speed of the alignment. The BLASTN program (for
5 nucleotide sequences) uses as defaults a wordlength (W) of 11, and expectation (E) of 10, and the BLOSUM62 scoring matrix (see Henikoff and Henikoff (1989) *Proc. Natl. Acad. Sci. USA* 89:10915) alignments, (B) of 50, expectation (E) of 10, M=5, N=-4 and a comparison of both strands.

Preferably, the "percentage of sequence identity" is determined by
10 comparing two optimally aligned sequences over a window of comparison of at least 20 positions, wherein the portion of the polynucleotide sequence in the comparison window may comprise additions or deletions (*i.e.*, gaps) of 20 percent or less, usually 5 to 15 percent, or 10 to 12 percent, as compared to the reference sequences (which does not comprise additions or deletions) for optimal alignment of the two sequences. The
15 percentage is calculated by determining the number of positions at which the identical nucleic acid bases occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the reference sequence (*i.e.*, the window size) and multiplying the results by 100 to yield the percentage of sequence identity.

20 It will be appreciated by those of ordinary skill in the art that, as a result of the degeneracy of the genetic code, there are many nucleotide sequences that encode a polypeptide as described herein. Some of these polynucleotides bear minimal homology to the nucleotide sequence of any native gene. Nonetheless, polynucleotides that vary due to differences in codon usage are specifically contemplated by the present
25 invention. Further, alleles of the genes comprising the polynucleotide sequences provided herein are within the scope of the present invention. Alleles are endogenous genes that are altered as a result of one or more mutations, such as deletions, additions and/or substitutions of nucleotides. The resulting mRNA and protein may, but need not, have an altered structure or function. Alleles may be identified using standard
30 techniques (such as hybridization, amplification and/or database sequence comparison).

Therefore, in another embodiment of the invention, a mutagenesis approach, such as site-specific mutagenesis, is employed for the preparation of immunogenic variants and/or derivatives of the polypeptides described herein. By this approach, specific modifications in a polypeptide sequence can be made through
5 mutagenesis of the underlying polynucleotides that encode them. These techniques provides a straightforward approach to prepare and test sequence variants, for example, incorporating one or more of the foregoing considerations, by introducing one or more nucleotide sequence changes into the polynucleotide.

Site-specific mutagenesis allows the production of mutants through the
10 use of specific oligonucleotide sequences which encode the DNA sequence of the desired mutation, as well as a sufficient number of adjacent nucleotides, to provide a primer sequence of sufficient size and sequence complexity to form a stable duplex on both sides of the deletion junction being traversed. Mutations may be employed in a selected polynucleotide sequence to improve, alter, decrease, modify, or otherwise
15 change the properties of the polynucleotide itself, and/or alter the properties, activity, composition, stability, or primary sequence of the encoded polypeptide.

In certain embodiments of the present invention, the inventors contemplate the mutagenesis of the disclosed polynucleotide sequences to alter one or more properties of the encoded polypeptide, such as the immunogenicity of a
20 polypeptide vaccine. The techniques of site-specific mutagenesis are well-known in the art, and are widely used to create variants of both polypeptides and polynucleotides. For example, site-specific mutagenesis is often used to alter a specific portion of a DNA molecule. In such embodiments, a primer comprising typically about 14 to about 25 nucleotides or so in length is employed, with about 5 to about 10 residues on both sides
25 of the junction of the sequence being altered.

As will be appreciated by those of skill in the art, site-specific mutagenesis techniques have often employed a phage vector that exists in both a single stranded and double stranded form. Typical vectors useful in site-directed mutagenesis include vectors such as the M13 phage. These phage are readily commercially-available
30 and their use is generally well-known to those skilled in the art. Double-stranded

plasmids are also routinely employed in site directed mutagenesis that eliminates the step of transferring the gene of interest from a plasmid to a phage.

In general, site-directed mutagenesis in accordance herewith is performed by first obtaining a single-stranded vector or melting apart of two strands of a double-stranded vector that includes within its sequence a DNA sequence that encodes the desired peptide. An oligonucleotide primer bearing the desired mutated sequence is prepared, generally synthetically. This primer is then annealed with the single-stranded vector, and subjected to DNA polymerizing enzymes such as *E. coli* polymerase I Klenow fragment, in order to complete the synthesis of the mutation-bearing strand. Thus, a heteroduplex is formed wherein one strand encodes the original non-mutated sequence and the second strand bears the desired mutation. This heteroduplex vector is then used to transform appropriate cells, such as *E. coli* cells, and clones are selected which include recombinant vectors bearing the mutated sequence arrangement.

The preparation of sequence variants of the selected peptide-encoding DNA segments using site-directed mutagenesis provides a means of producing potentially useful species and is not meant to be limiting as there are other ways in which sequence variants of peptides and the DNA sequences encoding them may be obtained. For example, recombinant vectors encoding the desired peptide sequence may be treated with mutagenic agents, such as hydroxylamine, to obtain sequence variants. Specific details regarding these methods and protocols are found in the teachings of Maloy *et al.*, 1994; Segal, 1976; Prokop and Bajpai, 1991; Kuby, 1994; and Maniatis *et al.*, 1982, each incorporated herein by reference, for that purpose.

As used herein, the term "oligonucleotide directed mutagenesis procedure" refers to template-dependent processes and vector-mediated propagation which result in an increase in the concentration of a specific nucleic acid molecule relative to its initial concentration, or in an increase in the concentration of a detectable signal, such as amplification. As used herein, the term "oligonucleotide directed mutagenesis procedure" is intended to refer to a process that involves the template-dependent extension of a primer molecule. The term template dependent process refers to nucleic acid synthesis of an RNA or a DNA molecule wherein the

sequence of the newly synthesized strand of nucleic acid is dictated by the well-known rules of complementary base pairing (see, for example, Watson, 1987). Typically, vector mediated methodologies involve the introduction of the nucleic acid fragment into a DNA or RNA vector, the clonal amplification of the vector, and the recovery of the amplified nucleic acid fragment. Examples of such methodologies are provided by U. S. Patent No. 4,237,224, specifically incorporated herein by reference in its entirety.

In another approach for the production of polypeptide variants of the present invention, recursive sequence recombination, as described in U.S. Patent No. 5,837,458, may be employed. In this approach, iterative cycles of recombination and screening or selection are performed to "evolve" individual polynucleotide variants of the invention having, for example, enhanced immunogenic activity.

In other embodiments of the present invention, the polynucleotide sequences provided herein can be advantageously used as probes or primers for nucleic acid hybridization. As such, it is contemplated that nucleic acid segments that comprise a sequence region of at least about 15 nucleotide long contiguous sequence that has the same sequence as, or is complementary to, a 15 nucleotide long contiguous sequence disclosed herein will find particular utility. Longer contiguous identical or complementary sequences, *e.g.*, those of about 20, 30, 40, 50, 100, 200, 500, 1000 (including all intermediate lengths) and even up to full length sequences will also be of use in certain embodiments.

The ability of such nucleic acid probes to specifically hybridize to a sequence of interest will enable them to be of use in detecting the presence of complementary sequences in a given sample. However, other uses are also envisioned, such as the use of the sequence information for the preparation of mutant species primers, or primers for use in preparing other genetic constructions.

Polynucleotide molecules having sequence regions consisting of contiguous nucleotide stretches of 10-14, 15-20, 30, 50, or even of 100-200 nucleotides or so (including intermediate lengths as well), identical or complementary to a polynucleotide sequence disclosed herein, are particularly contemplated as hybridization probes for use in, *e.g.*, Southern and Northern blotting. This would allow a gene

product, or fragment thereof, to be analyzed, both in diverse cell types and also in various bacterial cells. The total size of fragment, as well as the size of the complementary stretch(es), will ultimately depend on the intended use or application of the particular nucleic acid segment. Smaller fragments will generally find use in hybridization embodiments, wherein the length of the contiguous complementary region may be varied, such as between about 15 and about 100 nucleotides, but larger contiguous complementarity stretches may be used, according to the length complementary sequences one wishes to detect.

The use of a hybridization probe of about 15-25 nucleotides in length allows the formation of a duplex molecule that is both stable and selective. Molecules having contiguous complementary sequences over stretches greater than 15 bases in length are generally preferred, though, in order to increase stability and selectivity of the hybrid, and thereby improve the quality and degree of specific hybrid molecules obtained. One will generally prefer to design nucleic acid molecules having complementary stretches of 15 to 25 contiguous nucleotides, or even longer where desired.

Hybridization probes may be selected from any portion of any of the sequences disclosed herein. All that is required is to review the sequences set forth herein, or to any continuous portion of the sequences, from about 15-25 nucleotides in length up to and including the full length sequence, that one wishes to utilize as a probe or primer. The choice of probe and primer sequences may be governed by various factors. For example, one may wish to employ primers from towards the termini of the total sequence.

Small polynucleotide segments or fragments may be readily prepared by, for example, directly synthesizing the fragment by chemical means, as is commonly practiced using an automated oligonucleotide synthesizer. Also, fragments may be obtained by application of nucleic acid reproduction technology, such as the PCR™ technology of U. S. Patent 4,683,202 (incorporated herein by reference), by introducing selected sequences into recombinant vectors for recombinant production, and by other

recombinant DNA techniques generally known to those of skill in the art of molecular biology.

The nucleotide sequences of the invention may be used for their ability to selectively form duplex molecules with complementary stretches of the entire gene or gene fragments of interest. Depending on the application envisioned, one will typically desire to employ varying conditions of hybridization to achieve varying degrees of selectivity of probe towards target sequence. For applications requiring high selectivity, one will typically desire to employ relatively stringent conditions to form the hybrids, *e.g.*, one will select relatively low salt and/or high temperature conditions, such as provided by a salt concentration of from about 0.02 M to about 0.15 M salt at temperatures of from about 50°C to about 70°C. Such selective conditions tolerate little, if any, mismatch between the probe and the template or target strand, and would be particularly suitable for isolating related sequences.

Of course, for some applications, for example, where one desires to prepare mutants employing a mutant primer strand hybridized to an underlying template, less stringent (reduced stringency) hybridization conditions will typically be needed in order to allow formation of the heteroduplex. In these circumstances, one may desire to employ salt conditions such as those of from about 0.15 M to about 0.9 M salt, at temperatures ranging from about 20°C to about 55°C. Cross-hybridizing species can thereby be readily identified as positively hybridizing signals with respect to control hybridizations. In any case, it is generally appreciated that conditions can be rendered more stringent by the addition of increasing amounts of formamide, which serves to destabilize the hybrid duplex in the same manner as increased temperature. Thus, hybridization conditions can be readily manipulated, and thus will generally be a method of choice depending on the desired results.

According to another embodiment of the present invention, polynucleotide compositions comprising antisense oligonucleotides are provided. Antisense oligonucleotides have been demonstrated to be effective and targeted inhibitors of protein synthesis, and, consequently, provide a therapeutic approach by which a disease can be treated by inhibiting the synthesis of proteins that contribute to

the disease. The efficacy of antisense oligonucleotides for inhibiting protein synthesis is well established. For example, the synthesis of polygalacturonase and the muscarine type 2 acetylcholine receptor are inhibited by antisense oligonucleotides directed to their respective mRNA sequences (U. S. Patent 5,739,119 and U. S. Patent 5,759,829).

- 5 Further, examples of antisense inhibition have been demonstrated with the nuclear protein cyclin, the multiple drug resistance gene (MDG1), ICAM-1, E-selectin, STK-1, striatal GABA_A receptor and human EGF (Jaskulski *et al.*, Science. 1988 Jun 10;240(4858):1544-6; Vasanthakumar and Ahmed, Cancer Commun. 1989;1(4):225-32; Peris *et al.*, Brain Res Mol Brain Res. 1998 Jun 15;57(2):310-20; U. S. Patent 10 5,801,154; U.S. Patent 5,789,573; U. S. Patent 5,718,709 and U.S. Patent 5,610,288). Antisense constructs have also been described that inhibit and can be used to treat a variety of abnormal cellular proliferations, *e.g.* cancer (U. S. Patent 5,747,470; U. S. Patent 5,591,317 and U. S. Patent 5,783,683).

- Therefore, in certain embodiments, the present invention provides
- 15 oligonucleotide sequences that comprise all, or a portion of, any sequence that is capable of specifically binding to polynucleotide sequence described herein, or a complement thereof. In one embodiment, the antisense oligonucleotides comprise DNA or derivatives thereof. In another embodiment, the oligonucleotides comprise RNA or derivatives thereof. In a third embodiment, the oligonucleotides are modified DNAs
- 20 comprising a phosphorothioated modified backbone. In a fourth embodiment, the oligonucleotide sequences comprise peptide nucleic acids or derivatives thereof. In each case, preferred compositions comprise a sequence region that is complementary, and more preferably substantially-complementary, and even more preferably, completely complementary to one or more portions of polynucleotides disclosed herein.
- 25 Selection of antisense compositions specific for a given gene sequence is based upon analysis of the chosen target sequence and determination of secondary structure, T_m , binding energy, and relative stability. Antisense compositions may be selected based upon their relative inability to form dimers, hairpins, or other secondary structures that would reduce or prohibit specific binding to the target mRNA in a host cell. Highly
- 30 preferred target regions of the mRNA, are those which are at or near the AUG

translation initiation codon, and those sequences which are substantially complementary to 5' regions of the mRNA. These secondary structure analyses and target site selection considerations can be performed, for example, using v.4 of the OLIGO primer analysis software and/or the BLASTN 2.0.5 algorithm software (Altschul *et al.*, Nucleic Acids Res. 1997, 25(17):3389-402).

The use of an antisense delivery method employing a short peptide vector, termed MPG (27 residues), is also contemplated. The MPG peptide contains a hydrophobic domain derived from the fusion sequence of HIV gp41 and a hydrophilic domain from the nuclear localization sequence of SV40 T-antigen (Morris *et al.*, Nucleic Acids Res. 1997 Jul 15;25(14):2730-6). It has been demonstrated that several molecules of the MPG peptide coat the antisense oligonucleotides and can be delivered into cultured mammalian cells in less than 1 hour with relatively high efficiency (90%). Further, the interaction with MPG strongly increases both the stability of the oligonucleotide to nuclease and the ability to cross the plasma membrane.

According to another embodiment of the invention, the polynucleotide compositions described herein are used in the design and preparation of ribozyme molecules for inhibiting expression of the tumor polypeptides and proteins of the present invention in tumor cells. Ribozymes are RNA-protein complexes that cleave nucleic acids in a site-specific fashion. Ribozymes have specific catalytic domains that possess endonuclease activity (Kim and Cech, Proc Natl Acad Sci U S A. 1987 Dec;84(24):8788-92; Forster and Symons, Cell. 1987 Apr 24;49(2):211-20). For example, a large number of ribozymes accelerate phosphoester transfer reactions with a high degree of specificity, often cleaving only one of several phosphoesters in an oligonucleotide substrate (Cech *et al.*, Cell. 1981 Dec;27(3 Pt 2):487-96; Michel and Westhof, J Mol Biol. 1990 Dec 5;216(3):585-610; Reinhold-Hurek and Shub, Nature. 1992 May 14;357(6374):173-6). This specificity has been attributed to the requirement that the substrate bind via specific base-pairing interactions to the internal guide sequence ("IGS") of the ribozyme prior to chemical reaction.

Six basic varieties of naturally-occurring enzymatic RNAs are known presently. Each can catalyze the hydrolysis of RNA phosphodiester bonds *in trans* (and

thus can cleave other RNA molecules) under physiological conditions. In general, enzymatic nucleic acids act by first binding to a target RNA. Such binding occurs through the target binding portion of a enzymatic nucleic acid which is held in close proximity to an enzymatic portion of the molecule that acts to cleave the target RNA.

- 5 Thus, the enzymatic nucleic acid first recognizes and then binds a target RNA through complementary base-pairing, and once bound to the correct site, acts enzymatically to cut the target RNA. Strategic cleavage of such a target RNA will destroy its ability to direct synthesis of an encoded protein. After an enzymatic nucleic acid has bound and cleaved its RNA target, it is released from that RNA to search for another target and can
10 repeatedly bind and cleave new targets.

- The enzymatic nature of a ribozyme is advantageous over many technologies, such as antisense technology (where a nucleic acid molecule simply binds to a nucleic acid target to block its translation) since the concentration of ribozyme necessary to affect a therapeutic treatment is lower than that of an antisense
15 oligonucleotide. This advantage reflects the ability of the ribozyme to act enzymatically. Thus, a single ribozyme molecule is able to cleave many molecules of target RNA. In addition, the ribozyme is a highly specific inhibitor, with the specificity of inhibition depending not only on the base pairing mechanism of binding to the target RNA, but also on the mechanism of target RNA cleavage. Single mismatches, or base-
20 substitutions, near the site of cleavage can completely eliminate catalytic activity of a ribozyme. Similar mismatches in antisense molecules do not prevent their action (Woolf *et al.*, Proc Natl Acad Sci U S A. 1992 Aug 15;89(16):7305-9). Thus, the specificity of action of a ribozyme is greater than that of an antisense oligonucleotide binding the same RNA site.

- 25 The enzymatic nucleic acid molecule may be formed in a hammerhead, hairpin, a hepatitis δ virus, group I intron or RNaseP RNA (in association with an RNA guide sequence) or Neurospora VS RNA motif. Examples of hammerhead motifs are described by Rossi *et al.* Nucleic Acids Res. 1992 Sep 11;20(17):4559-65. Examples of hairpin motifs are described by Hampel *et al.* (Eur. Pat. Appl. Publ. No. EP 0360257),
30 Hampel and Tritz, Biochemistry 1989 Jun 13;28(12):4929-33; Hampel *et al.*, Nucleic

Acids Res. 1990 Jan 25;18(2):299-304 and U. S. Patent 5,631,359. An example of the hepatitis δ virus motif is described by Perrotta and Been, Biochemistry. 1992 Dec 1;31(47):11843-52; an example of the RNaseP motif is described by Guerrier-Takada *et al.*, Cell. 1983 Dec;35(3 Pt 2):849-57; Neurospora VS RNA ribozyme motif is described by Collins (Saville and Collins, Cell. 1990 May 18;61(4):685-96; Saville and Collins, Proc Natl Acad Sci U S A. 1991 Oct 1;88(19):8826-30; Collins and Olive, Biochemistry. 1993 Mar 23;32(11):2795-9); and an example of the Group I intron is described in (U. S. Patent 4,987,071). All that is important in an enzymatic nucleic acid molecule of this invention is that it has a specific substrate binding site which is complementary to one or more of the target gene RNA regions, and that it have nucleotide sequences within or surrounding that substrate binding site which impart an RNA cleaving activity to the molecule. Thus the ribozyme constructs need not be limited to specific motifs mentioned herein.

Ribozymes may be designed as described in Int. Pat. Appl. Publ. No. WO 93/23569 and Int. Pat. Appl. Publ. No. WO 94/02595, each specifically incorporated herein by reference) and synthesized to be tested *in vitro* and *in vivo*, as described. Such ribozymes can also be optimized for delivery. While specific examples are provided, those in the art will recognize that equivalent RNA targets in other species can be utilized when necessary.

Ribozyme activity can be optimized by altering the length of the ribozyme binding arms, or chemically synthesizing ribozymes with modifications that prevent their degradation by serum ribonucleases (see *e.g.*, Int. Pat. Appl. Publ. No. WO 92/07065; Int. Pat. Appl. Publ. No. WO 93/15187; Int. Pat. Appl. Publ. No. WO 91/03162; Eur. Pat. Appl. Publ. No. 92110298.4; U. S. Patent 5,334,711; and Int. Pat. Appl. Publ. No. WO 94/13688, which describe various chemical modifications that can be made to the sugar moieties of enzymatic RNA molecules), modifications which enhance their efficacy in cells, and removal of stem II bases to shorten RNA synthesis times and reduce chemical requirements.

Sullivan *et al.* (Int. Pat. Appl. Publ. No. WO 94/02595) describes the general methods for delivery of enzymatic RNA molecules. Ribozymes may be

administered to cells by a variety of methods known to those familiar to the art, including, but not restricted to, encapsulation in liposomes, by iontophoresis, or by incorporation into other vehicles, such as hydrogels, cyclodextrins, biodegradable nanocapsules, and bioadhesive microspheres. For some indications, ribozymes may be
5 directly delivered *ex vivo* to cells or tissues with or without the aforementioned vehicles. Alternatively, the RNA/vehicle combination may be locally delivered by direct inhalation, by direct injection or by use of a catheter, infusion pump or stent. Other routes of delivery include, but are not limited to, intravascular, intramuscular, subcutaneous or joint injection, aerosol inhalation, oral (tablet or pill form), topical,
10 systemic, ocular, intraperitoneal and/or intrathecal delivery. More detailed descriptions of ribozyme delivery and administration are provided in Int. Pat. Appl. Publ. No. WO 94/02595 and Int. Pat. Appl. Publ. No. WO 93/23569, each specifically incorporated herein by reference.

Another means of accumulating high concentrations of a ribozyme(s)
15 within cells is to incorporate the ribozyme-encoding sequences into a DNA expression vector. Transcription of the ribozyme sequences are driven from a promoter for eukaryotic RNA polymerase I (pol I), RNA polymerase II (pol II), or RNA polymerase III (pol III). Transcripts from pol II or pol III promoters will be expressed at high levels in all cells; the levels of a given pol II promoter in a given cell type will depend on the
20 nature of the gene regulatory sequences (enhancers, silencers, *etc.*) present nearby. Prokaryotic RNA polymerase promoters may also be used, providing that the prokaryotic RNA polymerase enzyme is expressed in the appropriate cells. Ribozymes expressed from such promoters have been shown to function in mammalian cells. Such transcription units can be incorporated into a variety of vectors for introduction into
25 mammalian cells, including but not restricted to, plasmid DNA vectors, viral DNA vectors (such as adenovirus or adeno-associated vectors), or viral RNA vectors (such as retroviral, semliki forest virus, sindbis virus vectors).

In another embodiment of the invention, peptide nucleic acids (PNAs) compositions are provided. PNA is a DNA mimic in which the nucleobases are
30 attached to a pseudopeptide backbone (Good and Nielsen, Antisense Nucleic Acid Drug

Dev. 1997 7(4) 431-37). PNA is able to be utilized in a number methods that traditionally have used RNA or DNA. Often PNA sequences perform better in techniques than the corresponding RNA or DNA sequences and have utilities that are not inherent to RNA or DNA. A review of PNA including methods of making,
5 characteristics of, and methods of using, is provided by Corey (*Trends Biotechnol* 1997 Jun;15(6):224-9). As such, in certain embodiments, one may prepare PNA sequences that are complementary to one or more portions of the ACE mRNA sequence, and such PNA compositions may be used to regulate, alter, decrease, or reduce the translation of ACE-specific mRNA, and thereby alter the level of ACE activity in a host cell to which
10 such PNA compositions have been administered.

PNAs have 2-aminoethyl-glycine linkages replacing the normal phosphodiester backbone of DNA (Nielsen *et al.*, *Science* 1991 Dec 6;254(5037):1497-500; Hanvey *et al.*, *Science*. 1992 Nov 27;258(5087):1481-5; Hyrup and Nielsen, *Bioorg Med Chem*. 1996 Jan;4(1):5-23). This chemistry has three important
15 consequences: firstly, in contrast to DNA or phosphorothioate oligonucleotides, PNAs are neutral molecules; secondly, PNAs are achiral, which avoids the need to develop a stereoselective synthesis; and thirdly, PNA synthesis uses standard Boc or Fmoc protocols for solid-phase peptide synthesis, although other methods, including a modified Merrifield method, have been used.

20 PNA monomers or ready-made oligomers are commercially available from PerSeptive Biosystems (Framingham, MA). PNA syntheses by either Boc or Fmoc protocols are straightforward using manual or automated protocols (Norton *et al.*, *Bioorg Med Chem*. 1995 Apr;3(4):437-45). The manual protocol lends itself to the production of chemically modified PNAs or the simultaneous synthesis of families of
25 closely related PNAs.

As with peptide synthesis, the success of a particular PNA synthesis will depend on the properties of the chosen sequence. For example, while in theory PNAs can incorporate any combination of nucleotide bases, the presence of adjacent purines can lead to deletions of one or more residues in the product. In expectation of this
30 difficulty, it is suggested that, in producing PNAs with adjacent purines, one should

repeat the coupling of residues likely to be added inefficiently. This should be followed by the purification of PNAs by reverse-phase high-pressure liquid chromatography, providing yields and purity of product similar to those observed during the synthesis of peptides.

5 Modifications of PNAs for a given application may be accomplished by coupling amino acids during solid-phase synthesis or by attaching compounds that contain a carboxylic acid group to the exposed N-terminal amine. Alternatively, PNAs can be modified after synthesis by coupling to an introduced lysine or cysteine. The ease with which PNAs can be modified facilitates optimization for better solubility or
10 for specific functional requirements. Once synthesized, the identity of PNAs and their derivatives can be confirmed by mass spectrometry. Several studies have made and utilized modifications of PNAs (for example, Norton *et al.*, Bioorg Med Chem. 1995 Apr;3(4):437-45; Petersen *et al.*, J Pept Sci. 1995 May-Jun;1(3):175-83; Orum *et al.*, Biotechniques. 1995 Sep;19(3):472-80; Footer *et al.*, Biochemistry. 1996 Aug
15 20;35(33):10673-9; Griffith *et al.*, Nucleic Acids Res. 1995 Aug 11;23(15):3003-8; Pardridge *et al.*, Proc Natl Acad Sci U S A. 1995 Jun 6;92(12):5592-6; Boffa *et al.*, Proc Natl Acad Sci U S A. 1995 Mar 14;92(6):1901-5; Gambacorti-Passerini *et al.*, Blood. 1996 Aug 15;88(4):1411-7; Armitage *et al.*, Proc Natl Acad Sci U S A. 1997 Nov 11;94(23):12320-5; Seeger *et al.*, Biotechniques. 1997 Sep;23(3):512-7). U.S.
20 Patent No. 5,700,922 discusses PNA-DNA-PNA chimeric molecules and their uses in diagnostics, modulating protein in organisms, and treatment of conditions susceptible to therapeutics.

 Methods of characterizing the antisense binding properties of PNAs are discussed in Rose (Anal Chem. 1993 Dec 15;65(24):3545-9) and Jensen *et al.*
25 (Biochemistry. 1997 Apr 22;36(16):5072-7). Rose uses capillary gel electrophoresis to determine binding of PNAs to their complementary oligonucleotide, measuring the relative binding kinetics and stoichiometry. Similar types of measurements were made by Jensen *et al.* using BIAcore™ technology.

 Other applications of PNAs that have been described and will be
30 apparent to the skilled artisan include use in DNA strand invasion, antisense inhibition,

mutational analysis, enhancers of transcription, nucleic acid purification, isolation of transcriptionally active genes, blocking of transcription factor binding, genome cleavage, biosensors, *in situ* hybridization, and the like.

Polynucleotide Identification, Characterization and Expression

5 Polynucleotides compositions of the present invention may be identified, prepared and/or manipulated using any of a variety of well established techniques (see generally, Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratories, Cold Spring Harbor, NY, 1989, and other like references). For example, a polynucleotide may be identified by screening a microarray of cDNAs for
10 tumor-associated expression (*i.e.*, expression that is at least two fold greater in a tumor than in normal tissue, as determined using a representative assay provided herein). Such screens may be performed, for example, using the microarray technology of Affymetrix, Inc. (Santa Clara, CA) according to the manufacturer's instructions (and essentially as described by Schena et al., *Proc. Natl. Acad. Sci. USA* 93:10614-10619,
15 1996 and Heller et al., *Proc. Natl. Acad. Sci. USA* 94:2150-2155, 1997). Alternatively, polynucleotides may be amplified from cDNA prepared from cells expressing the proteins described herein, such as tumor cells.

Many template dependent processes are available to amplify a target sequences of interest present in a sample. One of the best known amplification methods
20 is the polymerase chain reaction (PCR™) which is described in detail in U.S. Patent Nos. 4,683,195, 4,683,202 and 4,800,159, each of which is incorporated herein by reference in its entirety. Briefly, in PCR™, two primer sequences are prepared which are complementary to regions on opposite complementary strands of the target sequence. An excess of deoxynucleoside triphosphates is added to a reaction mixture
25 along with a DNA polymerase (*e.g.*, *Taq* polymerase). If the target sequence is present in a sample, the primers will bind to the target and the polymerase will cause the primers to be extended along the target sequence by adding on nucleotides. By raising and lowering the temperature of the reaction mixture, the extended primers will dissociate from the target to form reaction products, excess primers will bind to the

target and to the reaction product and the process is repeated. Preferably reverse transcription and PCR[™] amplification procedure may be performed in order to quantify the amount of mRNA amplified. Polymerase chain reaction methodologies are well known in the art.

5 Any of a number of other template dependent processes, many of which are variations of the PCR[™] amplification technique, are readily known and available in the art. Illustratively, some such methods include the ligase chain reaction (referred to as LCR), described, for example, in Eur. Pat. Appl. Publ. No. 320,308 and U.S. Patent No. 4,883,750; Qbeta Replicase, described in PCT Intl. Pat. Appl. Publ. No. 10 PCT/US87/00880; Strand Displacement Amplification (SDA) and Repair Chain Reaction (RCR). Still other amplification methods are described in Great Britain Pat. Appl. No. 2 202 328, and in PCT Intl. Pat. Appl. Publ. No. PCT/US89/01025. Other nucleic acid amplification procedures include transcription-based amplification systems (TAS) (PCT Intl. Pat. Appl. Publ. No. WO 88/10315), including nucleic acid sequence 15 based amplification (NASBA) and 3SR. Eur. Pat. Appl. Publ. No. 329,822 describes a nucleic acid amplification process involving cyclically synthesizing single-stranded RNA ("ssRNA"), ssDNA, and double-stranded DNA (dsDNA). PCT Intl. Pat. Appl. Publ. No. WO 89/06700 describes a nucleic acid sequence amplification scheme based on the hybridization of a promoter/primer sequence to a target single-stranded DNA 20 ("ssDNA") followed by transcription of many RNA copies of the sequence. Other amplification methods such as "RACE" (Frohman, 1990), and "one-sided PCR" (Ohara, 1989) are also well-known to those of skill in the art.

An amplified portion of a polynucleotide of the present invention may be used to isolate a full length gene from a suitable library (e.g., a tumor cDNA library) 25 using well known techniques. Within such techniques, a library (cDNA or genomic) is screened using one or more polynucleotide probes or primers suitable for amplification. Preferably, a library is size-selected to include larger molecules. Random primed libraries may also be preferred for identifying 5' and upstream regions of genes. Genomic libraries are preferred for obtaining introns and extending 5' sequences.

For hybridization techniques, a partial sequence may be labeled (*e.g.*, by nick-translation or end-labeling with ^{32}P) using well known techniques. A bacterial or bacteriophage library is then generally screened by hybridizing filters containing denatured bacterial colonies (or lawns containing phage plaques) with the labeled probe
5 (*see* Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratories, Cold Spring Harbor, NY, 1989). Hybridizing colonies or plaques are selected and expanded, and the DNA is isolated for further analysis. cDNA clones may be analyzed to determine the amount of additional sequence by, for example, PCR using a primer from the partial sequence and a primer from the vector. Restriction maps and
10 partial sequences may be generated to identify one or more overlapping clones. The complete sequence may then be determined using standard techniques, which may involve generating a series of deletion clones. The resulting overlapping sequences can then be assembled into a single contiguous sequence. A full length cDNA molecule can be generated by ligating suitable fragments, using well known techniques.

15 Alternatively, amplification techniques, such as those described above, can be useful for obtaining a full length coding sequence from a partial cDNA sequence. One such amplification technique is inverse PCR (*see* Triglia et al., *Nucl. Acids Res.* 16:8186, 1988), which uses restriction enzymes to generate a fragment in the known region of the gene. The fragment is then circularized by intramolecular ligation and
20 used as a template for PCR with divergent primers derived from the known region. Within an alternative approach, sequences adjacent to a partial sequence may be retrieved by amplification with a primer to a linker sequence and a primer specific to a known region. The amplified sequences are typically subjected to a second round of amplification with the same linker primer and a second primer specific to the known
25 region. A variation on this procedure, which employs two primers that initiate extension in opposite directions from the known sequence, is described in WO 96/38591. Another such technique is known as "rapid amplification of cDNA ends" or RACE. This technique involves the use of an internal primer and an external primer, which hybridizes to a polyA region or vector sequence, to identify sequences that are 5'
30 and 3' of a known sequence. Additional techniques include capture PCR (Lagerstrom et

al., *PCR Methods Applic. 1*:111-19, 1991) and walking PCR (Parker et al., *Nucl. Acids. Res. 19*:3055-60, 1991). Other methods employing amplification may also be employed to obtain a full length cDNA sequence.

In certain instances, it is possible to obtain a full length cDNA sequence
5 by analysis of sequences provided in an expressed sequence tag (EST) database, such as that available from GenBank. Searches for overlapping ESTs may generally be performed using well known programs (e.g., NCBI BLAST searches), and such ESTs may be used to generate a contiguous full length sequence. Full length DNA sequences may also be obtained by analysis of genomic fragments.

10 In other embodiments of the invention, polynucleotide sequences or fragments thereof which encode polypeptides of the invention, or fusion proteins or functional equivalents thereof, may be used in recombinant DNA molecules to direct expression of a polypeptide in appropriate host cells. Due to the inherent degeneracy of the genetic code, other DNA sequences that encode substantially the same or a
15 functionally equivalent amino acid sequence may be produced and these sequences may be used to clone and express a given polypeptide.

As will be understood by those of skill in the art, it may be advantageous in some instances to produce polypeptide-encoding nucleotide sequences possessing non-naturally occurring codons. For example, codons preferred by a particular
20 prokaryotic or eukaryotic host can be selected to increase the rate of protein expression or to produce a recombinant RNA transcript having desirable properties, such as a half-life which is longer than that of a transcript generated from the naturally occurring sequence.

Moreover, the polynucleotide sequences of the present invention can be
25 engineered using methods generally known in the art in order to alter polypeptide encoding sequences for a variety of reasons, including but not limited to, alterations which modify the cloning, processing, and/or expression of the gene product. For example, DNA shuffling by random fragmentation and PCR reassembly of gene fragments and synthetic oligonucleotides may be used to engineer the nucleotide
30 sequences. In addition, site-directed mutagenesis may be used to insert new restriction

sites, alter glycosylation patterns, change codon preference, produce splice variants, or introduce mutations, and so forth.

In another embodiment of the invention, natural, modified, or recombinant nucleic acid sequences may be ligated to a heterologous sequence to
5 encode a fusion protein. For example, to screen peptide libraries for inhibitors of polypeptide activity, it may be useful to encode a chimeric protein that can be recognized by a commercially available antibody. A fusion protein may also be engineered to contain a cleavage site located between the polypeptide-encoding sequence and the heterologous protein sequence, so that the polypeptide may be cleaved
10 and purified away from the heterologous moiety.

Sequences encoding a desired polypeptide may be synthesized, in whole or in part, using chemical methods well known in the art (see Caruthers, M. H. et al. (1980) *Nucl. Acids Res. Symp. Ser.* 215-223, Horn, T. et al. (1980) *Nucl. Acids Res. Symp. Ser.* 225-232). Alternatively, the protein itself may be produced using chemical
15 methods to synthesize the amino acid sequence of a polypeptide, or a portion thereof. For example, peptide synthesis can be performed using various solid-phase techniques (Roberge, J. Y. et al. (1995) *Science* 269:202-204) and automated synthesis may be achieved, for example, using the ABI 431A Peptide Synthesizer (Perkin Elmer, Palo Alto, CA).

20 A newly synthesized peptide may be substantially purified by preparative high performance liquid chromatography (*e.g.*, Creighton, T. (1983) *Proteins, Structures and Molecular Principles*, WH Freeman and Co., New York, N.Y.) or other comparable techniques available in the art. The composition of the synthetic peptides may be confirmed by amino acid analysis or sequencing (*e.g.*, the Edman degradation
25 procedure). Additionally, the amino acid sequence of a polypeptide, or any part thereof, may be altered during direct synthesis and/or combined using chemical methods with sequences from other proteins, or any part thereof, to produce a variant polypeptide.

In order to express a desired polypeptide, the nucleotide sequences encoding the polypeptide, or functional equivalents, may be inserted into appropriate
30 expression vector, *i.e.*, a vector which contains the necessary elements for the

transcription and translation of the inserted coding sequence. Methods which are well known to those skilled in the art may be used to construct expression vectors containing sequences encoding a polypeptide of interest and appropriate transcriptional and translational control elements. These methods include *in vitro* recombinant DNA
5 techniques, synthetic techniques, and *in vivo* genetic recombination. Such techniques are described, for example, in Sambrook, J. et al. (1989) Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Press, Plainview, N.Y., and Ausubel, F. M. et al. (1989) Current Protocols in Molecular Biology, John Wiley & Sons, New York. N.Y.

10 A variety of expression vector/host systems may be utilized to contain and express polynucleotide sequences. These include, but are not limited to, microorganisms such as bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors; insect cell systems infected with virus expression vectors (*e.g.*, baculovirus); plant cell
15 systems transformed with virus expression vectors (*e.g.*, cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or with bacterial expression vectors (*e.g.*, Ti or pBR322 plasmids); or animal cell systems.

The "control elements" or "regulatory sequences" present in an expression vector are those non-translated regions of the vector--enhancers, promoters,
20 5' and 3' untranslated regions--which interact with host cellular proteins to carry out transcription and translation. Such elements may vary in their strength and specificity. Depending on the vector system and host utilized, any number of suitable transcription and translation elements, including constitutive and inducible promoters, may be used. For example, when cloning in bacterial systems, inducible promoters such as the hybrid
25 lacZ promoter of the PBLUESCRIPT phagemid (Stratagene, La Jolla, Calif.) or PSPT1 plasmid (Gibco BRL, Gaithersburg, MD) and the like may be used. In mammalian cell systems, promoters from mammalian genes or from mammalian viruses are generally preferred. If it is necessary to generate a cell line that contains multiple copies of the sequence encoding a polypeptide, vectors based on SV40 or EBV may be
30 advantageously used with an appropriate selectable marker.

In bacterial systems, any of a number of expression vectors may be selected depending upon the use intended for the expressed polypeptide. For example, when large quantities are needed, for example for the induction of antibodies, vectors which direct high level expression of fusion proteins that are readily purified may be used. Such vectors include, but are not limited to, the multifunctional *E. coli* cloning and expression vectors such as BLUESCRIPT (Stratagene), in which the sequence encoding the polypeptide of interest may be ligated into the vector in frame with sequences for the amino-terminal Met and the subsequent 7 residues of β -galactosidase so that a hybrid protein is produced; pIN vectors (Van Heeke, G. and S. M. Schuster (1989) *J. Biol. Chem.* 264:5503-5509); and the like. pGEX Vectors (Promega, Madison, Wis.) may also be used to express foreign polypeptides as fusion proteins with glutathione S-transferase (GST). In general, such fusion proteins are soluble and can easily be purified from lysed cells by adsorption to glutathione-agarose beads followed by elution in the presence of free glutathione. Proteins made in such systems may be designed to include heparin, thrombin, or factor XA protease cleavage sites so that the cloned polypeptide of interest can be released from the GST moiety at will.

In the yeast, *Saccharomyces cerevisiae*, a number of vectors containing constitutive or inducible promoters such as alpha factor, alcohol oxidase, and PGH may be used. For reviews, see Ausubel et al. (supra) and Grant et al. (1987) *Methods Enzymol.* 153:516-544.

In cases where plant expression vectors are used, the expression of sequences encoding polypeptides may be driven by any of a number of promoters. For example, viral promoters such as the 35S and 19S promoters of CaMV may be used alone or in combination with the omega leader sequence from TMV (Takamatsu, N. (1987) *EMBO J.* 6:307-311. Alternatively, plant promoters such as the small subunit of RUBISCO or heat shock promoters may be used (Coruzzi, G. et al. (1984) *EMBO J.* 3:1671-1680; Broglie, R. et al. (1984) *Science* 224:838-843; and Winter, J. et al. (1991) *Results Probl. Cell Differ.* 17:85-105). These constructs can be introduced into plant cells by direct DNA transformation or pathogen-mediated transfection. Such techniques

are described in a number of generally available reviews (see, for example, Hobbs, S. or Murry, L. E. in McGraw Hill Yearbook of Science and Technology (1992) McGraw Hill, New York, N.Y.; pp. 191-196).

An insect system may also be used to express a polypeptide of interest.

- 5 For example, in one such system, *Autographa californica* nuclear polyhedrosis virus (AcNPV) is used as a vector to express foreign genes in *Spodoptera frugiperda* cells or in *Trichoplusia* larvae. The sequences encoding the polypeptide may be cloned into a non-essential region of the virus, such as the polyhedrin gene, and placed under control of the polyhedrin promoter. Successful insertion of the polypeptide-encoding sequence
- 10 will render the polyhedrin gene inactive and produce recombinant virus lacking coat protein. The recombinant viruses may then be used to infect, for example, *S. frugiperda* cells or *Trichoplusia* larvae in which the polypeptide of interest may be expressed (Engelhard, E. K. et al. (1994) *Proc. Natl. Acad. Sci.* 91 :3224-3227).

- In mammalian host cells, a number of viral-based expression systems are
- 15 generally available. For example, in cases where an adenovirus is used as an expression vector, sequences encoding a polypeptide of interest may be ligated into an adenovirus transcription/translation complex consisting of the late promoter and tripartite leader sequence. Insertion in a non-essential E1 or E3 region of the viral genome may be used to obtain a viable virus which is capable of expressing the polypeptide in infected host
- 20 cells (Logan, J. and Shenk, T. (1984) *Proc. Natl. Acad. Sci.* 81:3655-3659). In addition, transcription enhancers, such as the Rous sarcoma virus (RSV) enhancer, may be used to increase expression in mammalian host cells.

- Specific initiation signals may also be used to achieve more efficient translation of sequences encoding a polypeptide of interest. Such signals include the
- 25 ATG initiation codon and adjacent sequences. In cases where sequences encoding the polypeptide, its initiation codon, and upstream sequences are inserted into the appropriate expression vector, no additional transcriptional or translational control signals may be needed. However, in cases where only coding sequence, or a portion thereof, is inserted, exogenous translational control signals including the ATG initiation
- 30 codon should be provided. Furthermore, the initiation codon should be in the correct

reading frame to ensure translation of the entire insert. Exogenous translational elements and initiation codons may be of various origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of enhancers which are appropriate for the particular cell system which is used, such as those described in the literature (Scharf, D. et al. (1994) *Results Probl. Cell Differ.* 20:125-162).

In addition, a host cell strain may be chosen for its ability to modulate the expression of the inserted sequences or to process the expressed protein in the desired fashion. Such modifications of the polypeptide include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation. Post-translational processing which cleaves a "prepro" form of the protein may also be used to facilitate correct insertion, folding and/or function. Different host cells such as CHO, COS, HeLa, MDCK, HEK293, and WI38, which have specific cellular machinery and characteristic mechanisms for such post-translational activities, may be chosen to ensure the correct modification and processing of the foreign protein.

For long-term, high-yield production of recombinant proteins, stable expression is generally preferred. For example, cell lines which stably express a polynucleotide of interest may be transformed using expression vectors which may contain viral origins of replication and/or endogenous expression elements and a selectable marker gene on the same or on a separate vector. Following the introduction of the vector, cells may be allowed to grow for 1-2 days in an enriched media before they are switched to selective media. The purpose of the selectable marker is to confer resistance to selection, and its presence allows growth and recovery of cells which successfully express the introduced sequences. Resistant clones of stably transformed cells may be proliferated using tissue culture techniques appropriate to the cell type.

Any number of selection systems may be used to recover transformed cell lines. These include, but are not limited to, the herpes simplex virus thymidine kinase (Wigler, M. et al. (1977) *Cell* 11:223-32) and adenine phosphoribosyltransferase (Lowy, I. et al. (1990) *Cell* 22:817-23) genes which can be employed in tk.sup.- or aprt.sup.- cells, respectively. Also, antimetabolite, antibiotic or herbicide resistance can be used as the basis for selection; for example, dhfr which confers resistance to

methotrexate (Wigler, M. et al. (1980) *Proc. Natl. Acad. Sci.* 77:3567-70); npt, which confers resistance to the aminoglycosides, neomycin and G-418 (Colbere-Garapin, F. et al (1981) *J. Mol. Biol.* 150:1-14); and als or pat, which confer resistance to chlorsulfuron and phosphinotricin acetyltransferase, respectively (Murry, *supra*).

5 Additional selectable genes have been described, for example, trpB, which allows cells to utilize indole in place of tryptophan, or hisD, which allows cells to utilize histinol in place of histidine (Hartman, S. C. and R. C. Mulligan (1988) *Proc. Natl. Acad. Sci.* 85:8047-51). The use of visible markers has gained popularity with such markers as anthocyanins, beta-glucuronidase and its substrate GUS, and luciferase and its substrate

10 luciferin, being widely used not only to identify transformants, but also to quantify the amount of transient or stable protein expression attributable to a specific vector system (Rhodes, C. A. et al. (1995) *Methods Mol. Biol.* 55:121-131).

Although the presence/absence of marker gene expression suggests that the gene of interest is also present, its presence and expression may need to be

15 confirmed. For example, if the sequence encoding a polypeptide is inserted within a marker gene sequence, recombinant cells containing sequences can be identified by the absence of marker gene function. Alternatively, a marker gene can be placed in tandem with a polypeptide-encoding sequence under the control of a single promoter. Expression of the marker gene in response to induction or selection usually indicates

20 expression of the tandem gene as well.

Alternatively, host cells that contain and express a desired polynucleotide sequence may be identified by a variety of procedures known to those of skill in the art. These procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridizations and protein bioassay or immunoassay techniques which include,

25 for example, membrane, solution, or chip based technologies for the detection and/or quantification of nucleic acid or protein.

A variety of protocols for detecting and measuring the expression of polynucleotide-encoded products, using either polyclonal or monoclonal antibodies specific for the product are known in the art. Examples include enzyme-linked

30 immunosorbent assay (ELISA), radioimmunoassay (RIA), and fluorescence activated

cell sorting (FACS). A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering epitopes on a given polypeptide may be preferred for some applications, but a competitive binding assay may also be employed. These and other assays are described, among other places, in Hampton, R. et al. (1990; 5 Serological Methods, a Laboratory Manual, APS Press, St Paul, Minn.) and Maddox, D. E. et al. (1983; *J. Exp. Med.* 158:1211-1216).

A wide variety of labels and conjugation techniques are known by those skilled in the art and may be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting sequences related to 10 polynucleotides include oligolabeling, nick translation, end-labeling or PCR amplification using a labeled nucleotide. Alternatively, the sequences, or any portions thereof may be cloned into a vector for the production of an mRNA probe. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes in vitro by addition of an appropriate RNA polymerase such as T7, T3, or SP6 15 and labeled nucleotides. These procedures may be conducted using a variety of commercially available kits. Suitable reporter molecules or labels, which may be used include radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents as well as substrates, cofactors, inhibitors, magnetic particles, and the like.

Host cells transformed with a polynucleotide sequence of interest may be 20 cultured under conditions suitable for the expression and recovery of the protein from cell culture. The protein produced by a recombinant cell may be secreted or contained intracellularly depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides of the invention may be designed to contain signal sequences which direct secretion of the 25 encoded polypeptide through a prokaryotic or eukaryotic cell membrane. Other recombinant constructions may be used to join sequences encoding a polypeptide of interest to nucleotide sequence encoding a polypeptide domain which will facilitate purification of soluble proteins. Such purification facilitating domains include, but are not limited to, metal chelating peptides such as histidine-tryptophan modules that allow 30 purification on immobilized metals, protein A domains that allow purification on

immobilized immunoglobulin, and the domain utilized in the FLAGS extension/affinity purification system (Immunex Corp., Seattle, Wash.). The inclusion of cleavable linker sequences such as those specific for Factor XA or enterokinase (Invitrogen, San Diego, Calif.) between the purification domain and the encoded polypeptide may be used to facilitate purification. One such expression vector provides for expression of a fusion protein containing a polypeptide of interest and a nucleic acid encoding 6 histidine residues preceding a thioredoxin or an enterokinase cleavage site. The histidine residues facilitate purification on IMIAC (immobilized metal ion affinity chromatography) as described in Porath, J. et al. (1992, *Prot. Exp. Purif.* 3:263-281) while the enterokinase cleavage site provides a means for purifying the desired polypeptide from the fusion protein. A discussion of vectors which contain fusion proteins is provided in Kroll, D. J. et al. (1993; *DNA Cell Biol.* 12:441-453).

In addition to recombinant production methods, polypeptides of the invention, and fragments thereof, may be produced by direct peptide synthesis using solid-phase techniques (Merrifield J. (1963) *J. Am. Chem. Soc.* 85:2149-2154). Protein synthesis may be performed using manual techniques or by automation. Automated synthesis may be achieved, for example, using Applied Biosystems 431A Peptide Synthesizer (Perkin Elmer). Alternatively, various fragments may be chemically synthesized separately and combined using chemical methods to produce the full length molecule.

Antibody Compositions, Fragments Thereof and Other Binding Agents

According to another aspect, the present invention further provides binding agents, such as antibodies and antigen-binding fragments thereof, that exhibit immunological binding to a tumor polypeptide disclosed herein, or to a portion, variant or derivative thereof. An antibody, or antigen-binding fragment thereof, is said to "specifically bind," "immunologically bind," and/or is "immunologically reactive" to a polypeptide of the invention if it reacts at a detectable level (within, for example, an ELISA assay) with the polypeptide, and does not react detectably with unrelated polypeptides under similar conditions.

Immunological binding, as used in this context, generally refers to the non-covalent interactions of the type which occur between an immunoglobulin molecule and an antigen for which the immunoglobulin is specific. The strength, or affinity of immunological binding interactions can be expressed in terms of the dissociation constant (K_d) of the interaction, wherein a smaller K_d represents a greater affinity. Immunological binding properties of selected polypeptides can be quantified using methods well known in the art. One such method entails measuring the rates of antigen-binding site/antigen complex formation and dissociation, wherein those rates depend on the concentrations of the complex partners, the affinity of the interaction, and on geometric parameters that equally influence the rate in both directions. Thus, both the "on rate constant" (K_{on}) and the "off rate constant" (K_{off}) can be determined by calculation of the concentrations and the actual rates of association and dissociation. The ratio of K_{off}/K_{on} enables cancellation of all parameters not related to affinity, and is thus equal to the dissociation constant K_d . See, generally, Davies et al. (1990) Annual Rev. Biochem. 59:439-473.

An "antigen-binding site," or "binding portion" of an antibody refers to the part of the immunoglobulin molecule that participates in antigen binding. The antigen binding site is formed by amino acid residues of the N-terminal variable ("V") regions of the heavy ("H") and light ("L") chains. Three highly divergent stretches within the V regions of the heavy and light chains are referred to as "hypervariable regions" which are interposed between more conserved flanking stretches known as "framework regions," or "FRs". Thus the term "FR" refers to amino acid sequences which are naturally found between and adjacent to hypervariable regions in immunoglobulins. In an antibody molecule, the three hypervariable regions of a light chain and the three hypervariable regions of a heavy chain are disposed relative to each other in three dimensional space to form an antigen-binding surface. The antigen-binding surface is complementary to the three-dimensional surface of a bound antigen, and the three hypervariable regions of each of the heavy and light chains are referred to as "complementarity-determining regions," or "CDRs."

Binding agents may be further capable of differentiating between patients with and without a cancer, such as breast cancer, using the representative assays provided herein. For example, antibodies or other binding agents that bind to a tumor protein will preferably generate a signal indicating the presence of a cancer in at least about 20% of patients with the disease, more preferably at least about 30% of patients. Alternatively, or in addition, the antibody will generate a negative signal indicating the absence of the disease in at least about 90% of individuals without the cancer. To determine whether a binding agent satisfies this requirement, biological samples (*e.g.*, blood, sera, sputum, urine and/or tumor biopsies) from patients with and without a cancer (as determined using standard clinical tests) may be assayed as described herein for the presence of polypeptides that bind to the binding agent. Preferably, a statistically significant number of samples with and without the disease will be assayed. Each binding agent should satisfy the above criteria; however, those of ordinary skill in the art will recognize that binding agents may be used in combination to improve sensitivity.

Any agent that satisfies the above requirements may be a binding agent. For example, a binding agent may be a ribosome, with or without a peptide component, an RNA molecule or a polypeptide. In a preferred embodiment, a binding agent is an antibody or an antigen-binding fragment thereof. Antibodies may be prepared by any of a variety of techniques known to those of ordinary skill in the art. *See, e.g.*, Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, 1988. In general, antibodies can be produced by cell culture techniques, including the generation of monoclonal antibodies as described herein, or via transfection of antibody genes into suitable bacterial or mammalian cell hosts, in order to allow for the production of recombinant antibodies. In one technique, an immunogen comprising the polypeptide is initially injected into any of a wide variety of mammals (*e.g.*, mice, rats, rabbits, sheep or goats). In this step, the polypeptides of this invention may serve as the immunogen without modification. Alternatively, particularly for relatively short polypeptides, a superior immune response may be elicited if the polypeptide is joined to a carrier protein, such as bovine serum albumin or keyhole limpet hemocyanin. The immunogen

is injected into the animal host, preferably according to a predetermined schedule incorporating one or more booster immunizations, and the animals are bled periodically. Polyclonal antibodies specific for the polypeptide may then be purified from such antisera by, for example, affinity chromatography using the polypeptide coupled to a
5 suitable solid support.

Monoclonal antibodies specific for an antigenic polypeptide of interest may be prepared, for example, using the technique of Kohler and Milstein, *Eur. J. Immunol.* 6:511-519, 1976, and improvements thereto. Briefly, these methods involve the preparation of immortal cell lines capable of producing antibodies having the
10 desired specificity (*i.e.*, reactivity with the polypeptide of interest). Such cell lines may be produced, for example, from spleen cells obtained from an animal immunized as described above. The spleen cells are then immortalized by, for example, fusion with a myeloma cell fusion partner, preferably one that is syngeneic with the immunized animal. A variety of fusion techniques may be employed. For example, the spleen cells
15 and myeloma cells may be combined with a nonionic detergent for a few minutes and then plated at low density on a selective medium that supports the growth of hybrid cells, but not myeloma cells. A preferred selection technique uses HAT (hypoxanthine, aminopterin, thymidine) selection. After a sufficient time, usually about 1 to 2 weeks, colonies of hybrids are observed. Single colonies are selected and their culture
20 supernatants tested for binding activity against the polypeptide. Hybridomas having high reactivity and specificity are preferred.

Monoclonal antibodies may be isolated from the supernatants of growing hybridoma colonies. In addition, various techniques may be employed to enhance the yield, such as injection of the hybridoma cell line into the peritoneal cavity of a suitable
25 vertebrate host, such as a mouse. Monoclonal antibodies may then be harvested from the ascites fluid or the blood. Contaminants may be removed from the antibodies by conventional techniques, such as chromatography, gel filtration, precipitation, and extraction. The polypeptides of this invention may be used in the purification process in, for example, an affinity chromatography step.

A number of therapeutically useful molecules are known in the art which comprise antigen-binding sites that are capable of exhibiting immunological binding properties of an antibody molecule. The proteolytic enzyme papain preferentially cleaves IgG molecules to yield several fragments, two of which (the "F(ab)" fragments) each comprise a covalent heterodimer that includes an intact antigen-binding site. The enzyme pepsin is able to cleave IgG molecules to provide several fragments, including the "F(ab')₂" fragment which comprises both antigen-binding sites. An "Fv" fragment can be produced by preferential proteolytic cleavage of an IgM, and on rare occasions IgG or IgA immunoglobulin molecule. Fv fragments are, however, more commonly derived using recombinant techniques known in the art. The Fv fragment includes a non-covalent V_H::V_L heterodimer including an antigen-binding site which retains much of the antigen recognition and binding capabilities of the native antibody molecule. Inbar et al. (1972) Proc. Nat. Acad. Sci. USA 69:2659-2662; Hochman et al. (1976) Biochem 15:2706-2710; and Ehrlich et al. (1980) Biochem 19:4091-4096.

A single chain Fv ("sFv") polypeptide is a covalently linked V_H::V_L heterodimer which is expressed from a gene fusion including V_H- and V_L-encoding genes linked by a peptide-encoding linker. Huston et al. (1988) Proc. Nat. Acad. Sci. USA 85(16):5879-5883. A number of methods have been described to discern chemical structures for converting the naturally aggregated--but chemically separated--light and heavy polypeptide chains from an antibody V region into an sFv molecule which will fold into a three dimensional structure substantially similar to the structure of an antigen-binding site. See, *e.g.*, U.S. Pat. Nos. 5,091,513 and 5,132,405, to Huston et al.; and U.S. Pat. No. 4,946,778, to Ladner et al.

Each of the above-described molecules includes a heavy chain and a light chain CDR set, respectively interposed between a heavy chain and a light chain FR set which provide support to the CDRs and define the spatial relationship of the CDRs relative to each other. As used herein, the term "CDR set" refers to the three hypervariable regions of a heavy or light chain V region. Proceeding from the N-terminus of a heavy or light chain, these regions are denoted as "CDR1," "CDR2," and "CDR3" respectively. An antigen-binding site, therefore, includes six CDRs,

comprising the CDR set from each of a heavy and a light chain V region. A polypeptide comprising a single CDR, (*e.g.*, a CDR1, CDR2 or CDR3) is referred to herein as a "molecular recognition unit." Crystallographic analysis of a number of antigen-antibody complexes has demonstrated that the amino acid residues of CDRs form extensive
5 contact with bound antigen, wherein the most extensive antigen contact is with the heavy chain CDR3. Thus, the molecular recognition units are primarily responsible for the specificity of an antigen-binding site.

As used herein, the term "FR set" refers to the four flanking amino acid sequences which frame the CDRs of a CDR set of a heavy or light chain V region.
10 Some FR residues may contact bound antigen; however, FRs are primarily responsible for folding the V region into the antigen-binding site, particularly the FR residues directly adjacent to the CDRs. Within FRs, certain amino residues and certain structural features are very highly conserved. In this regard, all V region sequences contain an internal disulfide loop of around 90 amino acid residues. When the V regions fold into a
15 binding-site, the CDRs are displayed as projecting loop motifs which form an antigen-binding surface. It is generally recognized that there are conserved structural regions of FRs which influence the folded shape of the CDR loops into certain "canonical" structures--regardless of the precise CDR amino acid sequence. Further, certain FR residues are known to participate in non-covalent interdomain contacts which stabilize
20 the interaction of the antibody heavy and light chains.

A number of "humanized" antibody molecules comprising an antigen-binding site derived from a non-human immunoglobulin have been described, including chimeric antibodies having rodent V regions and their associated CDRs fused to human constant domains (Winter et al. (1991) *Nature* 349:293-299; Lobuglio et al. (1989)
25 *Proc. Nat. Acad. Sci. USA* 86:4220-4224; Shaw et al. (1987) *J Immunol.* 138:4534-4538; and Brown et al. (1987) *Cancer Res.* 47:3577-3583), rodent CDRs grafted into a human supporting FR prior to fusion with an appropriate human antibody constant domain (Riechmann et al. (1988) *Nature* 332:323-327; Verhoeyen et al. (1988) *Science* 239:1534-1536; and Jones et al. (1986) *Nature* 321:522-525), and rodent CDRs
30 supported by recombinantly veneered rodent FRs (European Patent Publication No.

519,596, published Dec. 23, 1992). These "humanized" molecules are designed to minimize unwanted immunological response toward rodent antihuman antibody molecules which limits the duration and effectiveness of therapeutic applications of those moieties in human recipients.

5 As used herein, the terms "veneered FRs" and "recombinantly veneered FRs" refer to the selective replacement of FR residues from, *e.g.*, a rodent heavy or light chain V region, with human FR residues in order to provide a xenogeneic molecule comprising an antigen-binding site which retains substantially all of the native FR polypeptide folding structure. Veneering techniques are based on the understanding that
10 the ligand binding characteristics of an antigen-binding site are determined primarily by the structure and relative disposition of the heavy and light chain CDR sets within the antigen-binding surface. Davies et al. (1990) *Ann. Rev. Biochem.* 59:439-473. Thus, antigen binding specificity can be preserved in a humanized antibody only wherein the CDR structures, their interaction with each other, and their interaction with the rest of
15 the V region domains are carefully maintained. By using veneering techniques, exterior (*e.g.*, solvent-accessible) FR residues which are readily encountered by the immune system are selectively replaced with human residues to provide a hybrid molecule that comprises either a weakly immunogenic, or substantially non-immunogenic veneered surface.

20 The process of veneering makes use of the available sequence data for human antibody variable domains compiled by Kabat et al., in *Sequences of Proteins of Immunological Interest*, 4th ed., (U.S. Dept. of Health and Human Services, U.S. Government Printing Office, 1987), updates to the Kabat database, and other accessible U.S. and foreign databases (both nucleic acid and protein). Solvent accessibilities of V
25 region amino acids can be deduced from the known three-dimensional structure for human and murine antibody fragments. There are two general steps in veneering a murine antigen-binding site. Initially, the FRs of the variable domains of an antibody molecule of interest are compared with corresponding FR sequences of human variable domains obtained from the above-identified sources. The most homologous human V
30 regions are then compared residue by residue to corresponding murine amino acids. The

residues in the murine FR which differ from the human counterpart are replaced by the residues present in the human moiety using recombinant techniques well known in the art. Residue switching is only carried out with moieties which are at least partially exposed (solvent accessible), and care is exercised in the replacement of amino acid residues which may have a significant effect on the tertiary structure of V region domains, such as proline, glycine and charged amino acids.

In this manner, the resultant "veneered" murine antigen-binding sites are thus designed to retain the murine CDR residues, the residues substantially adjacent to the CDRs, the residues identified as buried or mostly buried (solvent inaccessible), the residues believed to participate in non-covalent (*e.g.*, electrostatic and hydrophobic) contacts between heavy and light chain domains, and the residues from conserved structural regions of the FRs which are believed to influence the "canonical" tertiary structures of the CDR loops. These design criteria are then used to prepare recombinant nucleotide sequences which combine the CDRs of both the heavy and light chain of a murine antigen-binding site into human-appearing FRs that can be used to transfect mammalian cells for the expression of recombinant human antibodies which exhibit the antigen specificity of the murine antibody molecule.

In another embodiment of the invention, monoclonal antibodies of the present invention may be coupled to one or more therapeutic agents. Suitable agents in this regard include radionuclides, differentiation inducers, drugs, toxins, and derivatives thereof. Preferred radionuclides include ^{90}Y , ^{123}I , ^{125}I , ^{131}I , ^{186}Re , ^{188}Re , ^{211}At , and ^{212}Bi . Preferred drugs include methotrexate, and pyrimidine and purine analogs. Preferred differentiation inducers include phorbol esters and butyric acid. Preferred toxins include ricin, abrin, diphtheria toxin, cholera toxin, gelonin, *Pseudomonas* exotoxin, *Shigella* toxin, and pokeweed antiviral protein.

A therapeutic agent may be coupled (*e.g.*, covalently bonded) to a suitable monoclonal antibody either directly or indirectly (*e.g.*, via a linker group). A direct reaction between an agent and an antibody is possible when each possesses a substituent capable of reacting with the other. For example, a nucleophilic group, such as an amino or sulfhydryl group, on one may be capable of reacting with a carbonyl-

containing group, such as an anhydride or an acid halide, or with an alkyl group containing a good leaving group (*e.g.*, a halide) on the other.

Alternatively, it may be desirable to couple a therapeutic agent and an antibody via a linker group. A linker group can function as a spacer to distance an antibody from an agent in order to avoid interference with binding capabilities. A linker group can also serve to increase the chemical reactivity of a substituent on an agent or an antibody, and thus increase the coupling efficiency. An increase in chemical reactivity may also facilitate the use of agents, or functional groups on agents, which otherwise would not be possible.

It will be evident to those skilled in the art that a variety of bifunctional or polyfunctional reagents, both homo- and hetero-functional (such as those described in the catalog of the Pierce Chemical Co., Rockford, IL), may be employed as the linker group. Coupling may be effected, for example, through amino groups, carboxyl groups, sulfhydryl groups or oxidized carbohydrate residues. There are numerous references describing such methodology, *e.g.*, U.S. Patent No. 4,671,958, to Rodwell et al.

Where a therapeutic agent is more potent when free from the antibody portion of the immunoconjugates of the present invention, it may be desirable to use a linker group which is cleavable during or upon internalization into a cell. A number of different cleavable linker groups have been described. The mechanisms for the intracellular release of an agent from these linker groups include cleavage by reduction of a disulfide bond (*e.g.*, U.S. Patent No. 4,489,710, to Spitler), by irradiation of a photolabile bond (*e.g.*, U.S. Patent No. 4,625,014, to Senter et al.), by hydrolysis of derivatized amino acid side chains (*e.g.*, U.S. Patent No. 4,638,045, to Kohn et al.), by serum complement-mediated hydrolysis (*e.g.*, U.S. Patent No. 4,671,958, to Rodwell et al.), and acid-catalyzed hydrolysis (*e.g.*, U.S. Patent No. 4,569,789, to Blattler et al.).

It may be desirable to couple more than one agent to an antibody. In one embodiment, multiple molecules of an agent are coupled to one antibody molecule. In another embodiment, more than one type of agent may be coupled to one antibody. Regardless of the particular embodiment, immunoconjugates with more than one agent may be prepared in a variety of ways. For example, more than one agent may be

coupled directly to an antibody molecule, or linkers that provide multiple sites for attachment can be used. Alternatively, a carrier can be used.

A carrier may bear the agents in a variety of ways, including covalent bonding either directly or via a linker group. Suitable carriers include proteins such as albumins (*e.g.*, U.S. Patent No. 4,507,234, to Kato et al.), peptides and polysaccharides such as aminodextran (*e.g.*, U.S. Patent No. 4,699,784, to Shih et al.). A carrier may also bear an agent by noncovalent bonding or by encapsulation, such as within a liposome vesicle (*e.g.*, U.S. Patent Nos. 4,429,008 and 4,873,088). Carriers specific for radionuclide agents include radiohalogenated small molecules and chelating compounds. For example, U.S. Patent No. 4,735,792 discloses representative radiohalogenated small molecules and their synthesis. A radionuclide chelate may be formed from chelating compounds that include those containing nitrogen and sulfur atoms as the donor atoms for binding the metal, or metal oxide, radionuclide. For example, U.S. Patent No. 4,673,562, to Davison et al. discloses representative chelating compounds and their synthesis.

T Cell Compositions

The present invention, in another aspect, provides T cells specific for a tumor polypeptide disclosed herein, or for a variant or derivative thereof. Such cells may generally be prepared *in vitro* or *ex vivo*, using standard procedures. For example, T cells may be isolated from bone marrow, peripheral blood, or a fraction of bone marrow or peripheral blood of a patient, using a commercially available cell separation system, such as the Isolex™ System, available from Nexell Therapeutics, Inc. (Irvine, CA; see also U.S. Patent No. 5,240,856; U.S. Patent No. 5,215,926; WO 89/06280; WO 91/16116 and WO 92/07243). Alternatively, T cells may be derived from related or unrelated humans, non-human mammals, cell lines or cultures.

T cells may be stimulated with a polypeptide, polynucleotide encoding a polypeptide and/or an antigen presenting cell (APC) that expresses such a polypeptide. Such stimulation is performed under conditions and for a time sufficient to permit the generation of T cells that are specific for the polypeptide of interest. Preferably, a tumor

polypeptide or polynucleotide of the invention is present within a delivery vehicle, such as a microsphere, to facilitate the generation of specific T cells.

T cells are considered to be specific for a polypeptide of the present invention if the T cells specifically proliferate, secrete cytokines or kill target cells coated with the polypeptide or expressing a gene encoding the polypeptide. T cell specificity may be evaluated using any of a variety of standard techniques. For example, within a chromium release assay or proliferation assay, a stimulation index of more than two fold increase in lysis and/or proliferation, compared to negative controls, indicates T cell specificity. Such assays may be performed, for example, as described in Chen et al., *Cancer Res.* 54:1065-1070, 1994. Alternatively, detection of the proliferation of T cells may be accomplished by a variety of known techniques. For example, T cell proliferation can be detected by measuring an increased rate of DNA synthesis (e.g., by pulse-labeling cultures of T cells with tritiated thymidine and measuring the amount of tritiated thymidine incorporated into DNA). Contact with a tumor polypeptide (100 ng/ml - 100 µg/ml, preferably 200 ng/ml - 25 µg/ml) for 3 - 7 days will typically result in at least a two fold increase in proliferation of the T cells. Contact as described above for 2-3 hours should result in activation of the T cells, as measured using standard cytokine assays in which a two fold increase in the level of cytokine release (e.g., TNF or IFN-γ) is indicative of T cell activation (see Coligan et al., *Current Protocols in Immunology*, vol. 1, Wiley Interscience (Greene 1998)). T cells that have been activated in response to a tumor polypeptide, polynucleotide or polypeptide-expressing APC may be CD4⁺ and/or CD8⁺. Tumor polypeptide-specific T cells may be expanded using standard techniques. Within preferred embodiments, the T cells are derived from a patient, a related donor or an unrelated donor, and are administered to the patient following stimulation and expansion.

For therapeutic purposes, CD4⁺ or CD8⁺ T cells that proliferate in response to a tumor polypeptide, polynucleotide or APC can be expanded in number either *in vitro* or *in vivo*. Proliferation of such T cells *in vitro* may be accomplished in a variety of ways. For example, the T cells can be re-exposed to a tumor polypeptide, or a short peptide corresponding to an immunogenic portion of such a polypeptide, with or

without the addition of T cell growth factors, such as interleukin-2, and/or stimulator cells that synthesize a tumor polypeptide. Alternatively, one or more T cells that proliferate in the presence of the tumor polypeptide can be expanded in number by cloning. Methods for cloning cells are well known in the art, and include limiting
5 dilution.

Pharmaceutical Compositions

In additional embodiments, the present invention concerns formulation of one or more of the polynucleotide, polypeptide, T-cell and/or antibody compositions disclosed herein in pharmaceutically-acceptable carriers for administration to a cell or
10 an animal, either alone, or in combination with one or more other modalities of therapy.

It will be understood that, if desired, a composition as disclosed herein may be administered in combination with other agents as well, such as, *e.g.*, other proteins or polypeptides or various pharmaceutically-active agents. In fact, there is virtually no limit to other components that may also be included, given that the
15 additional agents do not cause a significant adverse effect upon contact with the target cells or host tissues. The compositions may thus be delivered along with various other agents as required in the particular instance. Such compositions may be purified from host cells or other biological sources, or alternatively may be chemically synthesized as described herein. Likewise, such compositions may further comprise substituted or
20 derivatized RNA or DNA compositions.

Therefore, in another aspect of the present invention, pharmaceutical compositions are provided comprising one or more of the polynucleotide, polypeptide, antibody, and/or T-cell compositions described herein in combination with a physiologically acceptable carrier. In certain preferred embodiments, the
25 pharmaceutical compositions of the invention comprise immunogenic polynucleotide and/or polypeptide compositions of the invention for use in prophylactic and therapeutic vaccine applications. Vaccine preparation is generally described in, for example, M.F. Powell and M.J. Newman, eds., "Vaccine Design (the subunit and adjuvant approach)," Plenum Press (NY, 1995). Generally, such compositions will comprise one or more

polynucleotide and/or polypeptide compositions of the present invention in combination with one or more immunostimulants.

It will be apparent that any of the pharmaceutical compositions described herein can contain pharmaceutically acceptable salts of the polynucleotides and polypeptides of the invention. Such salts can be prepared, for example, from
5 pharmaceutically acceptable non-toxic bases, including organic bases (*e.g.*, salts of primary, secondary and tertiary amines and basic amino acids) and inorganic bases (*e.g.*, sodium, potassium, lithium, ammonium, calcium and magnesium salts).

In another embodiment, illustrative immunogenic compositions, *e.g.*,
10 vaccine compositions, of the present invention comprise DNA encoding one or more of the polypeptides as described above, such that the polypeptide is generated *in situ*. As noted above, the polynucleotide may be administered within any of a variety of delivery systems known to those of ordinary skill in the art. Indeed, numerous gene delivery techniques are well known in the art, such as those described by Rolland, *Crit. Rev.*
15 *Therap. Drug Carrier Systems* 15:143-198, 1998, and references cited therein. Appropriate polynucleotide expression systems will, of course, contain the necessary regulatory DNA regulatory sequences for expression in a patient (such as a suitable promoter and terminating signal). Alternatively, bacterial delivery systems may involve the administration of a bacterium (such as *Bacillus-Calmette-Guerrin*) that expresses an
20 immunogenic portion of the polypeptide on its cell surface or secretes such an epitope.

Therefore, in certain embodiments, polynucleotides encoding immunogenic polypeptides described herein are introduced into suitable mammalian host cells for expression using any of a number of known viral-based systems. In one illustrative embodiment, retroviruses provide a convenient and effective platform for
25 gene delivery systems. A selected nucleotide sequence encoding a polypeptide of the present invention can be inserted into a vector and packaged in retroviral particles using techniques known in the art. The recombinant virus can then be isolated and delivered to a subject. A number of illustrative retroviral systems have been described (*e.g.*, U.S. Pat. No. 5,219,740; Miller and Rosman (1989) *BioTechniques* 7:980-990; Miller, A. D.
30 (1990) *Human Gene Therapy* 1:5-14; Scarpa et al. (1991) *Virology* 180:849-852; Burns

et al. (1993) Proc. Natl. Acad. Sci. USA 90:8033-8037; and Boris-Lawrie and Temin (1993) Cur. Opin. Genet. Develop. 3:102-109.

In addition, a number of illustrative adenovirus-based systems have also been described. Unlike retroviruses which integrate into the host genome, adenoviruses
5 persist extrachromosomally thus minimizing the risks associated with insertional mutagenesis (Haj-Ahmad and Graham (1986) J. Virol. 57:267-274; Bett et al. (1993) J. Virol. 67:5911-5921; Mittereder et al. (1994) Human Gene Therapy 5:717-729; Seth et al. (1994) J. Virol. 68:933-940; Barr et al. (1994) Gene Therapy 1:51-58; Berkner, K. L. (1988) BioTechniques 6:616-629; and Rich et al. (1993) Human Gene Therapy 4:461-
10 476).

Various adeno-associated virus (AAV) vector systems have also been developed for polynucleotide delivery. AAV vectors can be readily constructed using techniques well known in the art. See, *e.g.*, U.S. Pat. Nos. 5,173,414 and 5,139,941; International Publication Nos. WO 92/01070 and WO 93/03769; Lebkowski et al.
15 (1988) Molec. Cell. Biol. 8:3988-3996; Vincent et al. (1990) Vaccines 90 (Cold Spring Harbor Laboratory Press); Carter, B. J. (1992) Current Opinion in Biotechnology 3:533-539; Muzyczka, N. (1992) Current Topics in Microbiol. and Immunol. 158:97-129; Kotin, R. M. (1994) Human Gene Therapy 5:793-801; Shelling and Smith (1994) Gene Therapy 1:165-169; and Zhou et al. (1994) J. Exp. Med. 179:1867-1875.

20 Additional viral vectors useful for delivering the polynucleotides encoding polypeptides of the present invention by gene transfer include those derived from the pox family of viruses, such as vaccinia virus and avian poxvirus. By way of example, vaccinia virus recombinants expressing the novel molecules can be constructed as follows. The DNA encoding a polypeptide is first inserted into an
25 appropriate vector so that it is adjacent to a vaccinia promoter and flanking vaccinia DNA sequences, such as the sequence encoding thymidine kinase (TK). This vector is then used to transfect cells which are simultaneously infected with vaccinia. Homologous recombination serves to insert the vaccinia promoter plus the gene encoding the polypeptide of interest into the viral genome. The resulting TK.sup.(-)

recombinant can be selected by culturing the cells in the presence of 5-bromodeoxyuridine and picking viral plaques resistant thereto.

A vaccinia-based infection/transfection system can be conveniently used to provide for inducible, transient expression or coexpression of one or more polypeptides described herein in host cells of an organism. In this particular system, cells are first infected in vitro with a vaccinia virus recombinant that encodes the bacteriophage T7 RNA polymerase. This polymerase displays exquisite specificity in that it only transcribes templates bearing T7 promoters. Following infection, cells are transfected with the polynucleotide or polynucleotides of interest, driven by a T7 promoter. The polymerase expressed in the cytoplasm from the vaccinia virus recombinant transcribes the transfected DNA into RNA which is then translated into polypeptide by the host translational machinery. The method provides for high level, transient, cytoplasmic production of large quantities of RNA and its translation products. See, *e.g.*, Elroy-Stein and Moss, Proc. Natl. Acad. Sci. USA (1990) 87:6743-6747; Fuerst et al. Proc. Natl. Acad. Sci. USA (1986) 83:8122-8126.

Alternatively, avipoxviruses, such as the fowlpox and canarypox viruses, can also be used to deliver the coding sequences of interest. Recombinant avipox viruses, expressing immunogens from mammalian pathogens, are known to confer protective immunity when administered to non-avian species. The use of an Avipox vector is particularly desirable in human and other mammalian species since members of the Avipox genus can only productively replicate in susceptible avian species and therefore are not infective in mammalian cells. Methods for producing recombinant Avipoxviruses are known in the art and employ genetic recombination, as described above with respect to the production of vaccinia viruses. See, *e.g.*, WO 91/12882; WO 89/03429; and WO 92/03545.

Any of a number of alphavirus vectors can also be used for delivery of polynucleotide compositions of the present invention, such as those vectors described in U.S. Patent Nos. 5,843,723; 6,015,686; 6,008,035 and 6,015,694. Certain vectors based on Venezuelan Equine Encephalitis (VEE) can also be used, illustrative examples of which can be found in U.S. Patent Nos. 5,505,947 and 5,643,576.

Moreover, molecular conjugate vectors, such as the adenovirus chimeric vectors described in Michael et al. *J. Biol. Chem.* (1993) 268:6866-6869 and Wagner et al. *Proc. Natl. Acad. Sci. USA* (1992) 89:6099-6103, can also be used for gene delivery under the invention.

5 Additional illustrative information on these and other known viral-based delivery systems can be found, for example, in Fisher-Hoch et al., *Proc. Natl. Acad. Sci. USA* 86:317-321, 1989; Flexner et al., *Ann. N.Y. Acad. Sci.* 569:86-103, 1989; Flexner et al., *Vaccine* 8:17-21, 1990; U.S. Patent Nos. 4,603,112, 4,769,330, and 5,017,487; WO 89/01973; U.S. Patent No. 4,777,127; GB 2,200,651; EP 0,345,242; 10 WO 91/02805; Berkner, *Biotechniques* 6:616-627, 1988; Rosenfeld et al., *Science* 252:431-434, 1991; Kolls et al., *Proc. Natl. Acad. Sci. USA* 91:215-219, 1994; Kass-Eisler et al., *Proc. Natl. Acad. Sci. USA* 90:11498-11502, 1993; Guzman et al., *Circulation* 88:2838-2848, 1993; and Guzman et al., *Cir. Res.* 73:1202-1207, 1993.

In certain embodiments, a polynucleotide may be integrated into the 15 genome of a target cell. This integration may be in the specific location and orientation *via* homologous recombination (gene replacement) or it may be integrated in a random, non-specific location (gene augmentation). In yet further embodiments, the polynucleotide may be stably maintained in the cell as a separate, episomal segment of DNA. Such polynucleotide segments or "episomes" encode sequences sufficient to 20 permit maintenance and replication independent of or in synchronization with the host cell cycle. The manner in which the expression construct is delivered to a cell and where in the cell the polynucleotide remains is dependent on the type of expression construct employed.

In another embodiment of the invention, a polynucleotide is 25 administered/delivered as "naked" DNA, for example as described in Ulmer et al., *Science* 259:1745-1749, 1993 and reviewed by Cohen, *Science* 259:1691-1692, 1993. The uptake of naked DNA may be increased by coating the DNA onto biodegradable beads, which are efficiently transported into the cells.

In still another embodiment, a composition of the present invention can 30 be delivered via a particle bombardment approach, many of which have been described.

In one illustrative example, gas-driven particle acceleration can be achieved with devices such as those manufactured by Powderject Pharmaceuticals PLC (Oxford, UK) and Powderject Vaccines Inc. (Madison, WI), some examples of which are described in U.S. Patent Nos. 5,846,796; 6,010,478; 5,865,796; 5,584,807; and EP Patent No. 0500
5 799. This approach offers a needle-free delivery approach wherein a dry powder formulation of microscopic particles, such as polynucleotide or polypeptide particles, are accelerated to high speed within a helium gas jet generated by a hand held device, propelling the particles into a target tissue of interest.

In a related embodiment, other devices and methods that may be useful
10 for gas-driven needle-less injection of compositions of the present invention include those provided by Bioject, Inc. (Portland, OR), some examples of which are described in U.S. Patent Nos. 4,790,824; 5,064,413; 5,312,335; 5,383,851; 5,399,163; 5,520,639 and 5,993,412.

According to another embodiment, the pharmaceutical compositions
15 described herein will comprise one or more immunostimulants in addition to the immunogenic polynucleotide, polypeptide, antibody, T-cell and/or APC compositions of this invention. An immunostimulant refers to essentially any substance that enhances or potentiates an immune response (antibody and/or cell-mediated) to an exogenous antigen. One preferred type of immunostimulant comprises an adjuvant. Many
20 adjuvants contain a substance designed to protect the antigen from rapid catabolism, such as aluminum hydroxide or mineral oil, and a stimulator of immune responses, such as lipid A, *Bordetella pertussis* or *Mycobacterium tuberculosis* derived proteins. Certain adjuvants are commercially available as, for example, Freund's Incomplete Adjuvant and Complete Adjuvant (Difco Laboratories, Detroit, MI); Merck Adjuvant
25 65 (Merck and Company, Inc., Rahway, NJ); AS-2 (SmithKline Beecham, Philadelphia, PA); aluminum salts such as aluminum hydroxide gel (alum) or aluminum phosphate; salts of calcium, iron or zinc; an insoluble suspension of acylated tyrosine; acylated sugars; cationically or anionically derivatized polysaccharides; polyphosphazenes; biodegradable microspheres; monophosphoryl lipid A and quil A. Cytokines, such as

GM-CSF, interleukin-2, -7, -12, and other like growth factors, may also be used as adjuvants.

Within certain embodiments of the invention, the adjuvant composition is preferably one that induces an immune response predominantly of the Th1 type. High levels of Th1-type cytokines (*e.g.*, IFN- γ , TNF α , IL-2 and IL-12) tend to favor the induction of cell mediated immune responses to an administered antigen. In contrast, high levels of Th2-type cytokines (*e.g.*, IL-4, IL-5, IL-6 and IL-10) tend to favor the induction of humoral immune responses. Following application of a vaccine as provided herein, a patient will support an immune response that includes Th1- and Th2-type responses. Within a preferred embodiment, in which a response is predominantly Th1-type, the level of Th1-type cytokines will increase to a greater extent than the level of Th2-type cytokines. The levels of these cytokines may be readily assessed using standard assays. For a review of the families of cytokines, see Mosmann and Coffman, *Ann. Rev. Immunol.* 7:145-173, 1989.

Certain preferred adjuvants for eliciting a predominantly Th1-type response include, for example, a combination of monophosphoryl lipid A, preferably 3-de-O-acylated monophosphoryl lipid A, together with an aluminum salt. MPL[®] adjuvants are available from Corixa Corporation (Seattle, WA; *see*, for example, US Patent Nos. 4,436,727; 4,877,611; 4,866,034 and 4,912,094). CpG-containing oligonucleotides (in which the CpG dinucleotide is unmethylated) also induce a predominantly Th1 response. Such oligonucleotides are well known and are described, for example, in WO 96/02555, WO 99/33488 and U.S. Patent Nos. 6,008,200 and 5,856,462. Immunostimulatory DNA sequences are also described, for example, by Sato et al., *Science* 273:352, 1996. Another preferred adjuvant comprises a saponin, such as Quil A, or derivatives thereof, including QS21 and QS7 (Aquila Biopharmaceuticals Inc., Framingham, MA); Escin; Digitonin; or *Gypsophila* or *Chenopodium quinoa* saponins. Other preferred formulations include more than one saponin in the adjuvant combinations of the present invention, for example combinations of at least two of the following group comprising QS21, QS7, Quil A, β -escin, or digitonin.

Alternatively the saponin formulations may be combined with vaccine vehicles composed of chitosan or other polycationic polymers, polylactide and polylactide-co-glycolide particles, poly-N-acetyl glucosamine-based polymer matrix, particles composed of polysaccharides or chemically modified polysaccharides, liposomes and lipid-based particles, particles composed of glycerol monoesters, etc. The saponins may also be formulated in the presence of cholesterol to form particulate structures such as liposomes or ISCOMs. Furthermore, the saponins may be formulated together with a polyoxyethylene ether or ester, in either a non-particulate solution or suspension, or in a particulate structure such as a paucilamellar liposome or ISCOM. The saponins may also be formulated with excipients such as Carbopol^R to increase viscosity, or may be formulated in a dry powder form with a powder excipient such as lactose.


In one preferred embodiment, the adjuvant system includes the combination of a monophosphoryl lipid A and a saponin derivative, such as the combination of QS21 and 3D-MPL[®] adjuvant, as described in WO 94/00153, or a less reactogenic composition where the QS21 is quenched with cholesterol, as described in WO 96/33739. Other preferred formulations comprise an oil-in-water emulsion and tocopherol. Another particularly preferred adjuvant formulation employing QS21, 3D-MPL[®] adjuvant and tocopherol in an oil-in-water emulsion is described in WO 95/17210.

Another enhanced adjuvant system involves the combination of a CpG-containing oligonucleotide and a saponin derivative particularly the combination of CpG and QS21 is disclosed in WO 00/09159. Preferably the formulation additionally comprises an oil in water emulsion and tocopherol.

Additional illustrative adjuvants for use in the pharmaceutical compositions of the invention include Montanide ISA 720 (Seppic, France), SAF (Chiron, California, United States), ISCOMS (CSL), MF-59 (Chiron), the SBAS series of adjuvants (*e.g.*, SBAS-2 or SBAS-4, available from SmithKline Beecham, Rixensart, Belgium), Detox (Enhanzyn[®]) (Corixa, Hamilton, MT), RC-529 (Corixa, Hamilton, MT) and other aminoalkyl glucosaminide 4-phosphates (AGPs), such as those described

in pending U.S. Patent Application Serial Nos. 08/853,826 and 09/074,720, the disclosures of which are incorporated herein by reference in their entireties, and polyoxyethylene ether adjuvants such as those described in WO 99/52549A1.

Other preferred adjuvants include adjuvant molecules of the general
5 formula



wherein, n is 1-50, A is a bond or $-\text{C}(\text{O})-$, R is C_{1-50} alkyl or Phenyl C_{1-50} alkyl.

One embodiment of the present invention consists of a vaccine formulation comprising a polyoxyethylene ether of general formula (I), wherein n is
10 between 1 and 50, preferably 4-24, most preferably 9; the R component is C_{1-50} , preferably $\text{C}_4\text{-C}_{20}$ alkyl and most preferably C_{12} alkyl, and A is a bond. The concentration of the polyoxyethylene ethers should be in the range 0.1-20%, preferably from 0.1-10%, and most preferably in the range 0.1-1%. Preferred polyoxyethylene ethers are selected from the following group: polyoxyethylene-9-lauryl ether,
15 polyoxyethylene-9-stearyl ether, polyoxyethylene-8-stearyl ether, polyoxyethylene-4-lauryl ether, polyoxyethylene-35-lauryl ether, and polyoxyethylene-23-lauryl ether. Polyoxyethylene ethers such as polyoxyethylene lauryl ether are described in the Merck index (12th edition: entry 7717). These adjuvant molecules are described in WO 99/52549.

20 The polyoxyethylene ether according to the general formula (I) above may, if desired, be combined with another adjuvant. For example, a preferred adjuvant combination is preferably with CpG as described in the pending UK patent application GB 9820956.2.

According to another embodiment of this invention, an immunogenic
25 composition described herein is delivered to a host via antigen presenting cells (APCs), such as dendritic cells, macrophages, B cells, monocytes and other cells that may be engineered to be efficient APCs. Such cells may, but need not, be genetically modified to increase the capacity for presenting the antigen, to improve activation and/or maintenance of the T cell response, to have anti-tumor effects *per se* and/or to be
30 immunologically compatible with the receiver (*i.e.*, matched HLA haplotype). APCs

may generally be isolated from any of a variety of biological fluids and organs, including tumor and peritumoral tissues, and may be autologous, allogeneic, syngeneic or xenogeneic cells.

Certain preferred embodiments of the present invention use dendritic
5 cells or progenitors thereof as antigen-presenting cells. Dendritic cells are highly potent APCs (Banchereau and Steinman, *Nature* 392:245-251, 1998) and have been shown to be effective as a physiological adjuvant for eliciting prophylactic or therapeutic antitumor immunity (*see* Timmerman and Levy, *Ann. Rev. Med.* 50:507-529, 1999). In general, dendritic cells may be identified based on their typical shape (stellate *in situ*,
10 with marked cytoplasmic processes (dendrites) visible *in vitro*), their ability to take up, process and present antigens with high efficiency and their ability to activate naïve T cell responses. Dendritic cells may, of course, be engineered to express specific cell-surface receptors or ligands that are not commonly found on dendritic cells *in vivo* or *ex vivo*, and such modified dendritic cells are contemplated by the present invention. As
15 an alternative to dendritic cells, secreted vesicles antigen-loaded dendritic cells (called exosomes) may be used within a vaccine (*see* Zitvogel et al., *Nature Med.* 4:594-600, 1998).

Dendritic cells and progenitors may be obtained from peripheral blood, bone marrow, tumor-infiltrating cells, peritumoral tissues-infiltrating cells, lymph
20 nodes, spleen, skin, umbilical cord blood or any other suitable tissue or fluid. For example, dendritic cells may be differentiated *ex vivo* by adding a combination of cytokines such as GM-CSF, IL-4, IL-13 and/or TNF α to cultures of monocytes harvested from peripheral blood. Alternatively, CD34 positive cells harvested from peripheral blood, umbilical cord blood or bone marrow may be differentiated into
25 dendritic cells by adding to the culture medium combinations of GM-CSF, IL-3, TNF α , CD40 ligand, LPS, flt3 ligand and/or other compound(s) that induce differentiation, maturation and proliferation of dendritic cells.

Dendritic cells are conveniently categorized as "immature" and "mature" cells, which allows a simple way to discriminate between two well characterized
30 phenotypes. However, this nomenclature should not be construed to exclude all

possible intermediate stages of differentiation. Immature dendritic cells are characterized as APC with a high capacity for antigen uptake and processing, which correlates with the high expression of Fcγ receptor and mannose receptor. The mature phenotype is typically characterized by a lower expression of these markers, but a high expression of cell surface molecules responsible for T cell activation such as class I and class II MHC, adhesion molecules (*e.g.*, CD54 and CD11) and costimulatory molecules (*e.g.*, CD40, CD80, CD86 and 4-1BB).

APCs may generally be transfected with a polynucleotide of the invention (or portion or other variant thereof) such that the encoded polypeptide, or an immunogenic portion thereof, is expressed on the cell surface. Such transfection may take place *ex vivo*, and a pharmaceutical composition comprising such transfected cells may then be used for therapeutic purposes, as described herein. Alternatively, a gene delivery vehicle that targets a dendritic or other antigen presenting cell may be administered to a patient, resulting in transfection that occurs *in vivo*. *In vivo* and *ex vivo* transfection of dendritic cells, for example, may generally be performed using any methods known in the art, such as those described in WO 97/24447, or the gene gun approach described by Mahvi et al., *Immunology and cell Biology* 75:456-460, 1997. Antigen loading of dendritic cells may be achieved by incubating dendritic cells or progenitor cells with the tumor polypeptide, DNA (naked or within a plasmid vector) or RNA; or with antigen-expressing recombinant bacterium or viruses (*e.g.*, vaccinia, fowlpox, adenovirus or lentivirus vectors). Prior to loading, the polypeptide may be covalently conjugated to an immunological partner that provides T cell help (*e.g.*, a carrier molecule). Alternatively, a dendritic cell may be pulsed with a non-conjugated immunological partner, separately or in the presence of the polypeptide.

While any suitable carrier known to those of ordinary skill in the art may be employed in the pharmaceutical compositions of this invention, the type of carrier will typically vary depending on the mode of administration. Compositions of the present invention may be formulated for any appropriate manner of administration, including for example, topical, oral, nasal, mucosal, intravenous, intracranial, intraperitoneal, subcutaneous and intramuscular administration.

Carriers for use within such pharmaceutical compositions are biocompatible, and may also be biodegradable. In certain embodiments, the formulation preferably provides a relatively constant level of active component release. In other embodiments, however, a more rapid rate of release immediately upon administration may be desired. The formulation of such compositions is well within the level of ordinary skill in the art using known techniques. Illustrative carriers useful in this regard include microparticles of poly(lactide-co-glycolide), polyacrylate, latex, starch, cellulose, dextran and the like. Other illustrative delayed-release carriers include supramolecular biovectors, which comprise a non-liquid hydrophilic core (*e.g.*, a cross-linked polysaccharide or oligosaccharide) and, optionally, an external layer comprising an amphiphilic compound, such as a phospholipid (*see e.g.*, U.S. Patent No. 5,151,254 and PCT applications WO 94/20078, WO/94/23701 and WO 96/06638). The amount of active compound contained within a sustained release formulation depends upon the site of implantation, the rate and expected duration of release and the nature of the condition to be treated or prevented.

In another illustrative embodiment, biodegradable microspheres (*e.g.*, polylactate polyglycolate) are employed as carriers for the compositions of this invention. Suitable biodegradable microspheres are disclosed, for example, in U.S. Patent Nos. 4,897,268; 5,075,109; 5,928,647; 5,811,128; 5,820,883; 5,853,763; 5,814,344, 5,407,609 and 5,942,252. Modified hepatitis B core protein carrier systems, such as described in WO/99 40934, and references cited therein, will also be useful for many applications. Another illustrative carrier/delivery system employs a carrier comprising particulate-protein complexes, such as those described in U.S. Patent No. 5,928,647, which are capable of inducing a class I-restricted cytotoxic T lymphocyte responses in a host.

The pharmaceutical compositions of the invention will often further comprise one or more buffers (*e.g.*, neutral buffered saline or phosphate buffered saline), carbohydrates (*e.g.*, glucose, mannose, sucrose or dextrans), mannitol, proteins, polypeptides or amino acids such as glycine, antioxidants, bacteriostats, chelating agents such as EDTA or glutathione, adjuvants (*e.g.*, aluminum hydroxide), solutes that

render the formulation isotonic, hypotonic or weakly hypertonic with the blood of a recipient, suspending agents, thickening agents and/or preservatives. Alternatively, compositions of the present invention may be formulated as a lyophilizate.

The pharmaceutical compositions described herein may be presented in unit-dose or multi-dose containers, such as sealed ampoules or vials. Such containers are typically sealed in such a way to preserve the sterility and stability of the formulation until use. In general, formulations may be stored as suspensions, solutions or emulsions in oily or aqueous vehicles. Alternatively, a pharmaceutical composition may be stored in a freeze-dried condition requiring only the addition of a sterile liquid carrier immediately prior to use.

The development of suitable dosing and treatment regimens for using the particular compositions described herein in a variety of treatment regimens, including *e.g.*, oral, parenteral, intravenous, intranasal, and intramuscular administration and formulation, is well known in the art, some of which are briefly discussed below for general purposes of illustration.

In certain applications, the pharmaceutical compositions disclosed herein may be delivered *via* oral administration to an animal. As such, these compositions may be formulated with an inert diluent or with an assimilable edible carrier, or they may be enclosed in hard- or soft-shell gelatin capsule, or they may be compressed into tablets, or they may be incorporated directly with the food of the diet.

The active compounds may even be incorporated with excipients and used in the form of ingestible tablets, buccal tables, troches, capsules, elixirs, suspensions, syrups, wafers, and the like (see, for example, Mathiowitz *et al.*, Nature 1997 Mar 27;386(6623):410-4; Hwang *et al.*, Crit Rev Ther Drug Carrier Syst 1998;15(3):243-84; U. S. Patent 5,641,515; U. S. Patent 5,580,579 and U. S. Patent 5,792,451). Tablets, troches, pills, capsules and the like may also contain any of a variety of additional components, for example, a binder, such as gum tragacanth, acacia, cornstarch, or gelatin; excipients, such as dicalcium phosphate; a disintegrating agent, such as corn starch, potato starch, alginic acid and the like; a lubricant, such as magnesium stearate; and a sweetening agent, such as sucrose, lactose or saccharin may

be added or a flavoring agent, such as peppermint, oil of wintergreen, or cherry flavoring. When the dosage unit form is a capsule, it may contain, in addition to materials of the above type, a liquid carrier. Various other materials may be present as coatings or to otherwise modify the physical form of the dosage unit. For instance, 5 tablets, pills, or capsules may be coated with shellac, sugar, or both. Of course, any material used in preparing any dosage unit form should be pharmaceutically pure and substantially non-toxic in the amounts employed. In addition, the active compounds may be incorporated into sustained-release preparation and formulations.

Typically, these formulations will contain at least about 0.1% of the 10 active compound or more, although the percentage of the active ingredient(s) may, of course, be varied and may conveniently be between about 1 or 2% and about 60% or 70% or more of the weight or volume of the total formulation. Naturally, the amount of active compound(s) in each therapeutically useful composition may be prepared in such a way that a suitable dosage will be obtained in any given unit dose of the compound. 15 Factors such as solubility, bioavailability, biological half-life, route of administration, product shelf life, as well as other pharmacological considerations will be contemplated by one skilled in the art of preparing such pharmaceutical formulations, and as such, a variety of dosages and treatment regimens may be desirable.

For oral administration the compositions of the present invention may 20 alternatively be incorporated with one or more excipients in the form of a mouthwash, dentifrice, buccal tablet, oral spray, or sublingual orally-administered formulation. Alternatively, the active ingredient may be incorporated into an oral solution such as one containing sodium borate, glycerin and potassium bicarbonate, or dispersed in a dentifrice, or added in a therapeutically-effective amount to a composition that may 25 include water, binders, abrasives, flavoring agents, foaming agents, and humectants. Alternatively the compositions may be fashioned into a tablet or solution form that may be placed under the tongue or otherwise dissolved in the mouth.

In certain circumstances it will be desirable to deliver the pharmaceutical compositions disclosed herein parenterally, intravenously, intramuscularly, or even 30 intraperitoneally. Such approaches are well known to the skilled artisan, some of which

are further described, for example, in U. S. Patent 5,543,158; U. S. Patent 5,641,515 and U. S. Patent 5,399,363. In certain embodiments, solutions of the active compounds as free base or pharmacologically acceptable salts may be prepared in water suitably mixed with a surfactant, such as hydroxypropylcellulose. Dispersions may also be prepared in glycerol, liquid polyethylene glycols, and mixtures thereof and in oils. Under ordinary conditions of storage and use, these preparations generally will contain a preservative to prevent the growth of microorganisms.

Illustrative pharmaceutical forms suitable for injectable use include sterile aqueous solutions or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions (for example, see U. S. Patent 5,466,468). In all cases the form must be sterile and must be fluid to the extent that easy syringability exists. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms, such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (*e.g.*, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), suitable mixtures thereof, and/or vegetable oils. Proper fluidity may be maintained, for example, by the use of a coating, such as lecithin, by the maintenance of the required particle size in the case of dispersion and/or by the use of surfactants. The prevention of the action of microorganisms can be facilitated by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars or sodium chloride. Prolonged absorption of the injectable compositions can be brought about by the use in the compositions of agents delaying absorption, for example, aluminum monostearate and gelatin.

In one embodiment, for parenteral administration in an aqueous solution, the solution should be suitably buffered if necessary and the liquid diluent first rendered isotonic with sufficient saline or glucose. These particular aqueous solutions are especially suitable for intravenous, intramuscular, subcutaneous and intraperitoneal administration. In this connection, a sterile aqueous medium that can be employed will

be known to those of skill in the art in light of the present disclosure. For example, one dosage may be dissolved in 1 ml of isotonic NaCl solution and either added to 1000 ml of hypodermoclysis fluid or injected at the proposed site of infusion, (see for example, "Remington's Pharmaceutical Sciences" 15th Edition, pages 1035-1038 and 1570-
5 1580). Some variation in dosage will necessarily occur depending on the condition of the subject being treated. Moreover, for human administration, preparations will of course preferably meet sterility, pyrogenicity, and the general safety and purity standards as required by FDA Office of Biologics standards.

In another embodiment of the invention, the compositions disclosed
10 herein may be formulated in a neutral or salt form. Illustrative pharmaceutically-acceptable salts include the acid addition salts (formed with the free amino groups of the protein) and which are formed with inorganic acids such as, for example, hydrochloric or phosphoric acids, or such organic acids as acetic, oxalic, tartaric, mandelic, and the like. Salts formed with the free carboxyl groups can also be
15 derived from inorganic bases such as, for example, sodium, potassium, ammonium, calcium, or ferric hydroxides, and such organic bases as isopropylamine, trimethylamine, histidine, procaine and the like. Upon formulation, solutions will be administered in a manner compatible with the dosage formulation and in such amount as is therapeutically effective.

20 The carriers can further comprise any and all solvents, dispersion media, vehicles, coatings, diluents, antibacterial and antifungal agents, isotonic and absorption delaying agents, buffers, carrier solutions, suspensions, colloids, and the like. The use of such media and agents for pharmaceutical active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active
25 ingredient, its use in the therapeutic compositions is contemplated. Supplementary active ingredients can also be incorporated into the compositions. The phrase "pharmaceutically-acceptable" refers to molecular entities and compositions that do not produce an allergic or similar untoward reaction when administered to a human.

In certain embodiments, the pharmaceutical compositions may be
30 delivered by intranasal sprays, inhalation, and/or other aerosol delivery vehicles.

Methods for delivering genes, nucleic acids, and peptide compositions directly to the lungs *via* nasal aerosol sprays has been described, *e.g.*, in U. S. Patent 5,756,353 and U. S. Patent 5,804,212. Likewise, the delivery of drugs using intranasal microparticle resins (Takenaga *et al.*, J Controlled Release 1998 Mar 2;52(1-2):81-7) and
5 lysophosphatidyl-glycerol compounds (U. S. Patent 5,725,871) are also well-known in the pharmaceutical arts. Likewise, illustrative transmucosal drug delivery in the form of a polytetrafluoroethylene support matrix is described in U. S. Patent 5,780,045.

In certain embodiments, liposomes, nanocapsules, microparticles, lipid particles, vesicles, and the like, are used for the introduction of the compositions of the
10 present invention into suitable host cells/organisms. In particular, the compositions of the present invention may be formulated for delivery either encapsulated in a lipid particle, a liposome, a vesicle, a nanosphere, or a nanoparticle or the like. Alternatively, compositions of the present invention can be bound, either covalently or non-covalently, to the surface of such carrier vehicles.

15 The formation and use of liposome and liposome-like preparations as potential drug carriers is generally known to those of skill in the art (see for example, Lasic, Trends Biotechnol 1998 Jul;16(7):307-21; Takakura, Nippon Rinsho 1998 Mar;56(3):691-5; Chandran *et al.*, Indian J Exp Biol. 1997 Aug;35(8):801-9; Margalit, Crit Rev Ther Drug Carrier Syst. 1995;12(2-3):233-61; U.S. Patent 5,567,434; U.S.
20 Patent 5,552,157; U.S. Patent 5,565,213; U.S. Patent 5,738,868 and U.S. Patent 5,795,587, each specifically incorporated herein by reference in its entirety).

Liposomes have been used successfully with a number of cell types that are normally difficult to transfect by other procedures, including T cell suspensions, primary hepatocyte cultures and PC 12 cells (Renneisen *et al.*, J Biol Chem. 1990 Sep
25 25;265(27):16337-42; Muller *et al.*, DNA Cell Biol. 1990 Apr;9(3):221-9). In addition, liposomes are free of the DNA length constraints that are typical of viral-based delivery systems. Liposomes have been used effectively to introduce genes, various drugs, radiotherapeutic agents, enzymes, viruses, transcription factors, allosteric effectors and the like, into a variety of cultured cell lines and animals. Furthermore, the use of

liposomes does not appear to be associated with autoimmune responses or unacceptable toxicity after systemic delivery.

In certain embodiments, liposomes are formed from phospholipids that are dispersed in an aqueous medium and spontaneously form multilamellar concentric
5 bilayer vesicles (also termed multilamellar vesicles (MLVs)).

Alternatively, in other embodiments, the invention provides for pharmaceutically-acceptable nanocapsule formulations of the compositions of the present invention. Nanocapsules can generally entrap compounds in a stable and reproducible way (see, for example, Quintanar-Guerrero *et al.*, Drug Dev Ind Pharm.
10 1998 Dec;24(12):1113-28). To avoid side effects due to intracellular polymeric overloading, such ultrafine particles (sized around 0.1 μm) may be designed using polymers able to be degraded *in vivo*. Such particles can be made as described, for example, by Couvreur *et al.*, Crit Rev Ther Drug Carrier Syst. 1988;5(1):1-20; zur Muhlen *et al.*, Eur J Pharm Biopharm. 1998 Mar;45(2):149-55; Zambaux *et al.* J
15 Controlled Release. 1998 Jan 2;50(1-3):31-40; and U. S. Patent 5,145,684.

Cancer Therapeutic Methods

In further aspects of the present invention, the pharmaceutical compositions described herein may be used for the treatment of cancer, particularly for the immunotherapy of breast cancer. Within such methods, the pharmaceutical
20 compositions described herein are administered to a patient, typically a warm-blooded animal, preferably a human. A patient may or may not be afflicted with cancer. Accordingly, the above pharmaceutical compositions may be used to prevent the development of a cancer or to treat a patient afflicted with a cancer. Pharmaceutical compositions and vaccines may be administered either prior to or following surgical
25 removal of primary tumors and/or treatment such as administration of radiotherapy or conventional chemotherapeutic drugs. As discussed above, administration of the pharmaceutical compositions may be by any suitable method, including administration by intravenous, intraperitoneal, intramuscular, subcutaneous, intranasal, intradermal, anal, vaginal, topical and oral routes.

Within certain embodiments, immunotherapy may be active immunotherapy, in which treatment relies on the *in vivo* stimulation of the endogenous host immune system to react against tumors with the administration of immune response-modifying agents (such as polypeptides and polynucleotides as provided
5 herein).

Within other embodiments, immunotherapy may be passive immunotherapy, in which treatment involves the delivery of agents with established tumor-immune reactivity (such as effector cells or antibodies) that can directly or indirectly mediate antitumor effects and does not necessarily depend on an intact host
10 immune system. Examples of effector cells include T cells as discussed above, T lymphocytes (such as CD8⁺ cytotoxic T lymphocytes and CD4⁺ T-helper tumor-infiltrating lymphocytes), killer cells (such as Natural Killer cells and lymphokine-activated killer cells), B cells and antigen-presenting cells (such as dendritic cells and macrophages) expressing a polypeptide provided herein. T cell receptors and antibody
15 receptors specific for the polypeptides recited herein may be cloned, expressed and transferred into other vectors or effector cells for adoptive immunotherapy. The polypeptides provided herein may also be used to generate antibodies or anti-idiotypic antibodies (as described above and in U.S. Patent No. 4,918,164) for passive immunotherapy.

Effector cells may generally be obtained in sufficient quantities for adoptive immunotherapy by growth *in vitro*, as described herein. Culture conditions for expanding single antigen-specific effector cells to several billion in number with retention of antigen recognition *in vivo* are well known in the art. Such *in vitro* culture conditions typically use intermittent stimulation with antigen, often in the presence of
25 cytokines (such as IL-2) and non-dividing feeder cells. As noted above, immunoreactive polypeptides as provided herein may be used to rapidly expand antigen-specific T cell cultures in order to generate a sufficient number of cells for immunotherapy. In particular, antigen-presenting cells, such as dendritic, macrophage, monocyte, fibroblast and/or B cells, may be pulsed with immunoreactive polypeptides
30 or transfected with one or more polynucleotides using standard techniques well known

in the art. For example, antigen-presenting cells can be transfected with a polynucleotide having a promoter appropriate for increasing expression in a recombinant virus or other expression system. Cultured effector cells for use in therapy must be able to grow and distribute widely, and to survive long term *in vivo*. Studies
5 have shown that cultured effector cells can be induced to grow *in vivo* and to survive long term in substantial numbers by repeated stimulation with antigen supplemented with IL-2 (*see, for example, Cheever et al., Immunological Reviews 157:177, 1997*).

Alternatively, a vector expressing a polypeptide recited herein may be introduced into antigen presenting cells taken from a patient and clonally propagated *ex*
10 *vivo* for transplant back into the same patient. Transfected cells may be reintroduced into the patient using any means known in the art, preferably in sterile form by intravenous, intracavitary, intraperitoneal or intratumor administration.

Routes and frequency of administration of the therapeutic compositions described herein, as well as dosage, will vary from individual to individual, and may be
15 readily established using standard techniques. In general, the pharmaceutical compositions and vaccines may be administered by injection (*e.g.*, intracutaneous, intramuscular, intravenous or subcutaneous), intranasally (*e.g.*, by aspiration) or orally. Preferably, between 1 and 10 doses may be administered over a 52 week period. Preferably, 6 doses are administered, at intervals of 1 month, and booster vaccinations
20 may be given periodically thereafter. Alternate protocols may be appropriate for individual patients. A suitable dose is an amount of a compound that, when administered as described above, is capable of promoting an anti-tumor immune response, and is at least 10-50% above the basal (*i.e.*, untreated) level. Such response can be monitored by measuring the anti-tumor antibodies in a patient or by vaccine-
25 dependent generation of cytolytic effector cells capable of killing the patient's tumor cells *in vitro*. Such vaccines should also be capable of causing an immune response that leads to an improved clinical outcome (*e.g.*, more frequent remissions, complete or partial or longer disease-free survival) in vaccinated patients as compared to non-vaccinated patients. In general, for pharmaceutical compositions and vaccines
30 comprising one or more polypeptides, the amount of each polypeptide present in a dose

ranges from about 25 µg to 5 mg per kg of host. Suitable dose sizes will vary with the size of the patient, but will typically range from about 0.1 mL to about 5 mL.

In general, an appropriate dosage and treatment regimen provides the active compound(s) in an amount sufficient to provide therapeutic and/or prophylactic benefit. Such a response can be monitored by establishing an improved clinical outcome (e.g., more frequent remissions, complete or partial, or longer disease-free survival) in treated patients as compared to non-treated patients. Increases in preexisting immune responses to a tumor protein generally correlate with an improved clinical outcome. Such immune responses may generally be evaluated using standard proliferation, cytotoxicity or cytokine assays, which may be performed using samples obtained from a patient before and after treatment.

Cancer Detection and Diagnostic Compositions, Methods and Kits

In general, a cancer may be detected in a patient based on the presence of one or more breast tumor proteins and/or polynucleotides encoding such proteins in a biological sample (for example, blood, sera, sputum urine and/or tumor biopsies) obtained from the patient. In other words, such proteins may be used as markers to indicate the presence or absence of a cancer such as breast cancer. In addition, such proteins may be useful for the detection of other cancers. The binding agents provided herein generally permit detection of the level of antigen that binds to the agent in the biological sample. Polynucleotide primers and probes may be used to detect the level of mRNA encoding a tumor protein, which is also indicative of the presence or absence of a cancer. In general, a breast tumor sequence should be present at a level that is at least three fold higher in tumor tissue than in normal tissue

There are a variety of assay formats known to those of ordinary skill in the art for using a binding agent to detect polypeptide markers in a sample. See, e.g., Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, 1988. In general, the presence or absence of a cancer in a patient may be determined by (a) contacting a biological sample obtained from a patient with a binding agent; (b)

detecting in the sample a level of polypeptide that binds to the binding agent; and (c) comparing the level of polypeptide with a predetermined cut-off value.

In a preferred embodiment, the assay involves the use of binding agent immobilized on a solid support to bind to and remove the polypeptide from the remainder of the sample. The bound polypeptide may then be detected using a detection reagent that contains a reporter group and specifically binds to the binding agent/polypeptide complex. Such detection reagents may comprise, for example, a binding agent that specifically binds to the polypeptide or an antibody or other agent that specifically binds to the binding agent, such as an anti-immunoglobulin, protein G, protein A or a lectin. Alternatively, a competitive assay may be utilized, in which a polypeptide is labeled with a reporter group and allowed to bind to the immobilized binding agent after incubation of the binding agent with the sample. The extent to which components of the sample inhibit the binding of the labeled polypeptide to the binding agent is indicative of the reactivity of the sample with the immobilized binding agent. Suitable polypeptides for use within such assays include full length breast tumor proteins and polypeptide portions thereof to which the binding agent binds, as described above.

The solid support may be any material known to those of ordinary skill in the art to which the tumor protein may be attached. For example, the solid support may be a test well in a microtiter plate or a nitrocellulose or other suitable membrane. Alternatively, the support may be a bead or disc, such as glass, fiberglass, latex or a plastic material such as polystyrene or polyvinylchloride. The support may also be a magnetic particle or a fiber optic sensor, such as those disclosed, for example, in U.S. Patent No. 5,359,681. The binding agent may be immobilized on the solid support using a variety of techniques known to those of skill in the art, which are amply described in the patent and scientific literature. In the context of the present invention, the term "immobilization" refers to both noncovalent association, such as adsorption, and covalent attachment (which may be a direct linkage between the agent and functional groups on the support or may be a linkage by way of a cross-linking agent). Immobilization by adsorption to a well in a microtiter plate or to a membrane is

preferred. In such cases, adsorption may be achieved by contacting the binding agent, in a suitable buffer, with the solid support for a suitable amount of time. The contact time varies with temperature, but is typically between about 1 hour and about 1 day. In general, contacting a well of a plastic microtiter plate (such as polystyrene or polyvinylchloride) with an amount of binding agent ranging from about 10 ng to about 10 μ g, and preferably about 100 ng to about 1 μ g, is sufficient to immobilize an adequate amount of binding agent.

Covalent attachment of binding agent to a solid support may generally be achieved by first reacting the support with a bifunctional reagent that will react with both the support and a functional group, such as a hydroxyl or amino group, on the binding agent. For example, the binding agent may be covalently attached to supports having an appropriate polymer coating using benzoquinone or by condensation of an aldehyde group on the support with an amine and an active hydrogen on the binding partner (*see, e.g.,* Pierce Immunotechnology Catalog and Handbook, 1991, at A12-A13).

In certain embodiments, the assay is a two-antibody sandwich assay. This assay may be performed by first contacting an antibody that has been immobilized on a solid support, commonly the well of a microtiter plate, with the sample, such that polypeptides within the sample are allowed to bind to the immobilized antibody. Unbound sample is then removed from the immobilized polypeptide-antibody complexes and a detection reagent (preferably a second antibody capable of binding to a different site on the polypeptide) containing a reporter group is added. The amount of detection reagent that remains bound to the solid support is then determined using a method appropriate for the specific reporter group.

More specifically, once the antibody is immobilized on the support as described above, the remaining protein binding sites on the support are typically blocked. Any suitable blocking agent known to those of ordinary skill in the art, such as bovine serum albumin or Tween 20™ (Sigma Chemical Co., St. Louis, MO). The immobilized antibody is then incubated with the sample, and polypeptide is allowed to bind to the antibody. The sample may be diluted with a suitable diluent, such as

phosphate-buffered saline (PBS) prior to incubation. In general, an appropriate contact time (*i.e.*, incubation time) is a period of time that is sufficient to detect the presence of polypeptide within a sample obtained from an individual with breast cancer. Preferably, the contact time is sufficient to achieve a level of binding that is at least about 95% of that achieved at equilibrium between bound and unbound polypeptide. Those of ordinary skill in the art will recognize that the time necessary to achieve equilibrium may be readily determined by assaying the level of binding that occurs over a period of time. At room temperature, an incubation time of about 30 minutes is generally sufficient.

10 Unbound sample may then be removed by washing the solid support with an appropriate buffer, such as PBS containing 0.1% Tween 20™. The second antibody, which contains a reporter group, may then be added to the solid support. Preferred reporter groups include those groups recited above.

 The detection reagent is then incubated with the immobilized antibody-polypeptide complex for an amount of time sufficient to detect the bound polypeptide. An appropriate amount of time may generally be determined by assaying the level of binding that occurs over a period of time. Unbound detection reagent is then removed and bound detection reagent is detected using the reporter group. The method employed for detecting the reporter group depends upon the nature of the reporter group. For radioactive groups, scintillation counting or autoradiographic methods are generally appropriate. Spectroscopic methods may be used to detect dyes, luminescent groups and fluorescent groups. Biotin may be detected using avidin, coupled to a different reporter group (commonly a radioactive or fluorescent group or an enzyme). Enzyme reporter groups may generally be detected by the addition of substrate (generally for a specific period of time), followed by spectroscopic or other analysis of the reaction products.

 To determine the presence or absence of a cancer, such as breast cancer, the signal detected from the reporter group that remains bound to the solid support is generally compared to a signal that corresponds to a predetermined cut-off value. In one preferred embodiment, the cut-off value for the detection of a cancer is the average

mean signal obtained when the immobilized antibody is incubated with samples from patients without the cancer. In general, a sample generating a signal that is three standard deviations above the predetermined cut-off value is considered positive for the cancer. In an alternate preferred embodiment, the cut-off value is determined using a Receiver Operator Curve, according to the method of Sackett et al., *Clinical Epidemiology: A Basic Science for Clinical Medicine*, Little Brown and Co., 1985, p. 106-7. Briefly, in this embodiment, the cut-off value may be determined from a plot of pairs of true positive rates (*i.e.*, sensitivity) and false positive rates (100%-specificity) that correspond to each possible cut-off value for the diagnostic test result. The cut-off value on the plot that is the closest to the upper left-hand corner (*i.e.*, the value that encloses the largest area) is the most accurate cut-off value, and a sample generating a signal that is higher than the cut-off value determined by this method may be considered positive. Alternatively, the cut-off value may be shifted to the left along the plot, to minimize the false positive rate, or to the right, to minimize the false negative rate. In general, a sample generating a signal that is higher than the cut-off value determined by this method is considered positive for a cancer.

In a related embodiment, the assay is performed in a flow-through or strip test format, wherein the binding agent is immobilized on a membrane, such as nitrocellulose. In the flow-through test, polypeptides within the sample bind to the immobilized binding agent as the sample passes through the membrane. A second, labeled binding agent then binds to the binding agent-polypeptide complex as a solution containing the second binding agent flows through the membrane. The detection of bound second binding agent may then be performed as described above. In the strip test format, one end of the membrane to which binding agent is bound is immersed in a solution containing the sample. The sample migrates along the membrane through a region containing second binding agent and to the area of immobilized binding agent. Concentration of second binding agent at the area of immobilized antibody indicates the presence of a cancer. Typically, the concentration of second binding agent at that site generates a pattern, such as a line, that can be read visually. The absence of such a pattern indicates a negative result. In general, the amount of binding agent immobilized

on the membrane is selected to generate a visually discernible pattern when the biological sample contains a level of polypeptide that would be sufficient to generate a positive signal in the two-antibody sandwich assay, in the format discussed above. Preferred binding agents for use in such assays are antibodies and antigen-binding
5 fragments thereof. Preferably, the amount of antibody immobilized on the membrane ranges from about 25 ng to about 1 μ g, and more preferably from about 50 ng to about 500 ng. Such tests can typically be performed with a very small amount of biological sample.

Of course, numerous other assay protocols exist that are suitable for use
10 with the tumor proteins or binding agents of the present invention. The above descriptions are intended to be exemplary only. For example, it will be apparent to those of ordinary skill in the art that the above protocols may be readily modified to use tumor polypeptides to detect antibodies that bind to such polypeptides in a biological sample. The detection of such tumor protein specific antibodies may correlate with the
15 presence of a cancer.

A cancer may also, or alternatively, be detected based on the presence of T cells that specifically react with a tumor protein in a biological sample. Within certain methods, a biological sample comprising CD4⁺ and/or CD8⁺ T cells isolated from a patient is incubated with a tumor polypeptide, a polynucleotide encoding such a
20 polypeptide and/or an APC that expresses at least an immunogenic portion of such a polypeptide, and the presence or absence of specific activation of the T cells is detected. Suitable biological samples include, but are not limited to, isolated T cells. For example, T cells may be isolated from a patient by routine techniques (such as by Ficoll/Hypaque density gradient centrifugation of peripheral blood lymphocytes). T
25 cells may be incubated *in vitro* for 2-9 days (typically 4 days) at 37°C with polypeptide (e.g., 5 - 25 μ g/ml). It may be desirable to incubate another aliquot of a T cell sample in the absence of tumor polypeptide to serve as a control. For CD4⁺ T cells, activation is preferably detected by evaluating proliferation of the T cells. For CD8⁺ T cells, activation is preferably detected by evaluating cytolytic activity. A level of proliferation

that is at least two fold greater and/or a level of cytolytic activity that is at least 20% greater than in disease-free patients indicates the presence of a cancer in the patient.

As noted above, a cancer may also, or alternatively, be detected based on the level of mRNA encoding a tumor protein in a biological sample. For example, at
5 least two oligonucleotide primers may be employed in a polymerase chain reaction (PCR) based assay to amplify a portion of a tumor cDNA derived from a biological sample, wherein at least one of the oligonucleotide primers is specific for (*i.e.*, hybridizes to) a polynucleotide encoding the tumor protein. The amplified cDNA is then separated and detected using techniques well known in the art, such as gel
10 electrophoresis. Similarly, oligonucleotide probes that specifically hybridize to a polynucleotide encoding a tumor protein may be used in a hybridization assay to detect the presence of polynucleotide encoding the tumor protein in a biological sample.

To permit hybridization under assay conditions, oligonucleotide primers and probes should comprise an oligonucleotide sequence that has at least about 60%,
15 preferably at least about 75% and more preferably at least about 90%, identity to a portion of a polynucleotide encoding a tumor protein of the invention that is at least 10 nucleotides, and preferably at least 20 nucleotides, in length. Preferably, oligonucleotide primers and/or probes hybridize to a polynucleotide encoding a polypeptide described herein under moderately stringent conditions, as defined above.
20 Oligonucleotide primers and/or probes which may be usefully employed in the diagnostic methods described herein preferably are at least 10-40 nucleotides in length. In a preferred embodiment, the oligonucleotide primers comprise at least 10 contiguous nucleotides, more preferably at least 15 contiguous nucleotides, of a DNA molecule having a sequence as disclosed herein. Techniques for both PCR based assays and
25 hybridization assays are well known in the art (*see*, for example, Mullis et al., *Cold Spring Harbor Symp. Quant. Biol.*, 51:263, 1987; Erlich ed., *PCR Technology*, Stockton Press, NY, 1989).

One preferred assay employs RT-PCR, in which PCR is applied in conjunction with reverse transcription. Typically, RNA is extracted from a biological
30 sample, such as biopsy tissue, and is reverse transcribed to produce cDNA molecules.

PCR amplification using at least one specific primer generates a cDNA molecule, which may be separated and visualized using, for example, gel electrophoresis. Amplification may be performed on biological samples taken from a test patient and from an individual who is not afflicted with a cancer. The amplification reaction may be performed on several dilutions of cDNA spanning two orders of magnitude. A two-fold or greater increase in expression in several dilutions of the test patient sample as compared to the same dilutions of the non-cancerous sample is typically considered positive.

In another embodiment, the compositions described herein may be used as markers for the progression of cancer. In this embodiment, assays as described above for the diagnosis of a cancer may be performed over time, and the change in the level of reactive polypeptide(s) or polynucleotide(s) evaluated. For example, the assays may be performed every 24-72 hours for a period of 6 months to 1 year, and thereafter performed as needed. In general, a cancer is progressing in those patients in whom the level of polypeptide or polynucleotide detected increases over time. In contrast, the cancer is not progressing when the level of reactive polypeptide or polynucleotide either remains constant or decreases with time.

Certain *in vivo* diagnostic assays may be performed directly on a tumor. One such assay involves contacting tumor cells with a binding agent. The bound binding agent may then be detected directly or indirectly via a reporter group. Such binding agents may also be used in histological applications. Alternatively, polynucleotide probes may be used within such applications.

As noted above, to improve sensitivity, multiple tumor protein markers may be assayed within a given sample. It will be apparent that binding agents specific for different proteins provided herein may be combined within a single assay. Further, multiple primers or probes may be used concurrently. The selection of tumor protein markers may be based on routine experiments to determine combinations that results in optimal sensitivity. In addition, or alternatively, assays for tumor proteins provided herein may be combined with assays for other known tumor antigens.

The present invention further provides kits for use within any of the above diagnostic methods. Such kits typically comprise two or more components necessary for performing a diagnostic assay. Components may be compounds, reagents, containers and/or equipment. For example, one container within a kit may contain a
5 monoclonal antibody or fragment thereof that specifically binds to a tumor protein. Such antibodies or fragments may be provided attached to a support material, as described above. One or more additional containers may enclose elements, such as reagents or buffers, to be used in the assay. Such kits may also, or alternatively, contain a detection reagent as described above that contains a reporter group suitable for direct
10 or indirect detection of antibody binding.

Alternatively, a kit may be designed to detect the level of mRNA encoding a tumor protein in a biological sample. Such kits generally comprise at least one oligonucleotide probe or primer, as described above, that hybridizes to a polynucleotide encoding a tumor protein. Such an oligonucleotide may be used, for
15 example, within a PCR or hybridization assay. Additional components that may be present within such kits include a second oligonucleotide and/or a diagnostic reagent or container to facilitate the detection of a polynucleotide encoding a tumor protein.

The following Examples are offered by way of illustration and not by way of limitation.

20

EXAMPLE I

PREPARATION OF BREAST TUMOR-SPECIFIC CDNAS USING DIFFERENTIAL DISPLAY RT-PCR

This Example illustrates the preparation of cDNA molecules encoding
25 breast tumor-specific polypeptides using a differential display screen.

A. Preparation of B18Ag1 cDNA and Characterization of mRNA Expression

Tissue samples were prepared from breast tumor and normal tissue of a patient with breast cancer that was confirmed by pathology after removal from the patient. Normal RNA and tumor RNA was extracted from the samples and mRNA was

isolated and converted into cDNA using a (dT)₁₂AG (SEQ ID NO:130) anchored 3' primer. Differential display PCR was then executed using a randomly chosen primer (CTTCAACCTC) (SEQ ID NO:103). Amplification conditions were standard buffer containing 1.5 mM MgCl₂, 20 pmol of primer, 500 pmol dNTP, and 1 unit of *Taq* DNA polymerase (Perkin-Elmer, Branchburg, NJ). Forty cycles of amplification were performed using 94°C denaturation for 30 seconds, 42°C annealing for 1 minute, and 72°C extension for 30 seconds. An RNA fingerprint containing 76 amplified products was obtained. Although the RNA fingerprint of breast tumor tissue was over 98% identical to that of the normal breast tissue, a band was repeatedly observed to be specific to the RNA fingerprint pattern of the tumor. This band was cut out of a silver stained gel, subcloned into the T-vector (Novagen, Madison, WI) and sequenced.

The sequence of the cDNA, referred to as B18Ag1, is provided in SEQ ID NO:1. A database search of GENBANK and EMBL revealed that the B18Ag1 fragment initially cloned is 77% identical to the endogenous human retroviral element S71, which is a truncated retroviral element homologous to the Simian Sarcoma Virus (SSV). S71 contains an incomplete *gag* gene, a portion of the *pol* gene and an LTR-like structure at the 3' terminus (*see* Werner et al., *Virology* 174:225-238 (1990)). B18Ag1 is also 64% identical to SSV in the region corresponding to the P30 (*gag*) locus. B18Ag1 contains three separate and incomplete reading frames covering a region which shares considerable homology to a wide variety of *gag* proteins of retroviruses which infect mammals. In addition, the homology to S71 is not just within the *gag* gene, but spans several kb of sequence including an LTR.

B18Ag1-specific PCR primers were synthesized using computer analysis guidelines. RT-PCR amplification (94°C, 30 seconds; 60°C → 42°C, 30 seconds; 72°C, 30 seconds for 40 cycles) confirmed that B18Ag1 represents an actual mRNA sequence present at relatively high levels in the patient's breast tumor tissue. The primers used in amplification were B18Ag1-1 (CTG CCT GAG CCA CAA ATG) (SEQ ID NO:128) and B18Ag1-4 (CCG GAG GAG GAA GCT AGA GGA ATA) (SEQ ID NO:129) at a 3.5 mM magnesium concentration and a pH of 8.5, and B18Ag1-2 (ATG GCT ATT TTC GGG GCC TGA CA) (SEQ ID NO:126) and B18Ag1-3 (CCG GTA TCT CCT

CGT GGG TAT T) (SEQ ID NO:127) at 2 mM magnesium at pH 9.5. The same experiments showed exceedingly low to nonexistent levels of expression in this patient's normal breast tissue (*see* Figure 1). RT-PCR experiments were then used to show that B18Ag1 mRNA is present in nine other breast tumor samples (from Brazilian and American patients) but absent in, or at exceedingly low levels in, the normal breast tissue corresponding to each cancer patient. RT-PCR analysis has also shown that the B18Ag1 transcript is not present in various normal tissues (including lymph node, myocardium and liver) and present at relatively low levels in PBMC and lung tissue. The presence of B18Ag1 mRNA in breast tumor samples, and its absence from normal breast tissue, has been confirmed by Northern blot analysis, as shown in Figure 2.

The differential expression of B18Ag1 in breast tumor tissue was also confirmed by RNase protection assays. Figure 3 shows the level of B18Ag1 mRNA in various tissue types as determined in four different RNase protection assays. Lanes 1-12 represent various normal breast tissue samples, lanes 13-25 represent various breast tumor samples; lanes 26-27 represent normal prostate samples; lanes 28-29 represent prostate tumor samples; lanes 30-32 represent colon tumor samples; lane 33 represents normal aorta; lane 34 represents normal small intestine; lane 35 represents normal skin, lane 36 represents normal lymph node; lane 37 represents normal ovary; lane 38 represents normal liver; lane 39 represents normal skeletal muscle; lane 40 represents a first normal stomach sample, lane 41 represents a second normal stomach sample; lane 42 represents a normal lung; lane 43 represents normal kidney; and lane 44 represents normal pancreas. Interexperimental comparison was facilitated by including a positive control RNA of known β -actin message abundance in each assay and normalizing the results of the different assays with respect to this positive control.

RT-PCR and Southern Blot analysis has shown the B18Ag1 locus to be present in human genomic DNA as a single copy endogenous retroviral element. A genomic clone of approximately 12-18 kb was isolated using the initial B18Ag1 sequence as a probe. Four additional subclones were also isolated by XbaI digestion. Additional retroviral sequences obtained from the ends of the XbaI digests of these clones (located as shown in Figure 4) are shown as SEQ ID NO:3 - SEQ ID NO:10,

where SEQ ID NO:3 shows the location of the sequence labeled 10 in Figure 4, SEQ ID NO:4 shows the location of the sequence labeled 11-29, SEQ ID NO:5 shows the location of the sequence labeled 3, SEQ ID NO:6 shows the location of the sequence labeled 6, SEQ ID NO:7 shows the location of the sequence labeled 12, SEQ ID NO:8 shows the location of the sequence labeled 13, SEQ ID NO:9 shows the location of the sequence labeled 14 and SEQ ID NO:10 shows the location of the sequence labeled 11-22.

Subsequent studies demonstrated that the 12-18 kb genomic clone contains a retroviral element of about 7.75 kb, as shown in Figures 5A and 5B. The sequence of this retroviral element is shown in SEQ ID NO:141. The numbered line at the top of Figure 5A represents the sense strand sequence of the retroviral genomic clone. The box below this line shows the position of selected restriction sites. The arrows depict the different overlapping clones used to sequence the retroviral element. The direction of the arrow shows whether the single-pass subclone sequence corresponded to the sense or anti-sense strand. Figure 5B is a schematic diagram of the retroviral element containing B18Ag1 depicting the organization of viral genes within the element. The open boxes correspond to predicted reading frames, starting with a methionine, found throughout the element. Each of the six likely reading frames is shown, as indicated to the left of the boxes, with frames 1-3 corresponding to those found on the sense strand.

Using the cDNA of SEQ ID NO:1 as a probe, a longer cDNA was obtained (SEQ ID NO:227) which contains minor nucleotide differences (less than 1%) compared to the genomic sequence shown in SEQ ID NO:141.

B. Preparation of cDNA Molecules Encoding Other Breast Tumor-Specific Polypeptides

Normal RNA and tumor RNA was prepared and mRNA was isolated and converted into cDNA using a (dT)₁₂AG anchored 3' primer, as described above. Differential display PCR was then executed using the randomly chosen primers of SEQ ID NOs:87-125. Amplification conditions were as noted above, and bands observed to be specific to the RNA fingerprint pattern of the tumor were cut out of a silver stained

gel, subcloned into either the T-vector (Novagen, Madison, WI) or the pCRII vector (Invitrogen, San Diego, CA) and sequenced. The sequences are provided in SEQ ID NO:11 - SEQ ID NO:86. Of the 79 sequences isolated, 67 were found to be novel (SEQ ID NOs:11-26 and 28-77) (*see also* Figures 6-20).

- 5 An extended DNA sequence (SEQ ID NO:290) for the antigen B15Ag1 (originally identified partial sequence provided in SEQ ID NO:27) was obtained in further studies. Comparison of the sequence of SEQ ID NO:290 with those in the gene bank as described above, revealed homology to the known human β -A activin gene. Further studies led to the isolation of the full-length cDNA sequence for the antigen
- 10 B21GT2 (also referred to as B311D; originally identified partial cDNA sequence provided in SEQ ID NOs:56). The full-length sequence is provided in SEQ ID NO:307, with the corresponding amino acid sequence being provided in SEQ ID NO:308. Further studies led to the isolation of a splice variant of B311D. The B311D clone of SEQ ID NO:316 was sequenced and a XhoI/NotI fragment from this clone was gel
- 15 purified and 32P-cDTP labeled by random priming for use as a probe for further screening to obtain additional B311D gene sequence. Two fractions of a human breast tumor cDNA bacterial library were screened using standard techniques. One of the clones isolated in this manner yielded additional sequence which includes a poly A+ tail. The determined cDNA sequence of this clone (referred to as B311D_BT1_1A) is
- 20 provided in SEQ ID NO:317. The sequences of SEQ ID NOs:316 and 317 were found to share identity over a 464 bp region, with the sequences diverging near the poly A+ sequence of SEQ ID NO:317.

- Subsequent studies identified an additional 146 sequences (SEQ ID NOs:142-289), of which 115 appeared to be novel (SEQ ID NOs:142, 143, 146-152,
- 25 154-166, 168-176, 178-192, 194-198, 200-204, 206, 207, 209-214, 216, 218, 219, 221-240, 243-245, 247, 250, 251, 253, 255, 257-266, 268, 269, 271-273, 275, 276, 278, 280, 281, 284, 288 and 291). To the best of the inventors' knowledge none of the previously identified sequences have heretofore been shown to be expressed at a greater level in human breast tumor tissue than in normal breast tissue.

In further studies, several different splice forms of the antigen B11Ag1 (also referred to as B305D) were isolated, with each of the various splice forms containing slightly different versions of the B11Ag1 coding frame. Splice junction sequences define individual exons which, in various patterns and arrangements, make up the various splice forms. Primers were designed to examine the expression pattern of each of the exons using RT-PCR as described below. Each exon was found to show the same expression pattern as the original B11Ag1 clone, with expression being breast tumor-, normal prostate- and normal testis-specific. The determined cDNA sequences for the isolated protein coding exons are provided in SEQ ID NOs:292-298, respectively. The predicted amino acid sequences corresponding to the sequences of SEQ ID NOs:292 and 298 are provided in SEQ ID NOs:299 and 300. Additional studies using rapid amplification of cDNA ends (RACE), a 5' specific primer to one of the splice forms of B11Ag1 provided above and a breast adenocarcinoma, led to the isolation of three additional, related, splice forms referred to as isoforms B11C-15, B11C-8 and B11C-9,16. The determined cDNA sequences for these isoforms are provided in SEQ ID NO: 301-303, with the corresponding predicted amino acid sequences being provided in SEQ ID NOs:304-306.

The protein coding region of B11C-15 (SEQ ID NO: 301; also referred to as B305D isoform C) was used as a query sequence in a BLASTN search of the Genbank DNA database. A match was found to a genomic clone from chromosome 21 (Accession no. AP001465). The pairwise alignments provided in the BLASTN output were used to identify the putative exon, or coding, sequence of the chromosome 21 sequence that corresponds to the B305D sequence. Based on the BlastN pairwise alignments, the following pieces of GenBank record AP001465 were put together: base pairs 67978-68499, 72870-72987, 73144-73335, 76085-76206, 77905-78085, 80520-80624, 87602-87633. This sequence was then aligned with the B305D isoform C sequence using the DNA Star Seqman program and excess sequence was deleted in such a way as to maintain the sequence most similar to B305D. The final edited form of the chromosome 21 sequence was 96.5% identical to B305D. This resulting edited sequence from chromosome 21 was then translated and found to contain no stop codons

other than the final stop codon in the same position as that for B305D. As with B305D, the chromosome 21 sequence (provided in SEQ ID NO: 325) encoded a protein (SEQ ID NO: 326) with 384 amino acids. An alignment of this protein with the B305D isoform C protein (SEQ ID NO: 304) showed 90% amino acid identity.

5 The cDNA sequence of B305D isoform C (SEQ ID NO: 301) was used to identify homologs by searching the High Throughput Genome Sequencing (HTGS) database (NCBI, National Institutes for Health, Bethesda, MD). Homologs were identified on Chromosome 2 (Clone ID 9838181), Chromosome 10 (Clone ID 10933022), Chromosome 15 (Clone ID 11560284). These homologs shared greater
10 than 90% identity with B305D isoform C at the nucleic acid level. All three of these homologs encode 384 amino acid ORFs that share greater than 90% identity with the amino acid sequence of SEQ ID NO: 304. Further searching of the GenBank database with the sequence of SEQ ID NO: 301 yielded a partial sequence homolog on Chromosome 22 (Clone ID 5931507). cDNA sequences for the Chromosome 2, 10, 15
15 and 22 homologs were constructed based on the homology with B305D isoform C and the conserved sequences at intron-exon junctions. The cDNA sequences for the Chromosome 22, 2, 15 and 10 homologs are provided in SEQ ID NO: 327-330, respectively, with the corresponding amino acid sequences being provided in SEQ ID NO: 331, 334, 333 and 332, respectively.

20 In subsequent studies on B305D isoform A (cDNA sequence provided in SEQ ID NO:292), the cDNA sequence (provided in SEQ ID NO:313) was found to contain an additional guanine residue at position 884, leading to a frameshift in the open reading frame. The determined DNA sequence of this ORF is provided in SEQ ID NO:314. This frameshift generates a protein sequence (provided in SEQ ID NO:315) of
25 293 amino acids that contains the C-terminal domain common to the other isoforms of B305D but that differs in the N-terminal region.

EXAMPLE 2

PREPARATION OF B18AG1 DNA FROM HUMAN GENOMIC DNA

This Example illustrates the preparation of B18Ag1 DNA by
5 amplification from human genomic DNA.

B18Ag1 DNA may be prepared from 250 ng human genomic DNA using
20 pmol of B18Ag1 specific primers, 500 pmol dNTPS and 1 unit of *Taq* DNA
polymerase (Perkin Elmer, Branchburg, NJ) using the following amplification
parameters: 94°C for 30 seconds denaturing, 30 seconds 60°C to 42°C touchdown
10 annealing in 2°C increments every two cycles and 72°C extension for 30 seconds. The
last increment (a 42°C annealing temperature) should cycle 25 times. Primers were
selected using computer analysis. Primers synthesized were B18Ag1-1, B18Ag1-2,
B18Ag1-3, and B18Ag1-4. Primer pairs that may be used are 1+3, 1+4, 2+3, and 2+4.

Following gel electrophoresis, the band corresponding to B18Ag1 DNA
15 may be excised and cloned into a suitable vector.

EXAMPLE 3

PREPARATION OF B18AG1 DNA FROM BREAST TUMOR cDNA

20 This Example illustrates the preparation of B18Ag1 DNA by
amplification from human breast tumor cDNA.

First strand cDNA is synthesized from RNA prepared from human breast
tumor tissue in a reaction mixture containing 500 ng poly A⁺ RNA, 200 pmol of the
primer (T)₁₂AG (*i.e.*, TTT TTT TTT TTT AG) (SEQ ID NO:130), 1X first strand
25 reverse transcriptase buffer, 6.7 mM DTT, 500 mmol dNTPs, and 1 unit AMV or
MMLV reverse transcriptase (from any supplier, such as Gibco-BRL (Grand Island,
NY)) in a final volume of 30 µl. After first strand synthesis, the cDNA is diluted
approximately 25 fold and 1 µl is used for amplification as described in Example 2.
While some primer pairs can result in a heterogeneous population of transcripts, the
30 primers B18Ag1-2 (5'ATG GCT ATT TTC GGG GGC TGA CA) (SEQ ID NO:126)

and B18Ag1-3 (5'CCG GTA TCT CCT CGT GGG TAT T) (SEQ ID NO:127) yield a single 151 bp amplification product.

EXAMPLE 4

5 IDENTIFICATION OF B-CELL AND T-CELL EPITOPES OF B18Ag1

This Example illustrates the identification of B18Ag1 epitopes.

The B18Ag1 sequence can be screened using a variety of computer algorithms. To determine B-cell epitopes, the sequence can be screened for
10 hydrophobicity and hydrophilicity values using the method of Hopp, *Prog. Clin. Biol. Res. 172B*:367-77 (1985) or, alternatively, Cease et al., *J. Exp. Med.* 164:1779-84 (1986) or Spouge et al., *J. Immunol.* 138:204-12 (1987). Additional Class II MHC (antibody or B-cell) epitopes can be predicted using programs such as AMPHI (e.g., Margalit et al., *J. Immunol.* 138:2213 (1987)) or the methods of Rothbard and Taylor
15 (e.g., *EMBO J.* 7:93 (1988)).

Once peptides (15-20 amino acids long) are identified using these techniques, individual peptides can be synthesized using automated peptide synthesis equipment (available from manufacturers such as Perkin Elmer/Applied Biosystems Division, Foster City, CA) and techniques such as Merrifield synthesis. Following
20 synthesis, the peptides can be used to screen sera harvested from either normal or breast cancer patients to determine whether patients with breast cancer possess antibodies reactive with the peptides. Presence of such antibodies in breast cancer patient would confirm the immunogenicity of the specific B-cell epitope in question. The peptides can also be tested for their ability to generate a serologic or humoral immune in animals
25 (mice, rats, rabbits, chimps etc.) following immunization *in vivo*. Generation of a peptide-specific antiserum following such immunization further confirms the immunogenicity of the specific B-cell epitope in question.

To identify T-cell epitopes, the B18Ag1 sequence can be screened using different computer algorithms which are useful in identifying 8-10 amino acid motifs
30 within the B18Ag1 sequence which are capable of binding to HLA Class I MHC

molecules. (see, e.g., Rammensee et al., *Immunogenetics* 41:178-228 (1995)). Following synthesis such peptides can be tested for their ability to bind to class I MHC using standard binding assays (e.g., Sette et al., *J. Immunol.* 153:5586-92 (1994)) and more importantly can be tested for their ability to generate antigen reactive cytotoxic T-cells following *in vitro* stimulation of patient or normal peripheral mononuclear cells using, for example, the methods of Bakker et al., *Cancer Res.* 55:5330-34 (1995); Visseren et al., *J. Immunol.* 154:3991-98 (1995); Kawakami et al., *J. Immunol.* 154:3961-68 (1995); and Kast et al., *J. Immunol.* 152:3904-12 (1994). Successful *in vitro* generation of T-cells capable of killing autologous (bearing the same Class I MHC molecules) tumor cells following *in vitro* peptide stimulation further confirms the immunogenicity of the B18Ag1 antigen. Furthermore, such peptides may be used to generate murine peptide and B18Ag1 reactive cytotoxic T-cells following *in vivo* immunization in mice rendered transgenic for expression of a particular human MHC Class I haplotype (Vitiello et al., *J. Exp. Med.* 173:1007-15 (1991)).

A representative list of predicted B18Ag1 B-cell and T-cell epitopes, broken down according to predicted HLA Class I MHC binding antigen, is shown below:

Predicted Th Motifs (B-cell epitopes) (SEQ ID NOS.: 131-133)

SSGGRTFDDFHRYLLVGI
 QGAAQKPINLSKXIEVVQGHDE
 SPGVFLEHLQEAYRIYTPFDLSA

Predicted HLA A2.1 Motifs (T-cell epitopes) (SEQ ID NOS.: 134-140)

YLLVGIQGA
 GAAQKPINL
 NLSKXIEVV
 EVVQGHDES
 HLQEAYRIY
 NLAQVAQAA
 FVAQAAPDS

EXAMPLE 5

IDENTIFICATION OF T-CELL EPITOPES OF B11Ag1

This Example illustrates the identification of B11Ag1 (also referred to as B305D) epitopes. Four peptides, referred to as B11-8, B11-1, B11-5 and B11-12 (SEQ ID NOs:309-312, respectfully) were derived from the B11Ag1 gene.

Human CD8 T cells were primed *in vitro* to the peptide B11-8 using dendritic cells according to the protocol of Van Tsai et al. (*Critical Reviews in Immunology* 18:65-75, 1998). The resulting CD8 T cell cultures were tested for their ability to recognize the B11-8 peptide or a negative control peptide, presented by the B-LCL line, JY. Briefly, T cells were incubated with autologous monocytes in the presence of 10 ug/ml peptide, 10 ng/ml IL-7 and 10 ug/ml IL-2, and assayed for their ability to specifically lyse target cells in a standard 51-Cr release assay. As shown in Fig. 22, the bulk culture line demonstrated strong recognition of the B11-8 peptide with weaker recognition of the peptide B11-1.

A clone from this CTL line was isolated following rapid expansion using the monoclonal antibody OKT3 and human IL-2. As shown in Fig. 23, this clone (referred to as A1), in addition to being able to recognize specific peptide, recognized JY LCL transduced with the B11Ag1 gene. This data demonstrates that B11-8 is a naturally processed epitope of the B11Ag1 gene. In addition these T cells were further found to recognize and lyse, in an HLA-A2 restricted manner, an established tumor cell line naturally expressing B11Ag1 (Fig. 24). The T cells strongly recognize a lung adenocarcinoma (LT-140-22) naturally expressing B11Ag1 transduced with HLA-A2, as well as an A2+ breast carcinoma (CAMA-1) transduced with B11Ag1, but not untransduced lines or another negative tumor line (SW620).

These data clearly demonstrate that these human T cells recognize not only B11-specific peptides but also transduced cells, as well as naturally expressing tumor lines.

CTL lines raised against the antigens B11-5 and B11-12, using the procedures described above, were found to recognize corresponding peptide-coated targets.

EXAMPLE 6

CHARACTERIZATION OF BREAST TUMOR GENES DISCOVERED BY
DIFFERENTIAL DISPLAY PCR

5 The specificity and sensitivity of the breast tumor genes discovered by differential display PCR were determined using RT-PCR. This procedure enabled the rapid evaluation of breast tumor gene mRNA expression semiquantitatively without using large amounts of RNA. Using gene specific primers, mRNA expression levels in a variety of tissues were examined, including 8 breast tumors, 5 normal breasts, 2
10 prostate tumors, 2 colon tumors, 1 lung tumor, and 14 other normal adult human tissues, including normal prostate, colon, kidney, liver, lung, ovary, pancreas, skeletal muscle, skin, stomach and testes.

 To ensure the semiquantitative nature of the RT-PCR, β -actin was used as internal control for each of the tissues examined. Serial dilutions of the first strand
15 cDNAs were prepared and RT-PCR assays performed using β -actin specific primers. A dilution was then selected that enabled the linear range amplification of β -actin template, and which was sensitive enough to reflect the difference in the initial copy number. Using this condition, the β -actin levels were determined for each reverse transcription reaction from each tissue. DNA contamination was minimized by DNase
20 treatment and by assuring a negative result when using first strand cDNA that was prepared without adding reverse transcriptase.

 Using gene specific primers, the mRNA expression levels were determined in a variety of tissues. To date, 38 genes have been successfully examined by RT-PCR, five of which exhibit good specificity and sensitivity for breast tumors
25 (B15AG-1, B31GA1b, B38GA2a, B11A1a and B18AG1a). Figures 21A and 21B depict the results for three of these genes: B15AG-1 (SEQ ID NO:27), B31GA1b (SEQ ID NO:148) and B38GA2a (SEQ ID NO:157). Table I summarizes the expression level of all the genes tested in normal breast tissue and breast tumors, and also in other tissues.

TABLE I

Percentage of Breast Cancer Antigens that are Expressed in Various Tissues

5	Breast Tissues	Over-expressed in Breast Tumors	84%
		Equally Expressed in Normals and Tumor	16%
10	Other Tissues	Over-expressed in Breast Tumors but not in any Normal Tissues	9%
		Over-expressed in Breast Tumors but Expressed in Some Normal Tissues	30%
15		Over-expressed in Breast Tumors but Equally Expressed in All Other Tissues	61%

20

EXAMPLE 7

PREPARATION AND CHARACTERIZATION OF ANTIBODIES AGAINST BREAST TUMOR
POLYPEPTIDES

Polyclonal antibodies against the breast tumor antigen B305D were
25 prepared as follows.

The breast tumor antigen expressed in an *E. coli* recombinant expression system was grown overnight in LB broth with the appropriate antibiotics at 37 °C in a shaking incubator. The next morning, 10 ml of the overnight culture was added to 500 ml to 2x YT plus appropriate antibiotics in a 2L-baffled Erlenmeyer flask. When the
30 Optical Density (at 560 nm) of the culture reached 0.4-0.6, the cells were induced with IPTG (1 mM). Four hours after induction with IPTG, the cells were harvested by centrifugation. The cells were then washed with phosphate buffered saline and centrifuged again. The supernatant was discarded and the cells were either frozen for future use or immediately processed. Twenty ml of lysis buffer was added to the cell
35 pellets and vortexed. To break open the *E. coli* cells, this mixture was then run through the French Press at a pressure of 16,000 psi. The cells were then centrifuged again and

the supernatant and pellet were checked by SDS-PAGE for the partitioning of the recombinant protein. For proteins that localized to the cell pellet, the pellet was resuspended in 10 mM Tris pH 8.0, 1% CHAPS and the inclusion body pellet was washed and centrifuged again. This procedure was repeated twice more. The washed
5 inclusion body pellet was solubilized with either 8 M urea or 6 M guanidine HCl containing 10 mM Tris pH 8.0 plus 10 mM imidazole. The solubilized protein was added to 5 ml of nickel-chelate resin (Qiagen) and incubated for 45 min to 1 hour at room temperature with continuous agitation. After incubation, the resin and protein mixture were poured through a disposable column and the flow through was collected.
10 The column was then washed with 10-20 column volumes of the solubilization buffer. The antigen was then eluted from the column using 8M urea, 10 mM Tris pH 8.0 and 300 mM imidazole and collected in 3 ml fractions. A SDS-PAGE gel was run to determine which fractions to pool for further purification.

As a final purification step, a strong anion exchange resin such as
15 HiPrepQ (Biorad) was equilibrated with the appropriate buffer and the pooled fractions from above were loaded onto the column. Antigen was eluted off the column with a increasing salt gradient. Fractions were collected as the column was run and another SDS-PAGE gel was run to determine which fractions from the column to pool. The pooled fractions were dialyzed against 10 mM Tris pH 8.0. The protein was then vialled
20 after filtration through a 0.22 micron filter and the antigens were frozen until needed for immunization.

Four hundred micrograms of B305D antigen was combined with 100 micrograms of muramyldipeptide (MDP). Every four weeks rabbits were boosted with 100 micrograms mixed with an equal volume of Incomplete Freund's Adjuvant (IFA).
25 Seven days following each boost, the animal was bled. Sera was generated by incubating the blood at 4 °C for 12-24 hours followed by centrifugation.

Ninety-six well plates were coated with B305D antigen by incubating with 50 microliters (typically 1 microgram) of recombinant protein at 4 °C for 20 hours. 250 microliters of BSA blocking buffer was added to the wells and incubated at room
30 temperature for 2 hours. Plates were washed 6 times with PBS/0.01% Tween. Rabbit

sera was diluted in PBS. Fifty microliters of diluted sera was added to each well and incubated at room temperature for 30 min. Plates were washed as described above before 50 microliters of goat anti-rabbit horse radish peroxidase (HRP) at a 1:10000 dilution was added and incubated at room temperature for 30 min. Plates were again
5 washed as described above and 100 microliters of TMB microwell peroxidase substrate was added to each well. Following a 15 min incubation in the dark at room temperature, the colorimetric reaction was stopped with 100 microliters of 1N H₂SO₄ and read immediately at 450 nm. The polyclonal antibodies showed immunoreactivity to B305D.

10 Immunohistochemical (IHC) analysis of B305D expression in breast cancer and normal breast specimens was performed as follows. Paraffin-embedded formal fixed tissue was sliced into 8 micron sections. Steam heat induced epitope retrieval (SHIER) in 0.1 M sodium citrate buffer (pH 6.0) was used for optimal staining conditions. Sections were incubated with 10% serum/PBS for 5 minutes. Primary
15 antibody was added to each section for 25 min at indicated concentrations followed by a 25 min incubation with either an anti-rabbit or anti-mouse biotinylated antibody. Endogenous peroxidase activity was blocked by three 1.5 min incubations with hydrogen peroxide. The avidin biotin complex/horseradish peroxidase (ABC/HRP) systems was used along with DAB chromagen to visualize antigen expression. Slides
20 were counterstained with hematoxylin. B305D expression was detected in both breast tumor and normal breast tissue. However, the intensity of staining was much less in normal samples than in tumor samples and surface expression of B305D was observed only in breast tumor tissues.

A summary of real-time PCR and immunohistochemical analysis of
25 B305D expression in an extensive panel of normal tissues is presented in Table II below. These results demonstrate minimal expression of B305D in testis, inconclusive results in gall bladder, and no detection in all other tissues tested.

TABLE II

mRNA	IHC staining	Tissue type	Summary
Moderately positive	Positive	Testis	Nuclear staining of small minority of spermatids; spermatozoa negative; seminoma negative
Negative	Negative	Thymus	No expression
N/A	Negative	Artery	No expression
Negative	Negative	Skeletal muscle	No expression
Negative	Positive (weak staining)	Small bowel	No expression
Negative	Positive (weak staining)	Ovary	No expression
Negative		Pituitary	No expression
Negative	Positive (weak staining)	Stomach	No expression
Negative	Negative	Spinal cord	No expression
Negative	Negative	Spleen	No expression
Negative	Negative	Ureter	No expression
N/A	Negative	Gall bladder	Inconclusive
N/A	Negative	Placenta	No expression
Negative	Negative	Thyroid	No expression
Negative	Negative	Heart	No expression
Negative	Negative	Kidney	No expression
Negative	Negative	Liver	No expression
Negative	Negative	Brain-cerebellum	No expression
Negative	Negative	Colon	No expression
Negative	Negative	Skin	No expression
Negative	Negative	Bone marrow	No expression
N/A	Negative	Parathyroid	No expression
Negative	Negative	Lung	No expression
Negative	Negative	Esophagus	No expression
Negative	Positive (weak staining)	Uterus	No expression
Negative	Negative	Adrenal	No expression
Negative	Negative	Pancreas	No expression
N/A	Negative	Lymph node	No expression
Negative	Negative	Brain-cortex	No expression
N/A	Negative	Fallopian tube	No expression
Negative	Positive (weak staining)	Bladder	No expression
Negative	N/A	Bone	No expression
Negative	N/A	Salivary gland	No expression

Negative	N/A	Activated PBMC	No expression
Negative	N/A	Resting PBMC	No expression
Negative	N/A	Trachea	No expression
Negative	N/A	Vena cava	No expression
Negative	N/A	Retina	No expression
Negative	N/A	Cartilage	No expression

EXAMPLE 8

PROTEIN EXPRESSION OF BREAST TUMOR ANTIGENS

5 This example describes the expression and purification of the breast tumor antigen B305D in *E. coli* and in mammalian cells.

Expression of B305D isoform C-15 (SEQ ID NO:301; translated to 384 amino acids) in *E. coli* was achieved by cloning the open reading frame of B305D isoform C-15 downstream of the first 30 amino acids of the *M. tuberculosis* antigen Ra12 (SEQ ID NO:318) in pET17b. First, the internal EcoRI site in the B305D ORF was mutated without changing the protein sequence so that the gene could be cloned at the EcoRI site with Ra12. The PCR primers used for site-directed mutagenesis are shown in SEQ ID NO:319 (referred to as AW012) and SEQ ID NO:320 (referred to as AW013). The ORF of EcoRI site-modified B305D was then amplified by PCR using
10 the primers AW014 (SEQ ID NO:321) and AW015 (SEQ ID NO:322). The PCR product was digested with EcoRI and ligated to the Ra12/pET17b vector at the EcoRI site. The sequence of the resulting fusion construct (referred to as Ra12mB11C) was confirmed by DNA sequencing. The determined cDNA sequence for the fusion construct is provided in SEQ ID NO:323, with the amino acid sequence being provided
15 in SEQ ID NO:324.

The fusion construct was transformed into BL21(DE3)CodonPlus-RIL *E. coli* (Stratagene) and grown overnight in LB broth with kanamycin. The resulting culture was induced with IPTG. Protein was transferred to PVDF membrane and blocked with 5% non-fat milk (in PBS-Tween buffer), washed three times and
25 incubated with mouse anti-His tag antibody (Clontech) for 1 hour. The membrane was washed 3 times and probed with HRP-Protein A (Zymed) for 30 min. Finally, the

membrane was washed 3 times and developed with ECL (Amersham). Expression was detected by Western blot.

For recombinant expression in mammalian cells, B305D isoform C-15 (SEQ ID NO:301; translated to 384 amino acids) was subcloned into the mammalian expression vectors pCEP4 and pcDNA3.1 (Invitrogen). These constructs were transfected into HEK293 cells (ATCC) using Fugene 6 reagent (Roche). Briefly, the HEK cells were plated at a density of 100,000 cells/ml in DMEM (Gibco) containing 10% FBS (Hyclone) and grown overnight. The following day, 2 ul of Fugene 6 was added to 100 ul of DMEM containing no FBS and incubated for 15 minutes at room temperature. The Fugene 6/DMEM mixture was added to 1 ug of B305D/pCEP4 or B305D/pcDNA plasmid DNA and incubated for 15 minutes at room temperature. The Fugene/DNA mix was then added to the HEK293 cells and incubated for 48-72 hours at 37 °C with 7% CO₂. Cells were rinsed with PBS, the collected and pelleted by centrifugation.

For Western blot analysis, whole cell lysates were generated by incubating the cells in Triton-X100 containing lysis buffer for 30 minutes on ice. Lysates were then cleared by centrifugation at 10,000 rpm for 5 minutes at 4 °C. Samples were diluted with SDS_PAGE loading buffer containing beta-mercaptoethanol, and boiled for 10 minutes prior to loading the SDS_PAGE gel. Proteins were transferred to nitrocellulose and probed using Protein A purified anti-B305D rabbit polyclonal sera (prepared as described above) at a concentration of 1 ug/ml. The blot was revealed with a goat anti-rabbit Ig coupled to HRP followed by incubation in ECL substrate. Expression of B305D was detected in the the HEK293 lysates transfected with B305D, but not in control HEK293 cells transfected with vector alone.

For FACS analysis, cells were washed further with ice cold staining buffer and then incubated with a 1:100 dilution of a goat anti-rabbit Ig (H+L)-FITC reagent (Southern Biotechnology) for 30 minutes on ice. Following 3 washes, the cells were resuspended in staining buffer containing Propidium Iodide (PI), a vital stain that

allows for identification of permeable cells, and then analyzed by FACS. The FACS analysis showed surface expression of B305D protein.

From the foregoing it will be appreciated that, although specific
5 embodiments of the invention have been described herein for purposes of illustration, various modifications may be made without deviating from the spirit and scope of the invention. Accordingly, the invention is not limited except as by the appended claims.

CLAIMS

What is Claimed:

1. An isolated polynucleotide comprising a sequence selected from the group consisting of:

(a) sequences provided in SEQ ID NO: 1, 3-86, 142-298, 301-303, 307, 313, 314, 316, 317, 325 and 327-330;

(b) complements of the sequences provided in SEQ ID NO: 1, 3-86, 142-298, 301-303, 307, 313, 314, 316, 317, 325 and 327-330;

(c) sequences consisting of at least 20 contiguous residues of a sequence provided in SEQ ID NO: 1, 3-86, 142-298, 301-303, 307, 313, 314, 316, 317, 325 and 327-330;

(d) sequences that hybridize to a sequence provided in SEQ ID NO: 1, 3-86, 142-298, 301-303, 307, 313, 314, 316, 317, 325 and 327-330, under moderately stringent conditions;

(e) sequences having at least 75% identity to a sequence of SEQ ID NO: 1, 3-86, 142-298, 301-303, 307, 313, 314, 316, 317, 325 and 327-330;

(f) sequences having at least 90% identity to a sequence of SEQ ID NO: 1, 3-86, 142-298, 301-303, 307, 313, 314, 316, 317, 325 and 327-330; and

(g) degenerate variants of a sequence provided in SEQ ID NO: 1, 3-86, 142-298, 301-303, 307, 313, 314, 316, 317, 325 and 327-330.

2. An isolated polypeptide comprising an amino acid sequence selected from the group consisting of:

(a) SEQ ID NO: 299, 300, 304-306, 308-312, 314, 326 and 331-334;

(b) sequences encoded by a polynucleotide of claim 1;

(c) sequences having at least 70% identity to a sequence encoded by a polynucleotide of claim 1; and

(d) sequences having at least 90% identity to a sequence encoded by a polynucleotide of claim 1.

3. An expression vector comprising a polynucleotide of claim 1 operably linked to an expression control sequence.

4. A host cell transformed or transfected with an expression vector according to claim 3.

5. An isolated antibody, or antigen-binding fragment thereof, that specifically binds to a polypeptide of claim 2.

6. A method for detecting the presence of a cancer in a patient, comprising the steps of:

- (a) obtaining a biological sample from the patient;
- (b) contacting the biological sample with a binding agent that binds to a polypeptide of claim 2;
- (c) detecting in the sample an amount of polypeptide that binds to the binding agent; and
- (d) comparing the amount of polypeptide to a predetermined cut-off value and therefrom determining the presence of a cancer in the patient.

7. A fusion protein comprising at least one polypeptide according to claim 2.

8. An oligonucleotide that hybridizes to a sequence recited in SEQ ID NO: 1, 3-86, 142-298, 301-303, 307, 313, 314, 316, 317, 325 and 327-330 under moderately stringent conditions.

9. A method for stimulating and/or expanding T cells specific for a tumor protein, comprising contacting T cells with at least one component selected from the group consisting of:

- (a) polypeptides according to claim 2;
- (b) polynucleotides according to claim 1; and
- (c) antigen-presenting cells that express a polypeptide according to claim 2,

under conditions and for a time sufficient to permit the stimulation and/or expansion of T cells.

10. An isolated T cell population, comprising T cells prepared according to the method of claim 9.

11. A composition comprising a first component selected from the group consisting of physiologically acceptable carriers and immunostimulants, and a second component selected from the group consisting of:

- (a) polypeptides according to claim 2;
- (b) polynucleotides according to claim 1;
- (c) antibodies according to claim 5;
- (d) fusion proteins according to claim 7;
- (e) T cell populations according to claim 10; and
- (f) antigen presenting cells that express a polypeptide according to claim 2.

12. A method for stimulating an immune response in a patient, comprising administering to the patient a composition of claim 11.

13. A method for the treatment of a cancer in a patient, comprising administering to the patient a composition of claim 11.

14. A method for determining the presence of a cancer in a patient, comprising the steps of:

- (a) obtaining a biological sample from the patient;
- (b) contacting the biological sample with an oligonucleotide according to claim 8;
- (c) detecting in the sample an amount of a polynucleotide that hybridizes to the oligonucleotide; and
- (d) compare the amount of polynucleotide that hybridizes to the oligonucleotide to a predetermined cut-off value, and therefrom determining the presence of the cancer in the patient.

15. A diagnostic kit comprising at least one oligonucleotide according to claim 8.

16. A diagnostic kit comprising at least one antibody according to claim 5 and a detection reagent, wherein the detection reagent comprises a reporter group.

17. A method for inhibiting the development of a cancer in a patient, comprising the steps of:

- (a) incubating CD4⁺ and/or CD8⁺ T cells isolated from a patient with at least one component selected from the group consisting of: (i) polypeptides according to claim 2; (ii) polynucleotides according to claim 1; and (iii) antigen presenting cells that express a polypeptide of claim 2, such that T cell proliferate;
 - (b) administering to the patient an effective amount of the proliferated T cells,
- and thereby inhibiting the development of a cancer in the patient.




Fig. 1

2/19




Fig. 2




Fig. 3

GENOMIC CLONE MAP





Fig. 4



6/19

NUCLEOTIDE SEQUENCE OF THE REPRESENTATIVE
BREAST-TUMOR SPECIFIC cDNA B18Ag1

TTA GAG ACC CAA TTG GGA CCT AAT TGG GAC CCA AAT TTC TCA AGT GGA	48
Leu Glu Thr Gln Leu Gly Pro Asn Trp Asp Pro Asn Phe Ser Ser Gly	
1 5 10 15	
GGG AGA ACT TTT GAC GAT TTC CAC CGG TAT CTC CTC GTG GGT ATT CAG	96
Gly Arg Thr Phe Asp Asp Phe His Arg Tyr Leu Leu Val Gly Ile Gln	
20 25 30	
GGA GCT GCC CAG AAA CCT ATA AAC TTG TCT AAG GCG ATT GAA GTC GTC	144
Gly Ala Ala Gln Lys Pro Ile Asn Leu Ser Lys Ala Ile Glu Val Val	
35 40 45	
CAG GGG CAT GAT GAG TCA CCA GGA GTG TTT TTA GAG CAC CTC CAG GAG	192
Gln Gly His Asp Glu Ser Pro Gly Val Phe Leu Glu His Leu Gln Glu	
50 55 60	
GCT TAT CGG ATT TAC ACC CCT TTT GAC CTG GCA GCC CCC GAA AAT AGC	240
Ala Tyr Arg Ile Tyr Thr Pro Phe Asp Leu Ala Ala Pro Glu Asn Ser	
65 70 75 80	
CAT GCT CTT AAT TTG GCA TTT GTG GCT CAG GCA GCC CCA GAT AGT AAA	288
His Ala Leu Asn Leu Ala Phe Val Ala Gln Ala Ala Pro Asp Ser Lys	
85 90 95	
AGG AAA CTC CAA AAA CTA GAG GGA TTT TGC TGG AAT GAA TAC CAG TCA	336
Arg Lys Leu Gln Lys Leu Glu Gly Phe Cys Trp Asn Glu Tyr Gln Ser	
100 105 110	
GCT TTT AGA GAT AGC CTA AAA GGT TTT	363
Ala Phe Arg Asp Ser Leu Lys Gly Phe	
115 120	

Fig. 6

NUCLEOTIDE SEQUENCE OF THE REPRESENTATIVE
BREAST-TUMOR SPECIFIC cDNA B17Ag1

GC TGGGCACAGT GGCTCATACC TGTAATCCTG ACCGTTTCAG AGGCTCAGGT	60
CG CTTGAGCCCA AGATTTCAAG ACTAGTCTGG GTAACATAGT GAGACCCTAT	120
AA AAATAAAAAA ATGAGCCTGG TGTAGTGGCA CACACCAGCT GAGGAGGGAG	180
CT AGGAGA	196

Fig. 7

7/19

NUCLEOTIDE SEQUENCE OF THE REPRESENTATIVE
BREAST-TUMOR SPECIFIC cDNA B17Ag2

GC TTGGGGGCTC TGA CTAGAAA TTCAAGGAAC CTGGGATTCA AGTCCA ACTG 60
AC TTACTACTGTG GNC TCCAATA AACTGCTTCT TTCCTATTCC CTCTCTATTA 120
AA GGAAAACGAT GTCTGTGTAT AGCCAAGTCA GNTATCCTAA AAGGAGATAC 180
AT TAAATATCAG AATGTAAAAC CTGGGAACCA GGTTC C CAGC CTGGGATTAA 240
CA AGAAGACTGA ACAGTACTAC TGTGAAAAGC CCGAAGNGGC AATATGTTC A 300
TT GAAGGATGGC TGGGAGAATG AATGCTCTGT CCCCAGTCC CAAGCTCACT 360
CT CCTTTATAGC CTAGGAGA 388

*Fig. 8*NUCLEOTIDE SEQUENCE OF THE REPRESENTATIVE
BREAST-TUMOR SPECIFIC cDNA B13Ag2a

GC CTATAATCAT GTTTCATT ATTTTCACAT TTTATTAACC AATTTCTGTT 60
AA AATATGAGGG AAATATATGA AACAGGGAGG CAATGTT CAG ATAATTGATC 120
TG ATTTCTACAT CAGATGCTCT TTCCTTTCCT GTTATTTC TTTTATTTT C 180
GG TCGAATGTAA TAGCTTTGTT TCAAGAGAGA GTTTTGGCAG TTTCTGTAGC 240
CT GCTCATGTCT CCAGGCATCT ATTTGCACTT TAGGAGGTGT CGTGGGAGAC 300
CT ATTTTTTCCA TATTGGGCA ACTACTA 337

Fig. 9

8/19

NUCLEOTIDE SEQUENCE OF THE REPRESENTATIVE
BREAST-TUMOR SPECIFIC cDNA B13Ag1b

GC CATACAGTGC CTTTCCATTT ATTTAACCCC CACCTGAACG GCATAAACTG 60
GC TGGTGTTTTT TACTGTAAAC AATAAGGAGA CTTTGCTCTT CATTTAAACC 120
AT TTCATATTTT ACGCTCGAGG GTTTTACCG GTTCCTTTT ACACCTCTTA 180
TT TAAGTCGTTT GGAACAAGAT ATTTTTCCTT TCCTGGCAGC TTTTAACATT 240
TT TGTGCTGGG GGAAGTCTGG TCACTGTTT TCACAGTTGC AAATCAAGGC 300
CC AAGAAAAAAA AATTTTTTTG TTTTATTTGA AACTGGACCG GATAAACGGT 360
CG GCTGCTGTAT ATAGTTTTAA ATGGTTTATT GCACCTCCTT AAGTTGCACT 420
GG GGGGNTTTTG NATAGAAAGT NTTTANTCAC ANAGTCACAG GGAAGTTTNT 480
NA CTGAGCTAAA AAGGGCTGNT TTTGGGTGG GGCAGATGA AGGCTCACAG 540
TC TCTTAGAGGG GGAAGTCT A 571

Fig. 10

9/19

NUCLEOTIDE SEQUENCE OF THE REPRESENTATIVE
BREAST-TUMOR SPECIFIC cDNA B13Ag1a

TA ATAACTTAAA TATATTTTGA TCACCCACTG GGGTGATAAG ACAATAGATA 60
TT TCCAAAAAGC ATAAAACCAA AGTATCATAC CAAACCAAAT TCATACTGCT 120
CC GCACTGAAAC TTCACCTTCT AACTGTCTAC CTAACCAAAT TCTACCTTC 180
GG TGGGTGCTCA CTACTCTTTT TTTTTTTTTT TTTNTTTTGG AGATGGAGTC 240
CA GCCCAGGGGT GGAGTACAAT GGCACAACCT CAGCTCACTG NAACCTCCGC 300
TT CATGAGATTC TCCTGNTTCA GCCTTCCCAG TAGCTGGGAC TACAGGTGTG 360
TG CCTGGNTAAT CTTTTTTNGT TTTNNGGTAG AGATGGGGGT TTTACATGTT 420
TG GTNTCGAACT CCTGACCTCA AGTGATCCAC CCACCTCAGG CTCCCAAAGT 480
TA CAGACATGAG CCACTGNGCC CAGNCTGGT GCATGCTCAC TTCTCTAGGC 540

Fig. 11

10/19

NUCLEOTIDE SEQUENCE OF THE REPRESENTATIVE
BREAST-TUMOR SPECIFIC cDNA B11Ag1

TG CACATGCAGA ATATTCTATC GGTACTTCAG CTATTACTCA TTTTGATGGC 60
AG CCTATCCTCA AGATGAGTAT TTAGAAAGAA TTGATTTAGC GATAGACCAA 120
GC ACTCTGACTA CACGAAATTG TTCAGATGTG ATGGATTTAT GACAGTTGAT 180
GA GATTATTAAG TGATTATTTT AAAGGGAATC CATTAATTCC AGAATATCTT 240
TC AAGATGATAT AGAAATAGAA CAGAAAGAGA CTACAAATGA AGATGTATCA 300
TA TTGAAGAGCC TATAGTAGAA AATGAATTAG CTGCATTTAT TAGCCTTACA 360
TT TTCCTGATGA ATCTTATATT CAGCCATCGA CATAGCATTG CCTGATGGGC 420
GA ATAATAGAAA CTGGGTGCGG GGCTATTGAT GAATTCATCC NCAGTAAATT 480
AC AAAATATAAC TCGATTGCAT TTGGATGATG GAATACTAAA TCTGGCAAAA 540
GG AGCTACTAGT AACCTCTCTT TTTGAGATGC AAAATTTTCT TTTAGGGTTT 600
CT ACTTTACGGA TATTGGAGCA TAACGGGA 638

Fig. 12

11/19

NUCLEOTIDE SEQUENCE OF THE REPRESENTATIVE
BREAST-TUMOR SPECIFIC cDNA B3CA3c

ACTGATGGAT GTCGCCGGAG GCGAGGGGCC TTATCTGATG CTCGGCTGCC TGTCGTGAT 60
GTGCGCGGCG ATTGGGCTGT TTATCTCAA CACCGCCACG GCGGTGCTGA TGGCGCCTAT 120
TGCCTTAGCG GCGGCGAAGT CAATGGGCGT CTCACCCTAT CCTTTGCCA TGGTGGTGGC 180
GATGGCGGCT TCGGCGGCGT TTATGACCCC GGTCTCCTCG CCGGTTAACA CCCTGGTGCT 240
TGGCCCTGGC AAGTACTCAT TTAGCGATTT TGCAAAATA GGCCTG 286

*Fig. 13*NUCLEOTIDE SEQUENCE OF THE REPRESENTATIVE
BREAST-TUMOR SPECIFIC cDNA B9CG1

AG CAGCCCCCTTCTCAATTT CATCTGTAC TACCCTGGTG TAGTATCTCA 60
CA TTTTATAGC CTCCTCCCTG GTCTGTCTTT TGATTTTCCT GCCTGTAATC 120
AC ATAAGTCAA GTAAACATTT CTAAAGTGTG GTTATGCTCA TGCACTCCT 180
AA ATAGTTTCCA TTACCGTCTT AATAAAATTC GGATTTGTTT TTNCTATTN 240
CA CCTATGACCG AA 262

Fig. 14

12/19

NUCLEOTIDE SEQUENCE OF THE REPRESENTATIVE
BREAST-TUMOR SPECIFIC cDNA B9CG3

AG CAAAGCCAGT GGT TTGAGCT CTCTACTGTG TAAACTCCTA AACCAAGGCC 60
TA AATGGTGGCA GGATTTTAT TATAAACATG TACCATGCA AATTCCTAT 120
GA TATATTCTT TACATTTAAA CAATAAAAAT AATCTATTTT TAAAAGCCTA 180
AG TTAGGTAAGA GTGTTTAATG AGAGGGTATA AGGTATAAAT CACCAGTCAA 240
TG CCTATGACCG A 261

*Fig. 15*NUCLEOTIDE SEQUENCE OF THE REPRESENTATIVE
BREAST-TUMOR SPECIFIC cDNA B2CA2

CGACGTCGGT AAAATCGGAC ATGAAGCCAC CGCTGGTCTT TTCGTCCGAG CGATAGGCGC 60
CGGCCAGCCA GCGGAACGGT TGCCCGGATG GCGAAGCGAG CCGGAGTTCT TCGGACTGAG 120
TATGAATCTT GTTGTGAAAA TACTCGCCGC CTTCGTTCTGA CGACGTCGCG TCGAAATCTT 180
AATCATGGTT GAGCCGGATG CTGCCCCGA AGCCCT 276

Fig. 16

13/19

NUCLEOTIDE SEQUENCE OF THE REPRESENTATIVE
BREAST-TUMOR SPECIFIC cDNA B3CA1

CCCAGGTCAA CCAGGCTGCA ACACGCAGGT CCTTGGATTG GGCACGAAGC AGCGCTTCGC 60
TGTTTTCCAG GATTTTCAAC CAGTCGGTCT GGCCGTTCCTC ATGGAGCGAG AGCGCCTTGC 120
CCAGCTCATT TTCCAGCGCC TCGTATTCGC TGGAAAAACG CACATCCTCA CCCGCAAAGA 180
CATCCTTTGA AATCGGCTGT TCCGCGAGTT CCAGATANTG CGAGGAGAGC TTGCTCGAAT 240
AGGTCATCCT AACCCTTCAA TGCACACCAT GTGCGCCAAT GAATATCTTA ACAATTCAAC 300
TAGTTGGCAT AANAACCGAA CGAAAATCCC AATAGTCTGA AGAGCTCTTT TG 352

*Fig. 17*NUCLEOTIDE SEQUENCE OF THE REPRESENTATIVE
BREAST-TUMOR SPECIFIC cDNA B3CA2

CTGCATGTCC ACGGCCTGGA TTTACGGGTG GTCGGCGTTC ACCCCTGGCA GCTGGCGGCTC 60
TTCCCGACCA GGCCAGCAG GATGTGTGGG GCAAGGATAA CGGCGTGCGC ATCGCCTCGA 120
CCTATATGCC TACTGGCAAG GCCGAGCCCG TGGAAAGGCGG ATTCAGGTTC ANCGGTCGCT 180
GGAGCTTTTC CACCGGCTCC ATGCATTGTG ACTGGCTGTT TCTAGGCGGT CTGTTGCCCA 240
AGCGTGATGG TACGTCTGGC CTGGAGCATG TGACTTTCTG 280

Fig. 18

14/19

NUCLEOTIDE SEQUENCE OF THE REPRESENTATIVE
BREAST-TUMOR SPECIFIC cDNA B3CA3

AG GGAGCAAGGA GAAGGCATGG AGAGGCTCAN GCTGGTCCTG GCCTACGACT 60
CT GTCGCCGGGG ATGGTGGAGA ACTGAAGCGG GACCTCCTCG AGGTCTCTCG 120
TC NCCGTCCAGG AGGAGGGTCT TTCCGTGGTC TNGGAGGAGC GGGGGGAGAA 180
TC ATGGTCNACA TCCC 204

*Fig. 19*NUCLEOTIDE SEQUENCE OF THE REPRESENTATIVE
BREAST-TUMOR SPECIFIC cDNA B4CA1

TC AGGAGCGGGT AGAGTGGCAC CATTGAGGGG ATATTCAAAA ATATTATTTT 60
TG ATAGTTGCTG AGTTTTCTT TGACCCATGA GTTATATTGG AGTTTATTTT 120
CC AATCGCATGG ACATGTTAGA CTTATTTTCT GTTAATGATT NCTATTTTTA 180
GA TTTGAGAAAT TGGTTNTTAT TATATCAATT TTTGGTATTT GTTGAGTTTG 240
GC TTAGTATGTG ACCA 264

Fig. 20

15/19




Fig. 21A

16/19






Fig. 21B

17/19

*Fig. 22*

18/19

*Fig. 23*

19/19




Fig. 24

SEQUENCE LISTING

<110> Corixa Corporation
 Frudakis, Tony N.
 Reed, Steven G.
 Smith, John M.
 Misher, Linda E.
 Dillon, Davin C.
 Retter, Marc W.
 Wang, Aijun
 Skeiky, Yasir A.W.
 Harlocker, Susan L.
 Day, Craig H.

<120> COMPOSITIONS AND METHODS FOR THE
 THERAPY AND DIAGNOSIS OF BREAST CANCER

<130> 210121.41930PC

<140> PCT

<141> 2001-05-22

<160> 334

<170> FastSEQ for Windows Version 3.0

<210> 1

<211> 363

<212> DNA

<213> Homo sapien

<400> 1

ttagagaccc aattgggacc taattgggac ccaaatttct caagtggagg gagaactttt	60
gacgatttcc accggtatct cctcgtgggt attcagggag ctgcccagaa acctataaac	120
ttgtctaagg cgattgaagt cgtccagggg catgatgagt caccaggagt gtttttagag	180
cacctccagg aggcttatcg gatttacacc ccttttgacc tggcagcccc cgaaaatagc	240
catgctctta atttggcatt tgtggctcag gcagccccag atagtaaaag gaaactccaa	300
aaactagagg gattttgctg gaatgaatac cagtcagctt ttagagatag cctaaaaggt	360
ttt	363

<210> 2

<211> 121

<212> PRT

<213> Homo sapien

<400> 2

Leu Glu Thr Gln Leu Gly Pro Asn Trp Asp Pro Asn Phe Ser Ser Gly	
1 5 10 15	
Gly Arg Thr Phe Asp Asp Phe His Arg Tyr Leu Leu Val Gly Ile Gln	
20 25 30	
Gly Ala Ala Gln Lys Pro Ile Asn Leu Ser Lys Ala Ile Glu Val Val	
35 40 45	
Gln Gly His Asp Glu Ser Pro Gly Val Phe Leu Glu His Leu Gln Glu	
50 55 60	

Ala Tyr Arg Ile Tyr Thr Pro Phe Asp Leu Ala Ala Pro Glu Asn Ser
 65 70 75 80
 His Ala Leu Asn Leu Ala Phe Val Ala Gln Ala Ala Pro Asp Ser Lys
 85 90 95
 Arg Lys Leu Gln Lys Leu Glu Gly Phe Cys Trp Asn Glu Tyr Gln Ser
 100 105 110
 Ala Phe Arg Asp Ser Leu Lys Gly Phe
 115 120

<210> 3
 <211> 1080
 <212> DNA
 <213> Homo sapien
 <220>
 <221> misc_feature
 <222> (1)...(1080)
 <223> n = A,T,C or G

<400> 3
 tcttagaatc ttcatacccc gaactcttgg gaaaacttta atcagtcacc tacagtctac 60
 caccatttta ggaggagcaa agctacctca gtcctccgg agccgtttta agatccccc 120
 tcttcaaagc ctaacagatc aagcagctct ccggtgcaca acctgcgcc aggtaaatgc 180
 caaaaaaggt cctaaaccca gccaggcca ccgtctccaa gaaaactcac caggagaaaa 240
 gtgggaaatt gactttacag aagtaaaacc acaccgggct gggtagaaat accttctagt 300
 actggttagac accttctctg gatggactga agcatttgct accaaaaacg aaactgtcaa 360
 tatggtagtt aagtttttac tcaatgaaat catccctcga cgtgggctgc ctgttgccat 420
 aggtctgat aatggaacgg ccttcgcctt gtctatagtt taatcagtca gtaaggcggt 480
 aaacattcaa tggaagctcc attgtgccta tcgaccacga gctctgggca agtagaacgc 540
 atgaactgca ccctaaaaaa acactcttac aaaattaatc ttaaaaaccg gtgttaattg 600
 tgttagtctc cttcccttag cctacttag agttaagggt cacccttac tgggtgggt 660
 tctttacctt ttgaaatcat ntttnggaag gggctgccta tcttttctta actaaaaaan 720
 gccatttgg caaaaatttc ncaactaatt tntacgtnc taagtctccc caacaggtan 780
 aaaaatctnc tgcccttttc aaggaacct cccatccatt cctnaacaaa aggcctgccc 840
 ttcttcccc agttaactnt tttttnttaa aattcccaaa aaangaaccn cctgctggaa 900
 aaacncccc ctccaanccc cggccnaagn ggaagggtcc cttgaatccc nccccncna 960
 anggcccgga accnttaaan tngttccngg gggtnnggcc taaaagnccn atttggtaaa 1020
 cctanaaatt ttttcttttn taaaaaccac nntttntttt ttcttaaaaca aaacctntt 1080

<210> 4
 <211> 1087
 <212> DNA
 <213> Homo sapien
 <220>
 <221> misc_feature
 <222> (1)...(1087)
 <223> n = A,T,C or G

<400> 4
 tctagagctg cgcctggatc ccgccacagt gaggagacct gaagaccaga gaaaacacag 60
 caagtaggcc ctttaaaacta ctcacctgtg ttgtcttcta atttattctg ttttattttg 120
 tttccatcat ttttaaggggt taaaatcadc ttgttcagac ctcagcatat aaaatgacct 180
 atctgtagac ctcaggctcc aaccataccc caagagttgt ctgggtttgt ttaattact 240
 gccaggtttc agctgcagat atccctggaa ggaatattcc agattccctg agtagtttc 300
 aggttaaaat cctataggct tcttctgttt tgaggaagag ttcctgtcag agaaaaacat 360
 gattttggt ttttaacttt aatgcttggt aaacgctata aaaaaaattt tctaccctta 420
 gctttaaagt actgttagtg agaaattaaa attccttcag gaggattaaa ctgccatttc 480

3

```

agttacccta attccaaatg ttttggtggt tagaatcttc tttaatgttc ttgaagaagt      540
gttttatatt ttcccatcna gataaattct ctcnncctt nntttntnt ctnntttttt      600
aaaaaggant cttgctccgt .tgtccanget gggaattttt ttttggccaa tctccgctnc      660
cttgcaanaa tncgtcntcc caaaattacc ncctttttcc cacctccacc ccnnggaatt      720
acctggaatt anaggcccc ncccccccc cggctaattt gtttttgttt ttagtaaaaa      780
acgggtttcc tgttttagtt aggatggccc anntctgacc cctnatcnt cccctcngc      840
cctcnaatnt tnggnntang gcttaccccc ccngnngtt tttcctccat tnaaattttc      900
tntggantct tgaatnncgg gttttccctt ttaaaccnat tttttttttt nnnccccan      960
ttttncctcc cccntntnta angggggttt cccaanccgg gtccncccc angtccccaa     1020
tttttctccc cccccctctt ttttctttnc cccaaaantc ctatcttttc ctnnaaatat     1080
cnantnt

```

```

<210> 5
<211> 1010
<212> DNA
<213> Homo sapien

<220>
<221> misc_feature
<222> (1)...(1010)
<223> n = A,T,C or G

```

```

<400> 5
tctagaccaa gaaatgggag gatttttagag tgactgatga tttctctatc atctgcagtt      60
agtaaacatt ctccacagtt tatgcaaaaa gtaacaaaac cactgcagat gacaaacact     120
aggtaacaca catactatct cccaaatacc taccacaaag ctcaacaatt ttaaactggt     180
aggatcactg gctctaatac ccatgacatg aggtcaccac caaaccatca agcgctaaac     240
agacagaatg tttccactcc tgatccactg tgtgggaaga agcaccgaac ttaccactg      300
gggggcctgc ntcanaanaa aagcccatgc ccccggtnt ncctttnaac cggaacgaat     360
naaccaccca tccccacanc tcctctgttc ntgggccctg catcttgttg cctcntntnc     420
tttnggggan acntggggaa ggtaccccat ttcnttgacc ccncnanaaa acccngtgg      480
ccctttgccc tgattcncnt gggccttttc tcttttccct tttgggttgt ttaaattccc     540
aatgtcccn gaacctctc cntnctgccc aaaacctacc taaattnctc nctangnntt     600
ttcttggtgt tncctttcaa aggtnacctt ncctgttcan ncccnacnaa aattntttcc     660
ntatnntggn cccnnaaaaa nnnatcnnc cnaattgccc gaattggttn ggtttttcct     720
nctgggggaa accctttaaa tttccccctt ggccggcccc ccttttttcc ccccttnga      780
aggcaggngg ttcttccga acttccaatt ncaacagcgn tgccattgn tgaaaccctt     840
ttcctaaaat taaaaaatan cgggttnng nngcctctt tccctcng gnggngngng     900
aaantcctta cccnnaaaaa ggttgcttag ccccngtcc ccaactcccc nggaaaaatn     960
aaccttttcn aaaaaggaa tataantttt ccactccttn gttctcttcc                    1010

```

```

<210> 6
<211> 950
<212> DNA
<213> Homo sapien

<220>
<221> misc_feature
<222> (1)...(950)
<223> n = A,T,C or G

```

```

<400> 6
tctagagctc gcggccgca gctctaatac gactcactat agggcgtcga ctgatctca      60
gctcactgca atctctgccc ccggggtcat gcgattctcc tgcctcagcc ttccaagtag     120
ctgggattac aggcgtgcaa caccacaccc ggctaatttt gtatttttaa tagagatggg     180
gttttccctt gttggccann atggtctcna accctgacc tcnngtgatc ccccncccn      240
ngantcenna ctgctgggga tnnccgnnnn nnnccctccn ncncnnnnnn ncncnntccn      300
tntccttnc tcnnnnnnnn cnntcnntcc ncttctcnc cnnntntnt cncnncnncn      360

```

cnmncncnt	nccncnnnt	tcnctncnn	tntccnnnn	nttcnnnnn	cnmncntnn	420
ccntacntc	ntnnncnnnt	ccntctntnn	cctcnnnnnt	cnctncncnt	tntctectcn	480
ntnnnnnnct	ccnnnnntct	cntcncnnn	tnctcnnntn	ncncncccc	ncctcncnc	540
ctnnntnnn	cnncnnntcc	ntnccnttcn	nttcnnntnn	cnncntcncn	nnctntnttc	600
ccncnnttc	cttnccntn	nnntntcnnn	cnctcnnntc	ntttntctct	nnntcccnnc	660
tcnnttcnc	cnnttcnc	ccccnctnt	ctctcncnn	nnnnntntn	nnncntcnc	720
tntcncnttc	ntcnnntcnc	tnctntcnn	nnnnntcnc	tnccntntnt	ctnnntcnc	780
tcnctntcn	ccntccttn	ctntctctn	tntccttccc	ctcncctnct	cnctcncnc	840
ccntntntn	tnncncnnnt	ntnnncnnc	cntcctttn	tctctntnn	nnntnnctc	900
nnccntncc	ctnnntcnc	ntnntaccn	tnctntcnc	tctccttcc		950

<210> 7
 <211> 1086
 <212> DNA
 <213> Homo sapien
 <220>
 <221> misc_feature
 <222> (1)...(1086)
 <223> n = A,T,C or G

<400> 7		
tctagagctc	gcggccgcga gctcaattaa ccctactaa agggagtcga ctcgatcaga 60	
ctgttactgt	gtctatgtag aaagaagtag acataagaga ttccattttg ttctgtacta 120	
agaaaaattc	ttctgccttg agatgctgtt aatctgtaac cctagcccca accctgtgct 180	
cacagagaca	tgtgctgtgt tgactcaagg ttcaatggat ttagggctat gctttgttaa 240	
aaaagtgtt	gaagataata tgcttgtaa agtcatcac cattctctaa tctcaagtac 300	
ccagggacac	aatacactgc ggaaggccgc agggacctct gtctaggaaa gccaggtatt 360	
gtccaagatt	tctccccatg tgatagcctg agatatggcc tcatgggaag ggtaagacct 420	
gactgtcccc	cagcccagaca tccccagcc cgacatcccc cagcccagaca cccgaaaagg 480	
gtctgtgctg	aggaagatta ntaaaagagg aaggctcttt gcattgaagt aagaagaagg 540	
ctctgtctcc	tgctcgtccc tgggcaataa aatgtcttgg tgtaaaccac gaatgtatgt 600	
tctacttact	gagaatagga gaaaacatcc ttagggctgg aggtgagaca ccctggcggc 660	
atactgctct	ttaatgcacg agatgtttgt ntaattgcca tccagggcca ncccctttcc 720	
ttaacttttt	atganacaaa aactttgttc ncttttctg cgaacctctc cccctattan 780	
cctattggcc	tgcccatccc ctccccaaan ggtgaaaana tgttcntaaa tncgagggaa 840	
tccaaaacnt	tttcccgttg gtcccccttc caaccccgtc cctgggccnn tttctcccc 900	
aacntgtccc	ggntccttcn ttccncccc ctcccnngan aaaaaacccc gntnganggn 960	
gccccctcaa	attataacct ttccnaaaca aannngttcn aagggtgttt gnttcgggtg 1020	
cggctggcct	tgaggtcccc cctncacccc aatttggaan ccngtttttt ttattgcccc 1080	
ntcccc		1086

<210> 8
 <211> 1177
 <212> DNA
 <213> Homo sapien
 <220>
 <221> misc_feature
 <222> (1)...(1177)
 <223> n = A,T,C or G

<400> 8	
ncnntttaga	gtttgacaan ntaacaagc ngctcaggca gctgaaaaaa gccactgata 60
aagcatcctg	gagtatcaga gtttactgtt agatcagcct catttgactt cccctccac 120
atggtgttta	aatccagcta cactacttcc tgactcaaac tccactatc ctgttcatga 180
ctgtcaggaa	ctgtttgaaa ctactgaaac tggccgacct gatcttcaaa atgtgccct 240
aggaaagggtg	gatgccaccg tgttcacaga cagtaccnc ttcctcgaga agggactacg 300

```

aggggcccgt  gcancgttta  ccaaggagac  tnatgtgttg  tgggctcagg  ctttaccanc  360
aaacacctca  ncnncnaagg  ctgaattgat  cgcctcact  caggctctcg  gatggggtaa  420
gggatattaa  cgttaacact  gacagcaggt  acgcctttgc  tactgtgcat  gtacgtggag  480
ccatctacca  ggagcgtggg  ctactcactc  ggcaggtggc  tgnatccac  tgtaaangga  540
catcaaaagg  aaaacnnggc  tgttgcccg  ggtaaccana  aancgtatcn  ncagctcnaa  600
gatgctgtgt  tgactttcac  tcncncctct  taaacttgct  gccacantc  tcctttccca  660
accagatctg  cctgacaatc  ccctactca  aaaaaaaan  aanactggcc  ccgaaccna  720
accaataaaa  acggggangg  tnggtnganc  nncctgaccc  aaaaataatg  gatcccccg  780
gctgcaggaa  ttcaattcan  ctttatcnat  accccaacn  nggngggggg  ggccngtncc  840
cattncctct  ntattnatc  tttnncccc  cccccggcnt  cctttttnaa  ctggtgaaag  900
ggaaaacctg  ncttaccan  ttatcnctg  gacntcccc  ttccncggt  gnttanaaaa  960
aaaagccnc  antccntcc  naaatttgca  cngaaaggna  aggaatttaa  cctttatttt  1020
ttntccttt  antttgtnn  cccctttta  ccagggcgaa  cngccatcnt  ttaanaaaaa  1080
aaanagaang  tttattttc  cttngaacca  tcccaatana  aancaccgc  nggggaacgg  1140
gnggnaggc  cnctcacc  cttntgtng  gngggnc  1177

```

```

<210> 9
<211> 1146
<212> DNA
<213> Homo sapien

<220>
<221> misc_feature
<222> (1)... (1146)
<223> n = A,T,C or G

```

```

<400> 9
nccnnttnnt  gatgttgtct  ttttggcctc  tctttggata  ctttccctct  cttcagaggt  60
gaaaagggtc  aaaaggagct  gttgacagtc  atcccagggt  ggccaatgtg  tccagagtac  120
agactccatc  agtgagggtca  aagcctgggg  cttttcagag  aaggaggat  tatgggtttt  180
ccaattatac  aagtcagaag  tagaaagaag  ggacataaac  caggaagggg  gtggagcact  240
catcaccag  agggacttgt  gcctctctca  gtggtagtag  aggggctact  tcctccacc  300
acggttgcaa  ccaagaggca  atgggtgatg  agcctacagg  ggacatancc  gaggagacat  360
gggatgaccc  taaggagta  ggctggtttt  aaggcgggtg  gactgggtga  gggaaactct  420
cctcttcttc  agagagaagc  agtacagggc  gagctgaacc  ggctgaaggt  cgaggcgaaa  480
acacggtctg  gctcaggaag  accttggaag  taaaattatg  aatggtgcat  gaatggagcc  540
atggaagggg  tgctcctgac  caaactcagc  cattgatcaa  tgtagggaa  actgatcagg  600
gaagccggga  atttcattaa  caaccgcga  cacagctga  acattgtgag  gttcagtac  660
ccttcaaggg  gccactccac  tccaactttg  gccattctac  tttgcnaaat  ttccaaaact  720
tcctttttta  aggccgaatc  cntantccct  naaaaacnaa  aaaaaatctg  cncctattct  780
ggaaaaggcc  cancccttac  caggctggaa  gaaattttnc  cttttttttt  tttttgaagg  840
cntttnttaa  attgaacctn  aattcnccc  cccaaaaaaa  aaccncng  gggggcgat  900
ttccaaaaac  naattccctt  accaaaaaac  aaaaaccnc  cctnttccc  ttccnccctn  960
ttcttttaat  tagggagaga  tnaagcccc  caatttcng  gnetngatnn  gtttcccccc  1020
ccccatttt  ccnaaacttt  ttccancna  ggaancnc  ctttttttng  gtcngattna  1080
ncaaccttcc  aaacctttt  tccnaaaaa  nttgtntng  ngggaaaaan  acctnntttt  1140
atagan  1146

```

```

<210> 10
<211> 545
<212> DNA
<213> Homo sapien

```

```

<400> 10
cttcattggg  tacgggcccc  ctogaggtcg  acggtatcga  taagcttgat  atcgaattcc  60
tgcagcccg  gggatccact  agttctagag  tcaggaagaa  ccaccaacct  tcctgatttt  120
tattggctct  gagttctgag  gccagttttc  ttcttctgtt  gagtatgcgg  gattgtcagg  180
cagatctggc  tgtggaaagg  agactgtggg  cagcaagttt  agaggcgtga  ctgaaagtca  240

```



```

cactgcatct tgagctgctg aatcagcttt ctggttacca cgggcaacag ccgtgttttc 300
cttttgatgt cctttacagt ggattacagc cacctgctga ggtgagtagc ccacgctcct 360
ggtagatggc tccacgtaca tgcacagtag caaaggcgta cctgctgtca gtgttaacgt 420
taatatcctt accccatcgg agagcctgag tgagggcgat caattcagcc cttttgtgct 480
gaggtgtttg ctggttaagc cctgaacca caacacatct gtctccatgg taacagctgc 540
accgg 545

```

```

<210> 11
<211> 196
<212> DNA
<213> Homo sapien

```

```

<400> 11
tctcctaggc tgggcacagt ggctcatacc tgtaatcctg accgtttcag aggctcaggt 60
ggggggatcg cttgagccca agatttcaag actagtctgg gtaacatagt gagaccctat 120
ctctacgaaa aaataaaaaa atgagcctgg tgtagtggca cacaccagct gaggagggag 180
aatcgagcct aggaga 196

```

```

<210> 12
<211> 388
<212> DNA
<213> Homo sapien

```

```

<220>
<221> misc_feature
<222> (1)...(388)
<223> n = A,T,C or G

```

```

<400> 12
tctcctaggc ttgggggctc tgactagaaa ttcaaggaac ctgggattca agtccaactg 60
tgacaccaac ttactctgtg gnetccaata aactgcttct ttcctattcc ctctctatta 120
aataaaataa ggaaaacgat gtctgtgtat agccaagtca gntatcctaa aaggagatac 180
taagtacat taaatatcag aatgtaaaac ctgggaacca ggttcccagc ctgggattaa 240
actgacagca agaagactga acagtactac tgtgaaaagc ccgaagnngc aatatgttca 300
ctctaccgtt gaaggatggc tgggagaatg aatgctctgt ccccagttcc caagctcact 360
tactatacct cctttatagc ctaggaga 388

```

```

<210> 13
<211> 337
<212> DNA
<213> Homo sapien

```

```

<400> 13
tagtagttgc ctataatcat gtttctcatt attttcacat tttattaacc aatttctggt 60
taccctgaaa aatatgaggg aaatatatga aacagggagg caatgttcag ataattgatc 120
acaagatatg atttctacat cagatgctct ttcctttcct gtttatttcc tttttatttc 180
ggttgtgggg tcgaatgtaa tagctttgtt tcaagagaga gttttggcag tttctgtagc 240
ttctgacact gctcatgtct ccaggcatct atttgcactt taggaggtgt cgtgggagac 300
tgagaggtct attttttcca tatttgggca actacta 337

```

```

<210> 14
<211> 571
<212> DNA
<213> Homo sapien

```

```

<220>
<221> misc_feature
<222> (1)...(571)

```

<223> n = A,T,C or G

<400> 14

tagtagttgc	catacagtc	ctttccattt	atttaacccc	cacctgaacg	gcataaactg	60
agtgttcagc	tggtgttttt	tactgtaaac	aataaggaga	ctttgctcct	catttaaacc	120
aaaatcatat	ttcatatttt	acgctcgagg	gtttttaccg	gttccttttt	acactcctta	180
aaacagtttt	taagtcgttt	ggaacaagat	attttttctt	tcctggcagc	ttttaacatt	240
atagcaaat	tgtgtctggg	ggactgctgg	tcactgtttc	tcacagttgc	aaatcaaggc	300
atttgcaacc	aagaaaaaaa	aatttttttg	ttttatttga	aactggaccg	gataaacggt	360
gtttggagcg	gctgctgtat	atagttttaa	atggtttatt	gcacctcctt	aagttgcact	420
tatgtggggg	ggggnntttg	natagaaagt	ntttantcac	anagtcacag	ggacttttnt	480
cttttggnna	ctgagctaaa	aagggtgnt	tttcgggtgg	gggcagatga	aggctcacag	540
gaggcccttc	tcttagaggg	gggaactnct	a			571

<210> 15

<211> 548

<212> DNA

<213> Homo sapien

<220>

<221> misc_feature

<222> (1)...(548)

<223> n = A,T,C or G

<400> 15

tatatattta	ataacttaaa	tatatatttga	tcacccactg	gggtgataag	acaatagata	60
taaaagtatt	tccaaaaagc	ataaaaccaa	agtatcatac	caaaccaaat	tcatactgct	120
tccccacccc	gcactgaaac	ttcaccttct	aactgtctac	ctaaccaaat	tctacccttc	180
aagtcttttg	tgcggtgctca	ctactctttt	tttttttttt	ttnttttttg	agatggagtc	240
tggctgtgca	gccaggggt	ggagtacaat	ggcacaaact	cagctcactg	naacctccgc	300
ctcccagggt	catgagattc	tcctgnttca	gccttcccag	tagctgggac	tacaggtgtg	360
catcaccatg	cctggntaat	cttttttngt	tttngggtag	agatgggggt	tttacctgtt	420
ggccaggntg	gtntcgaaact	cctgacctca	agtgatccac	ccacctcagg	ctcccaaagt	480
gctaggatta	cagacatgag	ccactgngcc	cagncttggg	gcatgctcac	ttctctaggc	540
aactacta						548

<210> 16

<211> 638

<212> DNA

<213> Homo sapien

<220>

<221> misc_feature

<222> (1)...(638)

<223> n = A,T,C or G

<400> 16

ttccgttatg	cacatgcaga	atattctatc	ggtacttcag	ctattactca	ttttgatggc	60
gcaatccgag	cctatcctca	agatgagtat	ttagaaagaa	ttgatttagc	gatagaccaa	120
gctggtaagc	actctgacta	cacgaaattg	ttcagatgtg	atggatttat	gacagttgat	180
ctttggaaga	gattattaag	tgattatttt	aaaggggaatc	cattaattcc	agaatatctt	240
ggttttagctc	aagatgatat	agaaatagaa	cagaaagaga	ctacaaatga	agatgtatca	300
ccaactgata	ttgaagagcc	tatagtagaa	aatgaattag	ctgcatttat	tagccttaca	360
catagcgatt	ttcctgatga	atcttatatt	cagccatcga	catagcatta	cctgatgggc	420
aaccttacga	ataatagaaa	ctgggtgcgg	ggctattgat	gaattcatcc	ncagtaaatt	480
tggatatnac	aaaatataac	tcgattgcat	ttggatgatg	gaatactaaa	tctggcaaaa	540
gtaacttttg	agctactagt	aacctctctt	tttgagatgc	aaaattttct	tttagggttt	600
cttattctct	actttacgga	tattggagca	taacggga			638

<210> 17
 <211> 286
 <212> DNA
 <213> Homo sapien

<400> 17
 actgatggat gtcgccggag gcgagggggc ttatctgatg ctcggctgcc tgttcgtgat 60
 gtgcgcggcg attgggctgt ttatctcaaa caccgccacg gcggtgctga tggcgccctat 120
 tgccttagcg gcggcgaagt caatgggcgt ctcaccctat ccttttgcca tgggtggtggc 180
 gatggcggct tcggcggcgt ttatgacccc ggtctcctcg ccggttaaca ccctggtgct 240
 tggccctggc aagtactcat ttagcgattt tgtcaaaata ggcgtg 286

<210> 18
 <211> 262
 <212> DNA
 <213> Homo sapien

<220>
 <221> misc_feature
 <222> (1)...(262)
 <223> n = A,T,C or G

<400> 18
 tcgggtcatag cagccccttc ttctcaattt catctgtcac taccctgggt tagtatctca 60
 tagccttaca tttttatagc ctctccctg gtctgtcttt tgattttcct gcctgtaate 120
 catatcacac ataactgcaa gtaaacattt ctaaagtgtg gttatgctca tgtcactcct 180
 gtgncaagaa atagtttcca ttaccgtctt aataaaattc ggatttggtt tttncatatt 240
 tcactcttca cctatgaccg aa 262

<210> 19
 <211> 261
 <212> DNA
 <213> Homo sapien

<400> 19
 tcgggtcatag caaagccagt ggtttgagct ctctactgtg taaactccta aaccaaggcc 60
 atttatgata aatgggtggc ggatttttat tataaacatg taccatgca aatttcctat 120
 aactctgaga tatattcttc tacatttaaa caataaaaat aatctatttt taaaagccta 180
 atttcgtag ttaggtaaga gtgtttaatg agaggggtata aggtataaat caccagtcaa 240
 cgtttctctg cctatgaccg a 261

<210> 20
 <211> 294
 <212> DNA
 <213> Homo sapien

<220>
 <221> misc_feature
 <222> (1)...(294)
 <223> n = A,T,C or G

<400> 20
 tacaacgagg cgacgtcggg aaaatcggac atgaagccac cgctgggtctt ttcgtccgag 60
 cgatagggcg cggccagcca gcggaacggt tgcccggatg gcgaagcgag ccggagtctt 120
 tcggactgag tatgaatctt gttgtgaaaa tactcgccgc cttcgttcga cgacgtcgcg 180
 tcgaaatctt cgantcctt acgatcgaag tcttcgtggg cgacgatcgc ggtcagttcc 240
 gccccaccga aatcatggtt gagccggatg ctgnccccga agncctcgtt tgtn 294

<210> 21
 <211> 208
 <212> DNA
 <213> Homo sapien

<220>
 <221> misc_feature
 <222> (1)...(208)
 <223> n = A,T,C or G

<400> 21
 ttggtaaagg gcatggacgc agacgcctga cgtttggtcg aaaatctttc attgattcgt 60
 atcaatgaat aggaaaattc ccaaagaggg aatgtcctgt tgctcgccag tttttntgtt 120
 gttctcatgg anaaggcaan gagctcttca gactattggn attntcgttc ggtcttctgc 180
 caactagtcg ncttgcnang atcttcat 208

<210> 22
 <211> 287
 <212> DNA
 <213> Homo sapien

<220>
 <221> misc_feature
 <222> (1)...(287)
 <223> n = A,T,C or G

<400> 22
 nccnttgagc tgagtgattg agatntgtaa tgggtgtaag ggtgattcag gcggattagg 60
 gtggcgggtc acccggcagt gggctctccc acaggccagc aggatttggg gcaggtacgg 120
 ngtgcgcac gctcgactat atgctatggc aggcgagccg tgggaaggngg atcaggtcac 180
 ggcgctggag ctttccacgg tccatgnatt gngatggctg ttctaggcgg ctggttgccaa 240
 gcgtgatggt acgctggctg gagcattgat ttctggtgcc aaggtgg 287

<210> 23
 <211> 204
 <212> DNA
 <213> Homo sapien

<220>
 <221> misc_feature
 <222> (1)...(204)
 <223> n = A,T,C or G

<400> 23
 ttgggtaaag ggagcaagga gaaggcatgg agaggctcan gctggctcctg gcctacgact 60
 gggccaagct gtcgccgggg atggtggaga actgaagcgg gacctcctcg aggtcctccg 120
 ncgttacttc nccgtccagg aggagggctc ttccgtggtc tnggaggagc ggggggagaa 180
 gatnctctc atggtcnaca tccc 204

<210> 24
 <211> 264
 <212> DNA
 <213> Homo sapien

<220>
 <221> misc_feature
 <222> (1)...(264)

<223> n = A,T,C or G

<400> 24

tggaattggtc	aggagcgggt	agagtggcac	cattgagggg	atattcaaaa	atattatattt	60
gtcctaaatg	atagttgctg	agtttttctt	tgacccatga	gttatattgg	agtttatttt	120
ttaaatttcc	aatcgcatgg	acatgttaga	cttatittct	gttaatgatt	nctatittta	180
ttaaattgga	tttgagaaat	tggttnttat	tatatcaatt	tttggtattt	gttgagtttg	240
acattatagc	ttagtatgtg	acca				264

<210> 25

<211> 376

<212> DNA

<213> Homo sapien

<220>

<221> misc_feature

<222> (1)...(376)

<223> n = A,T,C or G

<400> 25

ttacaacgag	gggaaactcc	gtctctacaa	aaattaaaaa	attagccagg	tgtggtggtg	60
tgcacccgca	atcccagcta	cttgggaggt	tgagacacaa	gantcaccta	natgtgggag	120
gtcaaggttg	catgagtcac	gattgtgcca	ctgcactcca	gcctgggtga	cagaccgaga	180
ccctgcctca	anaganaang	aataggaagt	tcagaaatcn	tggntgtggn	gccagcaat	240
ctgcatctat	ncaaccctg	caggcaangc	tgatgcagcc	tangttcaag	agctgctgtt	300
tctggaggca	gcagttnggg	cttccatcca	gtatcacggc	cacactcgca	cnagccatct	360
gtcctccgtn	tgtnac					376

<210> 26

<211> 372

<212> DNA

<213> Homo sapien

<220>

<221> misc_feature

<222> (1)...(372)

<223> n = A,T,C or G

<400> 26

ttacaacgag	gggaaactcc	gtctctacaa	aaattaaaaa	attagccagg	tgtggtggtg	60
tgcacctgta	atcccagcta	cttgggcggc	tgagacacaa	gaaccaccta	aatgtgggag	120
ggtcaaggtt	gcatgagtca	tgatcgcgcc	actgcactcc	agcctgggtg	acagactgag	180
accctgcctc	aaaagaaaaa	gaataggaa	ttcagaaacc	ctgggtgtgg	ngcccagcaa	240
tctgcattta	aacaatccct	gcaggcaatg	ctgatgcagc	ctaagttcaa	gagctgctgt	300
tctggaggca	gnagtaagg	cttccatcca	gcacacgggn	caacactgca	aaagcacctg	360
tcctcgttgg	ta					372

<210> 27

<211> 477

<212> DNA

<213> Homo sapien

<400> 27

ttctgtccac	atctacaagt	tttattttatt	ttgtgggttt	tcagggtgac	taagtttttc	60
cctacattga	aaagagaagt	tgctaaaagg	tgacacaggaa	atcatttttt	taagtgaata	120
tgataaatatg	ggtccgtgct	taatacaact	gagacatatt	tgttctctgt	tttttttagag	180
tcacctctta	aagtccaatc	ccacaatggt	gaaaaaaaaa	tagaaagtat	ttgttctacc	240
tttaaggaga	ctgcagggat	tctccttgaa	aacggagtat	ggaatcaatc	ttaaataaat	300

atgaaattgg	ttggtcttct	gggataagaa	attcccaact	cagtgtgctg	aaattcacct	360
gacttttttt	gggaaaaaat	agtcgaaaat	gtcaatttgg	tccataaaat	acatgttact	420
attaaaagat	atttaaagac	aaattctttc	agagctctaa	gattgggtgtg	gacagaa	477

<210> 28

<211> 438

<212> DNA

<213> Homo sapien

<220>

<221> misc_feature

<222> (1)...(438)

<223> n = A,T,C or G

<400> 28

tctncaacct	cttgantgtc	aaaaaccttn	taggctatct	ctaaaagctg	actggtattc	60
attccagcaa	aatccctcta	gtttttggag	tttcctttta	ctatctgggg	ctgcctgagc	120
cacaaatgcc	aaattaagag	catggctatt	ttcgggggct	gacaggtcaa	aaggggtgta	180
aatccgataa	gcctcctgga	ggtgctctaa	aaacactcct	ggtgactcat	catgcccctg	240
gacgacttca	atcgncttag	acaagtttat	aggtttcttg	gcagctccct	gaatacccac	300
gaggagatgc	cgggtgaaat	cgtaaaaagt	tctccctoca	cttgagaaat	ttgggtccca	360
attaggctcc	aattgggtct	ctaataccta	ttcctctagc	ttcctcctcc	ggnctattgg	420
ttgatgtgag	gttgaaga					438

<210> 29

<211> 620

<212> DNA

<213> Homo sapien

<220>

<221> misc_feature

<222> (1)...(620)

<223> n = A,T,C or G

<400> 29

aagagggtag	cagccccaag	ccttgacaac	ttccataggg	tgtcaagcct	gtgggtgcac	60
agaagtcaaa	aattgagttt	tgggatcctc	agcctagatt	tcagaggata	taaagaaaca	120
cctaacacct	agatattcag	acaaaagttt	actacaggga	tgaagctttc	acggaaaacc	180
tctactagga	aagtacagaa	gagaaatgtg	ggtttgagac	cccaaacag	aatccctct	240
agaacactgc	ctaataaaac	tgtgagaaga	tggccactgt	catccagaca	ccagaatgat	300
agaccaccca	aaaacttatg	ccatattgcc	tataaaacct	acagacactc	aatgccagcc	360
ccatgaaaaa	aaaactgaga	agaagactgt	nccctacaat	gccaccggag	cagaactgcc	420
ccaggccatg	gaagcacagc	tcttatatca	atgtgacctg	gatgttgaga	catggaatcc	480
nangaaatcn	ttttaanact	tccacggtn	aatgactgcc	ctattanatt	cngaacttan	540
atccnggcct	gtgacctctt	tgctttggcc	attccccctt	tttggaatgg	ctnttttttt	600
cccatgcctg	tncctcttta					620

<210> 30

<211> 100

<212> DNA

<213> Homo sapien

<400> 30

ttacaacgag	ggggtcaatg	tcataaatgt	cacaataaaa	caatctcttc	tttttttttt	60
tttttttttt	tttttttttt	tttttttttt	tttttttttt			100

<210> 31

<211> 762

12

<212> DNA

<213> Homo sapien

<220>

<221> misc_feature

<222> (1)...(762)

<223> n = A,T,C or G

<400> 31

tagtctatgc	gccggacaga	gcagaattaa	attggaagtt	gccctccgga	ctttctaccc	60
acactcttcc	tgaaaagaga	aagaaaagag	gcaggaaaga	ggttaggatt	tcattttcaa	120
gagtcagcta	attaggagag	cagagttag	acagcagtag	gcaccccatg	atacaacca	180
tggacaaagt	ccctgttttag	taactgccag	acatgatcct	gctcagggtt	tgaaatctct	240
ctgccataa	aagatggaga	gcaggagtgc	catccacatc	aacacgtgtc	caagaaagag	300
tctcagggag	acaaggggtat	caaaaaacaa	gattctttaat	gggaaggaaa	tcaaaccaaa	360
aaattagatt	tttctctaca	tatatataat	atacagatat	ttaacacatt	attccagagg	420
tggctccagt	ccttggggct	tgagagatgg	tgaaaacttt	tgttccacat	taacttctgc	480
tctcaaattc	tgaagtatat	cagaatggga	caggcaatgt	tttgctccac	actggggcac	540
agacccaaat	ggttctgtgc	ccgaagaaga	gaagcccga	agacatgaag	gatgcttaag	600
gggggttggg	aaagccaaat	tggtantatc	ttttcctcct	gcctgtgttc	cngaagtctc	660
cnctgaagga	attcttaaaa	ccctttgtga	ggaaatgcc	ccttaccatg	acaantggtc	720
ccattgcttt	tagggngatg	gaaacaccaa	gggttttgat	cc		762

<210> 32

<211> 276

<212> DNA

<213> Homo sapien

<400> 32

tagtctatgc	gtgtattaac	ctcccctccc	tcagtaacaa	ccaaagaggc	aggagctggt	60
attaccaacc	ccatttttaca	gatgcatcaa	taatgacaga	gaagtgaagt	gacttgcgca	120
cacaaccagt	aaattggcag	agtcagattt	gaatccatgg	agtctggtct	gcactttcaa	180
tcaccgaata	ccctttctaa	gaaacgtgtg	ctgaatgagt	gcatggataa	atcagtgtct	240
actcaacatc	tttgcttaga	tatcccgcat	agacta			276

<210> 33

<211> 477

<212> DNA

<213> Homo sapien

<400> 33

tagtagttgc	caaataattt	aaaatttacc	cagaagtgat	tgaaaacttt	ttggaaacaa	60
aaacaaataa	agccaaaagg	taaaataaaa	atatctttgc	actctcgtaa	ttacctatcc	120
ataacttttt	caccgtaagc	tctcctgctt	gttagtgtag	tgtggttata	ttaaactttt	180
tagttattat	tttttattca	cttttccact	agaaagtcac	tattgattta	gcacacatgt	240
tgatctcatt	tcattttttc	tttttatagg	caaaatttga	tgctatgcaa	caaaaatact	300
caagccatt	atcttttttc	ccccgaaat	ctgaaaattg	caggggacag	agggaagtta	360
tccattataa	aaattgtaaa	tatgttcagt	ttatgtttta	aaatgcacaa	aacataagaa	420
aattgtgttt	acttgagctg	ctgattgtaa	gcagttttat	ctcaggggca	actacta	477

<210> 34

<211> 631

<212> DNA

<213> Homo sapien

<400> 34

tagtagttgc	caattcagat	gatcagaaat	gctgctttcc	tcagcattgt	cttggttaaac	60
cgcagcccat	ttggaacttt	ggcagtgaga	agccaaaagg	aagaggtgaa	tgacatatat	120

```

atatatatat attcaatgaa agtaaaatgt atatgctcat atacttttcta gttatcagaa      180
tgagttaagc tttatgccat tgggctgctg catattttaa tcagaagata aaagaaaatc      240
tgggcatttt tagaatgtga tacatgtttt tttaaaactg ttaaataatta tttcgatatt      300
tgtctaagaa ccggaatgtt cttaaaattt actaaaacag tattgtttga ggaagagaaa      360
actgtactgt ttgccattat tacagtogta caagtgcacg tcaagtcacc cactctctca      420
ggcatcagta tccacctcat agctttacac attttgacgg ggaatattgc agcatcctca      480
ggcctgacat ctgggaaagg ctcagatcca cctactgctc cttgctcggt gatttgtttt      540
aaaatattgt gcctgggtgc acttttaagc cacagccctg cctaaaagcc agcagagAAC      600
agaaccgcga ccattctata ggcaactact a                                     631

```

```

<210> 35
<211> 578
<212> DNA
<213> Homo sapien

```

```

<400> 35
tagtagttgc catcccatat tacagaaggc tctgtataca tgacttattt ggaagtgatc      60
tgttttctct ccaaaccat ttatcgtaat ttcaccagtc ttggatcaat cttggtttcc      120
actgatacca tgaaacctac ttggagcaga cattgcacag ttttctgtgg taaaaactaa      180
agggttattt gctaagctgt catcttatgc ttagtatttt ttttttacag tggggaattg      240
ctgagattac attttgttat tcattagata ctttgggata acttgacact gtcttctttt      300
tttcgctttt aattgctatc atcatgcttt tgaaacaaga acacattagt cctcaagtat      360
tacataagct tgcttggtac gcctgggtgt ttaaaggact atctttggcc tcaggttcac      420
aagaatgggc aaagtgtttc cttatgttct gtagttctca ataaaagatt gccaggggcc      480
gggtactgtg gctcgactg taatcccagc actttgggaa gctgaggctg gcggtacatg      540
ttagggcagg tgttcgaaac cagcctgggc aactacta                                     578

```

```

<210> 36
<211> 583
<212> DNA
<213> Homo sapien

```

```

<400> 36
tagtagttgc ctgtaatccc agcaactcag gaggctgggg caggagaatc agttgaacct      60
gggaggcaga agttgtaatt agcaaagatc gcaccattgc acttcagcct gggcaacaag      120
agttagattc catctcaaaa acaaaaaaaa gaaaaagaaa agaaaaggaa aaaacgtata      180
aaccagccca aaacaaaatg atcattcttt taataagcaa gactaattta atgtgtttat      240
ttaatcaaaag cagttgaatc ttctgagtta ttggtgaaaa taccatgta gttaatttag      300
ggttcttact tgggtgaacg tttgatgttc acaggttata aaatggttaa caaggaaaat      360
gatgcataaa gaatcttata aactactaaa aataaataaa atataaatgg ataggtgcta      420
tggatggagt ttttgtgtaa tttaaaatct tgaagtcatt ttggatgctc attggttgct      480
tggtaatttc cattaggaaa aggttatgat atggggaac tgtttctgga aattgcggaa      540
tgtttctcat ctgtaaatg ctagtatctc agggcaacta cta                                     583

```

```

<210> 37
<211> 716
<212> DNA
<213> Homo sapien

```

```

<220>
<221> misc_feature
<222> (1)...(716)
<223> n = A,T,C or G

```

```

<400> 37
gatctactag tcatntggat tctatccatg gcagctaagc ctttctgaat ggattctact      60
gctttcttgt tctttaatcc agacccttat atatgtttat gttcacaggc agggcaatgt      120
ttagtgaaaa caattctaaa ttttttattt tgcattttca tgctaatttc cgtcacactc      180

```


cagcaggctt	cctgggagaa	taaggagaaa	tacagctaaa	gacattgtcc	ctgcttactt	240
acagccta	ggtatgcaaa	accacttcaa	taaagtaaca	ggaaaagtac	taaccaggta	300
gaatggacca	aaactgatat	agaaaaatca	gaggaagaga	ggaacaaata	tttactgagt	360
cctagaatgt	acaaggcttt	ttaattacat	attttatgta	aggcctgcaa	aaaacagggtg	420
agtaatcaac	atttgtccca	ttttacatat	aaggaaactg	aagcttaaat	tgaataat	480
aatgcataga	ttttatagtt	agaccatggt	caggtcoccta	tggtatactt	actagctgta	540
tgaatatgag	aaaaataattt	tggtattttc	ttggcatcag	tattttcatc	tgcaaaataa	600
agctaaagtt	atttagcaaa	cagtcagcat	agtgcctgat	acatagtagg	tgctccaaac	660
atgattacnc	tantattngg	tattanaaaa	atccaatata	ggcntggata	aaaccg	716

<210> 38

<211> 688

<212> DNA

<213> Homo sapien

<220>

<221> misc_feature

<222> (1)...(688)

<223> n = A,T,C or G

<400> 38

ttctgtccac	atatcatccc	actttaattg	ttaatcagca	aaactttcaa	tgaaaaatca	60
tccatttttaa	ccaggatcac	accaggaaac	tgaagggtga	ttttttttta	ccttaaaaaa	120
aaaaaaaaa	accaaacaaa	ccaaaacaga	ttaacagcaa	agagttctaa	aaaatttaca	180
tttctcttac	aactgtcatt	cagagaacaa	tagttcttaa	gtctgttaaa	tcttggcatt	240
aaacagagaaa	cttgatgaan	agttgtactt	ggaatattgt	ggattttttt	ttttgtctaa	300
tctcccccta	ttgttttgcc	aacagtaatt	taagttttgt	tggaacatcc	ccgtagttga	360
agtgtaaaaca	atgtatagga	aggaatatat	gataagatga	tgcatcacat	atgcattaca	420
tgtagggacc	ttcacaaact	catgcactca	gaaaacatgc	ttgaagagga	ggagaggacg	480
gcccagggtc	accatccagg	tgccctgagg	acagagaatg	cagaagtggc	actgttgaaa	540
tttagaagac	catgtgtgaa	tggtttcagg	cctgggatgt	ttgccaccaa	gaagtgcctc	600
cgagaaattt	ctttcccatt	tggaatacac	ggtggcctga	tgggtacggt	gggtgaccca	660
acgaagaaaa	tgaaattctg	ccctttcc				688

<210> 39

<211> 585

<212> DNA

<213> Homo sapien

<220>

<221> misc_feature

<222> (1)...(585)

<223> n = A,T,C or G

<400> 39

tagtagttgc	cgcnnaccta	aaanttggaa	agcatgatgt	ctaggaaaca	tantaaaata	60
gggtatgcct	atgtgctaca	gagagatgtt	agcattttaa	gtgcatantt	ttatgtat	120
tgacaaatgc	atatnccctc	ataatccaca	actgattacg	aagctattac	aattaaaaag	180
tttggccggg	cgtggtgggc	ggtggctgac	gcctgtaatc	ccagcacttt	gggaggccga	240
ggcacgcgga	tcacgaggtc	gggagttcaa	gaccatcctg	gctaacacgg	tgaaagtcca	300
tctctactaa	aaatacga	aaattacccc	ggcgtgggtg	cgggcgcctg	tagtcccagc	360
tactccggag	gctgaggcag	gagaatggcg	tgaacccagg	acacggagct	tgcaagtgtc	420
caacatcacg	tcactgccct	ccagcctggg	ggacaggaac	aagantcccc	tcctcanaaa	480
agaaaaatac	tactnatant	ttcnacttta	ttttaantta	cacagaactn	cctcttggtg	540
cccccttacc	attcatctca	cccacctcct	atagggcacn	nctaa		585

<210> 40

<211> 475

15

<212> DNA

<213> Homo sapien

<400> 40

tctgtccaca	ccaatcttag	aagctctgaa	aagaatttgt	ctttaaatat	cttttaatatag	60
taacatgtat	tttatggacc	aaattgacat	tttcgactgt	tttttccaaa	aaagtcaggt	120
gaatttcagc	acactgagtt	gggaatttct	tatcccagaa	gaccaacca	tttcatatatt	180
atttaagatt	gattccatac	tccgttttca	aggagaatcc	ctgcagtctc	cttaaaggta	240
gaacaaatac	ttcctatttt	tttttcacca	ttgtgggatt	ggactttaag	aggtgactct	300
aaaaaaacag	agaacaaata	tgtctcagtt	gtattaagca	cggacccata	ttatcatatt	360
cacttaaaaa	aatgatttcc	tgtgcacctt	ttggcaactt	ctcttttcaa	tgtagggaaa	420
aacttagtca	ccctgaaaac	ccacaaaata	aataaaactt	gtagatgtgg	acaga	475

<210> 41

<211> 423

<212> DNA

<213> Homo sapien

<400> 41

taagagggta	catcgggtaa	gaacgtaggc	acatctagag	cttagagaag	tctggggtag	60
gaaaaaatc	taagtattta	taagggtata	ggtaacattt	aaaagtaggg	ctagctgaca	120
ttatttagaa	agaacacata	cggagagata	agggcaaagg	actaagacca	gaggaacact	180
aatatttagt	gatcacttcc	attcttggtta	aaaatagtaa	cttttaagtt	agcttcaagg	240
aagatttttg	gccatgatta	gttgtcaaaa	gttagttctc	ttgggtttat	attactaatt	300
ttgttttaag	atccttggtta	gtgctttaat	aaagtcattg	tatatcaaac	gctctaaaac	360
attgtagcat	gttaaagtgc	acaatatact	taccatttgt	tgtatatggc	tgtaccctct	420
cta						423

<210> 42

<211> 527

<212> DNA

<213> Homo sapien

<220>

<221> misc_feature

<222> (1)...(527)

<223> n = A,T,C or G

<400> 42

tctcctaggc	taatgtgtgt	gtttctgtaa	aagtaaaaag	ttaaaaattt	taaaaataga	60
aaaaagctta	tagaataaga	atatgaagaa	agaaaatatt	tttgtacatt	tgcacaatga	120
gtttatgttt	taagctaagt	gttattacaa	aagagccaaa	aagggtttta	aaattaaaac	180
gtttgtaaag	ttacagtacc	cttatgttaa	tttataattg	aagaaagaaa	aacttttttt	240
tataaatgta	gtgtagccta	agcatacagt	atttataaag	tctggcagtg	ttcaataatg	300
tcctaggcct	tcacattcac	tactgactc	acccagagca	acttccagtc	ctgtaagctc	360
cattcgtggt	aagtgccta	tacagggtgca	ccatttattt	tacagtattt	ttactgtacc	420
ttctctatgt	ttccatatgt	ttcgatatac	aaataccact	ggttactatn	gcccnacagg	480
taattccagt	aacacggcct	gtatacgtct	ggtancccta	gngaaga		527

<210> 43

<211> 331

<212> DNA

<213> Homo sapien

<400> 43

tcttcaacct	cgtaggacaa	ctctcatatg	cctgggcact	atttttaggt	tactaccttg	60
gctgcccttc	tttaagaaaa	aaaaaagaag	aaaaaagaac	ttttccacaa	gtttctcttc	120
ctctagttagg	aaaattagag	aatcatgtt	tttaattttg	tgttatttca	gatcacaaat	180

16

tcaaacactt	gtaaacatta	agcttctgtt	caatccccctg	ggaagaggat	tcattctgat	240
atttacggtt	caaaagaagt	tgtaatat	tgcttggaac	acagagaacc	agttattaac	300
ttcctactac	tattatataa	taaataataa	c			331

<210> 44
 <211> 592
 <212> DNA
 <213> Homo sapien

<220>
 <221> misc_feature
 <222> (1)...(592)
 <223> n = A,T,C or G

<400> 44						
ggcttagtag	ttgccaggca	aaatarcgtt	gattctcctc	aggagccacc	cccaacaccc	60
ctgtttgctt	ctagacctat	acctagacta	aagtcccagc	agacccttag	aggtgagggt	120
cagagtgacc	cttgaggaga	tgtgctacac	tagaaaagaa	ctgcttgagt	tttctaattt	180
atataagcag	aaatctggag	aagagtcata	ggaatggata	ttaagggtgt	gagataatgg	240
cggaaggaat	atagagttgg	atcaggctgg	acttattgat	ttgaaccac	taagtagaga	300
ttctgctttt	gatgttcag	ctcagggagt	taaaaaaggt	tttaatggtt	ctaatagttt	360
atttgcttgg	ttagctgaaa	tatggataaa	agatggccca	ctgtgagcaa	gctggaaatg	420
cctgatctct	ctcagtttaa	tgtagaggaa	gggatccaaa	agtttaggga	ganttggatg	480
ctggraktgg	attggtcact	ttgrgacct	cccwtcccag	ctgggagggt	ccagaagata	540
cacccttgac	caacgctttg	cgaaatggat	ttgtgatggc	ggcaactact	aa	592

<210> 45
 <211> 567
 <212> DNA
 <213> Homo sapien

<220>
 <221> misc_feature
 <222> (1)...(567)
 <223> n = A,T,C or G

<400> 45						
ggcttagtag	ttgccattgc	gagtgccttc	tcaacgagcg	ttgaacatgg	cggattgtct	60
agattcaacg	gatttgagtt	ttaccagcaa	agcgaaccaa	gcgcggccca	gagaattatg	120
ggttgggttg	ctttgaaaag	atggaaatcc	tgtaggccta	gtcagaaaag	ccttcttgca	180
gaacagttgg	ttctcgggcg	aacgctcatc	aagatgccca	ttggaaaggc	tagcgtgtat	240
ttgggagagc	ctgatagcgt	gtcttctgat	gatgtttgtg	cttggacagt	gacaaaagat	300
atgcaaagca	agtcggaact	agacgtcaag	cttcgtgagc	aaattattgt	agactcctac	360
ttatactgtg	aggaatgata	gccaagggtg	gggactttta	gactaagggtg	gtttgtactt	420
gcgccgatga	tcccaggcag	aaagamctga	tcgctagttt	tatacgggca	actactaagc	480
cgaattccag	cacactggcg	gccgttacta	attggatccg	anctcggtag	cagcttgatg	540
catascttga	gttwtctata	ntgtcnc				567

<210> 46
 <211> 908
 <212> DNA
 <213> Homo sapien

<220>
 <221> misc_feature
 <222> (1)...(908)
 <223> n = A,T,C or G

```

<400> 46
gagcgaaaga ccgagggcag ngntangng cgangaagcg gagagggcca aaaagcaacc      60
gctttccccc gggggtgccg attcattaag gcaggtggag gacaggtttc ccgatggaag      120
gcggcagggg cgcaagcaat taatgtgagt aggccattca ttagcaccgg ggcttaacat      180
ttaagcttcg ggttggtatg tgggtgggaat tgtgagcgga taacaatttc acacaggaaa      240
cagctatgac catgattacg ccaagctatt taggtgacat tatagaataa ctcaagttat      300
gcatcaagct tggtagcgag ttccgatcca ctagtaacgg ccgccagtgt gtggaattcg      360
gcttagtagt tgccgaccat ggagtgtac ctaggctaga atacctgagy tcctccctag      420
cctcactcac attaaattgt atcttttcta cattagatgt cctcagcgcc ttatttctgc      480
tggacwatcg ataaattaat cctgatagga tgatagcagc agattaatta ctgagagtat      540
gttaattgtg catccctcct atataacgta ttgcatcttt aatggagcaa ttctggagat      600
aatccctgaa ggcaaaggaa tgaatcttga ggtgagaaa gccagaatca gtgtccagct      660
gcagttgtgg gagaagggtga tattatgtat gtctcagaag tgacaccata tgggcaacta      720
ctaagcccgga attccagcac actggcgggc gttactaatg gatccgagct cgtaccaag      780
cttgatgcat agcttgagta tctatagtgt cactaaatag cctggcggtta tcatggtcat      840
agctgtttcc tgtgtgaaat tgttatccgc tccaattcc cccaccata cgagccggaa      900
cataaagt

```

<210> 47

<211> 480

<212> DNA

<213> Homo sapien

<220>

<221> misc_feature

<222> (1)...(480)

<223> n = A,T,C or G

```

<400> 47
tgccaacaag gaaagtttta aatttcccct tgaggattct tggatgatcat caaatcagtc      60
ggtttttaag gttgttttct gtcaaataac tctaacttta agccaaacag tatatggaag      120
cacagataka atattacaca gataaaagag gatttgatct aaagtaraga tagttggggg      180
ctttaatttc tggaacctag gtctcccat cttcttctgt gctgaggaaac ttcttggaag      240
cggggattct aaagttcttt ggaagacagt ttgaaacca ccatgttgtt ctcagtacct      300
ttatttttaa aaagtaggtg aacattttga gagagaaaag ggcttggttg agatgaagtc      360
ccccccccc cttttttttt ttttagctga aatagatacc ctatgttnaa rgaarggatt      420
attatttacc atgcaytar scacatgctc tttgatgggc nyctccstac cctccttaag      480

```

<210> 48

<211> 591

<212> DNA

<213> Homo sapien

```

<400> 48
aagaggttac cgagtggaat ttccgcttca ctagtctggt gtggctagtc ggtttcgtgg      60
tggccaacat tacgaacttc caactcaacc gttcttggtg gttcaagcgg gagtaccggc      120
gaggatggtg gcgtgaattc tggcctttct ttgccgtggg atcggtagcc gccatcatcg      180
gtatgtttat caagatcttc tttactaacc cgacctctcc gatttacctg cccgagccgt      240
ggtttaacga ggggaggggg atccagtcac gcgagtactg gtcccagatc ttccgcatcg      300
tcgtgacaat gcctatcaac ttctgtctca ataagttgtg gaccttccga acggtgaagc      360
actccgaaaa cgtccgggtg ctgctgtgct gtgactccca aaatcttgat aacaacaagg      420
taaccgaatc gcgctaagga accccggcat ctcgggtact ctgcatatgc gtaccctta      480
agccgaattc cagcacactg gcggccgtta ctaattggat ccgaactccg taaccaagcc      540
tgatgcgtaa cttgagttat tctatagtgt ccctaaaata acctggcggt a          591

```

<210> 49

<211> 454

<212> DNA

<213> Homo sapien

<400> 49

aagaggggtac	ctgccttgaa	attttaaagt	ctaaggaaar	tgggagatga	ttaagagttg	60
gtgtggcyta	gtcacaccaa	aatgtattta	ttacatcctg	ctcctttcta	gttgacagga	120
aagaaagctg	ctgtggggaa	aggaggata	aatactgaag	ggatttacta	aacaaatgtc	180
catcacagag	ttttcctttt	tttttttttg	agacagagtc	ttgctctgtc	acccaggctg	240
gaatgaagwg	gtatgatctc	agttgaatgc	aacctctacc	tcctaggttc	aagcgattct	300
catgcctcag	cctcctgagc	agctgggact	ataggcgcat	gtaccatgc	caggctaatt	360
tttatatttt	tattagagac	ggggtgttgc	catgttggcc	aggcaggctc	cgaactcctg	420
ggcctcagat	gatctgcccc	accgtaccct	ctta			454

<210> 50

<211> 463

<212> DNA

<213> Homo sapien

<400> 50

aagaggggtac	caaaaaaaag	aaaaaggaaa	aaaagaaaaa	caacttgat	aaggctttct	60
gctgcataca	gctttttttt	tttaaataaa	tggtgccaac	aaatgttttt	gcattcacac	120
caattgctgg	ttttgaaatc	gtactcttca	aaggtatttg	tgcagatcaa	tccaatagtg	180
atgccccgta	ggttttgtgg	actgcccacg	ttgtctacct	tctcatgtag	gagccattga	240
gagactgttt	ggacatgcct	gtgttcacgt	agccgtgatg	tccggggggc	gtgtacatca	300
tggtaccgtg	gggtggggtc	tgcatctggc	gctgggcata	tggctgggtg	cccatcatgc	360
ccatctgcat	ctgcataggg	tattggggcg	tttgatccat	atagccatga	ttgctgtggt	420
agccactggt	catcattggc	tgggacatgc	tgttaccctc	tta		463

<210> 51

<211> 399

<212> DNA

<213> Homo sapien

<400> 51

cttcaacctc	ccaaagtgtc	gggattacag	gactgagcca	ccacgctcag	cctaagcctc	60
tttttacta	ccctctaagc	gatctaccac	agtgatgagg	ggctaaagag	cagtgaatt	120
tgattacaat	aatggaactt	agatttatta	attaacaatt	tttccttagc	atgttggttc	180
cataattatt	aagagtatgg	acttacttag	aaatgagctt	tcattttaag	aatttcactc	240
ttgaccttct	ctattagtct	gagcagtag	acactatacg	tattttattt	aactaaccta	300
ccttgagcta	ttacttttta	aaaggctata	tacatgaatg	tgtattgtca	actgtaaagc	360
cccacagtat	ttaattatat	catgatgtct	ttgagggtg			399

<210> 52

<211> 392

<212> DNA

<213> Homo sapien

<400> 52

cttcaacctc	aatcaacctt	ggtaattgat	aaaatcatca	cttaactttc	tgatataatg	60
gcaataatta	tctgagaaaa	aaaagtgtg	aaagattaaa	cttgcatttc	tctcagaatc	120
ttgaaggata	tttgaataat	tcaaaagcgg	aatcagtagt	atcagccgaa	gaaactcact	180
tagctagaac	gttggaacca	tggatctaag	tccttgccct	tccactaacc	agctgattgg	240
ttttgtgtaa	acctcctaca	cgcttgggct	tggtgcctc	atttgtcaaa	gtaaaggctg	300
aaataggaag	ataatgaacc	gtgtcttttt	ggtctctttt	ccatccatta	ctctgatttt	360
acaaagaggc	ctgtattccc	ctggtgaggt	tg			392

<210> 53

<211> 179

<212> DNA

<213> Homo sapien

<220>

<221> misc_feature

<222> (1)...(179)

<223> n = A,T,C or G

<400> 53

ttcgggtgat	gcctcctcag	gctacagtga	agactggatt	acagaaaggt	gccagcgaga	60
tttcagattc	ctgtaaacct	ctaaagaaaa	ggagtcgcgc	ctcaactgat	gtagaaatga	120
ctagttcagc	atacngagac	acntctgact	ccgattctag	aggactgagt	gacctgcan	179

<210> 54

<211> 112

<212> DNA

<213> Homo sapien

<220>

<221> misc_feature

<222> (1)...(112)

<223> n = A,T,C or G

<400> 54

ttcgggtgat	gcctcctcag	gctacatcat	natagaagca	aagtagaana	atcnngtttg	60
tgcatTTTTCC	cacanacaaa	attcaaata	ntggaagaaa	ttggganagt	at	112

<210> 55

<211> 225

<212> DNA

<213> Homo sapien

<400> 55

tgagcttccg	cttctgacaa	ctcaatagat	aatcaaagga	caactttaac	agggattcac	60
aaaggagtat	atccaaatgc	caataaacat	ataaaaagga	attcagcttc	atcatcatca	120
gaagwatgca	aattaaaacc	ataatgagaa	accactatgt	cccactagaa	tagataaaat	180
cttaaaagac	tggtaaaacc	aagtgttgg	aaggcaagag	gagca		225

<210> 56

<211> 175

<212> DNA

<213> Homo sapien

<400> 56

gctcctcttg	ccttaccac	acattctcaa	aaacctgtta	gagtcctaag	cattctcctg	60
ttagtattgg	gattttaccc	ctgtcctata	aagatgttat	gtacaaaaa	tgaagtggag	120
ggccataccc	tgaggaggag	gagggatctc	tagtgttgtc	agaagcggaa	gctca	175

<210> 57

<211> 223

<212> DNA

<213> Homo sapien

<400> 57

agccatttac	cacccatgga	tgaatggatt	ttgtaattct	agctgttgta	ttttgtgaat	60
ttgttaattt	tggtgttttt	ctgtgaaaca	catacattgg	atatgggagg	taaaggagtg	120
tcccagttgc	tcctgggtcac	tccctttata	gccattactg	tcttgtttct	tgtaactcag	180
gttaggtttt	ggtctctctt	gctccactgc	aaaaaaaaa	aaa		223

<210> 58
 <211> 211
 <212> DNA
 <213> Homo sapien

<400> 58
 gttcgaaggt gaacgtgtag gtagcggatc tcacaactgg ggaactgtca aagacgaatt 60
 aactgacttg gatcaatcaa atgtgactga ggaaacacct gaaggtgaag aacatcatcc 120
 agtggcagac actgaaaata aggagaatga agttgaagag gtaaaagagg aggggtccaaa 180
 agagatgact ttggatgggt ggtaaattggc t 211

<210> 59
 <211> 208
 <212> DNA
 <213> Homo sapien

<400> 59
 gctcctcttg ccttaccaac ttgacacca tcataacca tgtggccagg ttgacagccc 60
 aggctgcaca tcaggggact gcctcgcaat acttcatgct gttgctgctg actgatggtg 120
 ctgtgacgga tgtggaagcc acacgtgagg ctgtggtgcg tgcctcgaac ctgcccattg 180
 cagtgatcat tatgggtgggt aaatggct 208

<210> 60
 <211> 171
 <212> DNA
 <213> Homo sapien

<400> 60
 agccatttac caccataact aaattctagt tcaaactcca acttcttcca taaaacatct 60
 aaccactgac accagttggc aatagcttct tccttcttta acctcttaga gtatttatgg 120
 tcaatgccac acatttctgc aactgaataa agttggtaag gcaagaggag c 171

<210> 61
 <211> 134
 <212> DNA
 <213> Homo sapien

<220>
 <221> misc_feature
 <222> (1)...(134)
 <223> n = A,T,C or G

<400> 61
 cggtgatgc ctctcaggc ttgtgtgtgt ccaactcnact cactggcctc ttctccagca 60
 actggtgaan atgtcctcan gaaaancncc acacgcnct cagggtgggg tgggaancat 120
 canaatcatc nggc 134

<210> 62
 <211> 145
 <212> DNA
 <213> Homo sapien

<400> 62
 agaggttaca tatgcaacag tatataaagg aagaagtgca ctgagaggaa cttcatcaag 60
 gccatttaac caataagtga tagagtcaag gctcaacca ggtgtgacgg attccaggtc 120
 ccaagctcct tactggtacc ctctt 145

<210> 63

<211> 297
 <212> DNA
 <213> Homo sapien

<400> 63
 tgcactgaga ggaattcaaa gggtttatgc caaagaacaa accagtcctc tgcagcctaa 60
 ctcatttggt tttgggctgc gaagccatgt agagggcgat caggcagtag atggtcctc 120
 ccacagtcag cgccatggtg gtccggtaaa gcatttggtc aggcaggcct cgtttcaggt 180
 agacgggcac acatcagctt tctggaaaaa cttttgtagc tctggagctt tgtttttccc 240
 agcataatca tacactgtgg aatcggaggt cagtttagtt ggtaaggcaa gaggagc 297

<210> 64
 <211> 300
 <212> DNA
 <213> Homo sapien

<400> 64
 gcactgagag gaacttccaa tactatgttg aataggagtg gtgagagagg gcacccctgt 60
 cttgtgccgg ttttcaaagg gaatgcttcc agcttttgcc cattcagtat aatattaaag 120
 aatgttttac cattttctgt cttgcctgtt tttctgtgtt tttgttggtc tcttcattct 180
 ccatttttag gcctttacat gtttaggaata tatttctttt aatgatactt cacctttggt 240
 atcttttgtg agactctact catagtgtga taagcactgg gttggttaagg caagaggagc 300

<210> 65
 <211> 203
 <212> DNA
 <213> Homo sapien

<400> 65
 gctcctcttg ccttaccac tcacccagta tgcagcaat tttatcrgct ttacctacga 60
 aacagcctgt atccaaacac ttaacacact cacctgaaaa gttcaggcaa caatcgctt 120
 ctcatgggtc tctctgctcc agttctgaac ctttctcttt tcctagaaca tgcatttarg 180
 tcgatagaag ttcctctcag tgc 203

<210> 66
 <211> 344
 <212> DNA
 <213> Homo sapien

<400> 66
 tacggggacc cctgcattga gaaagcgaga ctcactctga agctgaaatg ctggtgccct 60
 tgcaagtgtg gtagcaggag ttctgtgctt tgtgggctaa ggctcctgga tgacccctga 120
 catggagaag gcagagttgt gtgccccttc tcatggcctc gtcaaggcat catggactgc 180
 cacacacaaa atgccgtttt tattaacgac atgaaattga aggagagaac acaattcact 240
 gatgtggctc gtaaccatgg atatggtcac atacagaggt gtgattatgt aaaggttaat 300
 tccaccacc tcatgtggaa actagcctca atgcaggggt ccca 344

<210> 67
 <211> 157
 <212> DNA
 <213> Homo sapien

<400> 67
 gcactgagag gaacttcgta gggaggttga actggctgct gaggaggggg aacaacaggg 60
 taaccagact gatagccatt ggatggataa tatggtggtt gaggaggagc actacttata 120
 gcagagggtt gtgtatagcc tgaggaggca tcaccgc 157

<210> 68

<211> 137
 <212> DNA
 <213> Homo sapien

<400> 68
 gcactgagag gaacttctag aaagtgaaag tctagacata aaataaaata aaaattttaa 60
 actcaggaga gacagcccag cacggtggct cagcctgta atcccagaac ttggggagcc 120
 tgaggaggca tcacccg 137

<210> 69
 <211> 137
 <212> DNA
 <213> Homo sapien

<400> 69
 cgggtgatgc ctctcaggc tgtatcttga agactatcga ctggacttct tatcaactga 60
 agaatccggt aaaaatacca gttgtattat ttctacctgt caaaatccat ttcaaagtgt 120
 gaagttcctc tcagtgc 137

<210> 70
 <211> 220
 <212> DNA
 <213> Homo sapien

<220>
 <221> misc_feature
 <222> (1)...(220)
 <223> n = A,T,C or G

<400> 70
 agcatgttga gccagacac gcaatctgaa tgagtgtgca cctcaagtaa atgtctacac 60
 gctgcctggt ctgacatggc acaccatcnc gtggagggca casctctgct cngcctacwa 120
 cgagggcant ctcatwgaca gggtccaccc accaaactgc aagaggctca nnaagtactr 180
 ccagggtmya sggacmasgg tgggaytyca ycacwcatct 220

<210> 71
 <211> 353
 <212> DNA
 <213> Homo sapien

<220>
 <221> misc_feature
 <222> (1)...(353)
 <223> n = A,T,C or G

<400> 71
 cgttagggtc tctatccact gctaaacat acacctgggt aaacaggac catttaacat 60
 tcccanctaa atatgccaaag tgacttcaca tgtttatctt aaagatgtcc aaaacgcaac 120
 tgattttctc cctaaacct gtgatgggtg gatgattaan cctgagtggt ctacagcaag 180
 ttaagtgcaa ggtgctaaat gaangtgacc tgagatacag catctacaag gcagtacctc 240
 tcaacncagg gcaactttgc ttctcanagg gcatttagca gtgtctgaag taatttctgt 300
 attacaactc acggggcggg ggggtgaatat ctantggana gnagacccta acg 353

<210> 72
 <211> 343
 <212> DNA
 <213> Homo sapien

<400> 72
gcaactgagag gaacttccaa tacyatkac agagtgaaca rgcarccyac agaacaggag 60
aaaatgttyg caatctctcc atctgacaaa aggctaatat ccagawtcta awaggaactt 120
aaacaaattt atgagaaaag aacaracaac ctcaawcaaaa agtgggtgaa ggawatgcts 180
aaargaagac atytattcag ccagtaaaca yatgaaaaaa aggctcatsa tcaactgawca 240
ttagagaaat gcaaatacaa accacaatga gataccatct yayrccagtt agaayggtga 300
tcattaaaaar stcaggaaac aacagatgct ggacaagggtg tca 343

<210> 73

<211> 321

<212> DNA

<213> Homo sapien

<220>

<221> misc_feature

<222> (1)...(321)

<223> n = A,T,C or G

<400> 73
gcaactgagag gaacttcaga gagagagaga gagttccacc ctgtacttgg ggagagaaac 60
agaagggtgag aaagtctttg gttctgaagc agcttctaag atcttttcat ttgcttcatt 120
tcaaagttcc catgctgcc aagtgccatc ctttggggta ctgttttctg agctccagtg 180
ataactcatt tatacaagg agataccag aaaaaaagtg agcaaatctt aaaaagggtg 240
cttgagttca gccttaata ccatcttgaa atgacacaga gaaagaanga tgttgggtgg 300
gagtgatag agaccctaac g 321

<210> 74

<211> 321

<212> DNA

<213> Homo sapien

<400> 74
gcaactgagag gaacttcaga gagagagaga gagttccacc ctgtacttgg ggagagaaac 60
agaagggtgag aaagtctttg gttctgaagc agcttctaag atcttttcat ttgcttcatt 120
tcaaagttcc catgctgcc aagtgccatc ctttggggta ctgttttctg agctccagtg 180
ataactcatt tatacaagg agataccag aaaaaaagtg agcaaatctt aaaaagggtg 240
cttgagttca gyccttaata ccatcttgaa atgamacaga gaaagaagga tgttgggtgg 300
gagtgatag agaccctaac g 321

<210> 75

<211> 317

<212> DNA

<213> Homo sapien

<400> 75
gcaactgagag gaacttccac atgcactgag aaatgcatgt tcacaaggac tgaagtctgg 60
aactcagttt ctcaagttcca atcctgattc aggtgtttac cagctacaca accttaagca 120
agtcagataa ccttagcttc ctcatatgca aaatgagaat gaaaagtact catcgctgaa 180
ttgttttgag gattagaaaa acatctggca tgcagtagaa attcaattag tattcatttt 240
cattcttcta aattaaacaa ataggatttt tagtggtgga acttcagaca ccagaaatgg 300
gagtgatag agaccct 317

<210> 76

<211> 244

<212> DNA

<213> Homo sapien

<400> 76

```

cgttagggtc tctatccact cccactactg atcaaaactct atttatttaa ttatttttat    60
catacttttaa gttctgggat acacgtgcag catgcgcagg tttgttgcat aggtatacac    120
ttgccatggt gggttgctgc acccatcagt ccatcatcta cattaggat ttctcctaata    180
gctatccctc ccctagcccc ttacaccccc aacaggctct agtgtgtgaa gttcctctca    240
gtgc                                                                    244

```

```

<210> 77
<211> 254
<212> DNA
<213> Homo sapien

```

```

<400> 77
cgttagggtc tctatccact gaaatctgaa gcacaggagg aagagaagca gtyctagtga    60
gatggcaagt tcwtttacca cactcttttaa catttygttt agttttaacc tttatttatg    120
gataataaag gttaatatta ataatgattt attttaaggc attcccraat ttgcataatt    180
ctccttttgg agataccctt ttatctccag tgcaagtctg gatcaaagtg atasamagaa    240
gttcctctca gtgc                                                                    254

```

```

<210> 78
<211> 355
<212> DNA
<213> Homo sapien

```

```

<220>
<221> misc_feature
<222> (1)...(355)
<223> n = A,T,C or G

```

```

<400> 78
ttcgatacag gcaaacatga actgcaggag ggtggtgacg atcatgatgt tgccgatggt    60
ccggatggnc acgaagacgc actggancac gtgcttacgt ccttttgctc tgttgatggc    120
cctgagggga cgcaggaccc ttatgaccct cagaatcttc acaacgggag atggcactgg    180
attgantccc antgacacca gagacacccc aaccaccagn atatcantat attgatgtag    240
ttcctgtaga nggccccctt gtggaggaaa gctccatnag ttggtcatct tcaacaggat    300
ctcaacagtt tccgatggct gtgatgggca tagtcatant taaccntgtn tcgaa        355

```

```

<210> 79
<211> 406
<212> DNA
<213> Homo sapien

```

```

<400> 79
taagagggta ccagcagaaa ggtagtatc atcagatagc atcttatacg agtaatatgc    60
ctgctatattg aagtgttaatt gagaaggaaa attttagcgt gctcactgac ctgcctgtag    120
ccccagtgc agctaggatg tgcattctcc agccatcaag agactgagtc aagttgttcc    180
ttaagtcala acagcagact cagctctgac attctgattc gaatgacact gttcaggaat    240
cggaatcctg tcgattagac tggacagctt gtggcaagtg aatttgccctg taacaagcca    300
gattttttta aatttatatt gtaaataatg tgtgtgtgtg tgtgtgtata tatatatata    360
tgtacagtta tctaagttta tttaaaagtt gtttggtacc ctctta                    406

```

```

<210> 80
<211> 327
<212> DNA
<213> Homo sapien

```

```

<400> 80
tttttttttt ttactcggc tcagtctaata cctttttgta gtcactcata ggccagactt    60
agggtctagga tgatgattaa taagagggat gacataacta ttagtggcag gttagtgtgt    120

```

```

tgtagggctc atggtagggg taaaaggagg gcaatttcta gatcaaataa taagaaggta      180
atagctacta agaagaattt tatggagaaa gggacgcggg cgggggatat agggtcgaag      240
ccgcactcgt aaggggtgga tttttctatg tagccgttga gttgtggtag tcaaatgta      300
ataattatta gtagtaagcc taggaga                                           327

```

<210> 81
 <211> 318
 <212> DNA
 <213> Homo sapien

```

<400> 81
tagtctatgc ggttgattcg gcaatccatt atttgctgga ttttgtcatg tgttttgcc      60
attgcattca taatttatta tgcatttatg ctgttatctc ctaagtcatg gtatataatc      120
catgcttttt atgttttgtc tgacataaac tottatcaga gccctttgca cacagggatt      180
caataaatat taacacagtc tacatttatt tgggtgaatat tgcatatctg ctgtactgaa      240
agcacattaa gtaacaaagg caagtggagaa gaatgaaaag cactactcac aacagttatc      300
atgattgcgc atagacta                                           318

```

<210> 82
 <211> 338
 <212> DNA
 <213> Homo sapien

```

<400> 82
tcttcaacct ctactccac taatagcttt ttgatgactt ctagcaagcc tcgctaacct      60
cgcttacc cccactatta acctactggg agaactctct gtgctagtaa ccacgttctc      120
ctgatcaaat atcactctcc tacttacagg actcaacata ctagtacag ccctatactc      180
cctctacata ttaccacaa cacaatgggg ctactcacc caccacatta acaacataaa      240
accctcattc acacgagaaa acaccctcat gttcatacac ctatcccca ttctcctcct      300
atccctcaac ccgacatca ttaccgggtt ttcctctt                                           338

```

<210> 83
 <211> 111
 <212> DNA
 <213> Homo sapien

```

<400> 83
agccatttac ccccatcca caaaaaaaaaa aaaaaaaaag aaaaatatca aggaataaaa      60
atagactttg aacaaaaagg aacatttgct ggcctgagga ggcacaccc g                                           111

```

<210> 84
 <211> 224
 <212> DNA
 <213> Homo sapien

```

<400> 84
tcgggtgatg cctcctcagg ccaagaagat aaagcttcag acccctaaca catttccaaa      60
aaggaagaaa ggagaaaaaa gggcatcatc cccgttcoga agggtcaggg aggaggaaat      120
tgaggtgatg tcacgagttg cggacaactc ctttgatgcc aagcgaggtg cagccggaga      180
ctggggagag cgagccaatc aggttttgaa gttcctctca gtgc                                           224

```

<210> 85
 <211> 348
 <212> DNA
 <213> Homo sapien

```

<400> 85
gcactgagag gaacttcgtt ggaaacgggt ttttttcatg taaggctaga cagaagaatt      60

```

```

ctcagtaact tccttgtggt gtgtgtattc aactcacasa gttgaacgat cctttacaca 120
gagcagactt gtaacactct twttgtggaa tttgcaagtg gagatttcag scgctttgaa 180
gtsaaaggta gaaaaggaaa tatcttccta taaaaactag acagaatgat tctcagaaac 240
tccttttgta tgtgtgcggt caactcacag agtttaacct ttcwtttcat agaagcagtt 300
aggaaacact ctgtttgtaa agtctgcaag tggatagaga ccctaacg 348

```

<210> 86

<211> 293

<212> DNA

<213> Homo sapien

<400> 86

```

gcactgagag gaacttcytc gtgwtgktgc yattcaactc acagagttga asswtsmttt 60
acabagwkca ggcttkcaaa cactcttttt gtmgaatytc caagwggaka tttsrrccrc 120
tttgwggycw wysktmgaaw mggrwatatc ttcwyatmra amctagacag aaksattctc 180
akaawstyyy ytgtagawgs tgcrttcaac tcacagagkt kaacmwtyct kytsatrgag 240
cagttwkgaa actctmtttc tttggattct gcaagtggat agagacccta acg 293

```

<210> 87

<211> 10

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer for amplification from breast tumor cDNA

<400> 87

```

ctcctaggt 10

```

<210> 88

<211> 10

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer for amplification from breast tumor cDNA

<400> 88

```

agtagttgcc 10

```

<210> 89

<211> 11

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer for amplification from breast tumor cDNA

<400> 89

```

ttccgttatg c 11

```

<210> 90

<211> 10

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer for amplification from breast tumor cDNA

<400> 90
tggtaaaggg 10

<210> 91
<211> 10
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer for amplification from breast tumor cDNA

<400> 91
tcggtcatag 10

<210> 92
<211> 10
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer for amplification from breast tumor cDNA

<400> 92
tacaacgagg 10

<210> 93
<211> 10
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer for amplification from breast tumor cDNA

<400> 93
tggattggtc 10

<210> 94
<211> 10
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer for amplification from breast tumor cDNA

<400> 94
ctttctaccc 10

<210> 95
<211> 10
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer for amplification from breast tumor cDNA

<400> 95
ttttggctcc 10

<210> 96
<211> 10
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer for amplification from breast tumor cDNA

<400> 96
ggaaccaatc 10

<210> 97
<211> 10
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer for amplification from breast tumor cDNA

<400> 97
tcgatacagg 10

<210> 98
<211> 10
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer for amplification from breast tumor cDNA

<400> 98
ggtactaagg 10

<210> 99
<211> 10
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer for amplification from breast tumor cDNA

<400> 99
agtctatgcg 10

<210> 100
<211> 10
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer for amplification from breast tumor cDNA

<400> 100
ctatccatgg 10

<210> 101
<211> 10

<212> DNA
<213> Artificial Sequence

<220>
<223> Primer for amplification from breast tumor cDNA

<400> 101
tctgtccaca 10

<210> 102
<211> 10
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer for amplification from breast tumor cDNA

<400> 102
aagagggtac 10

<210> 103
<211> 10
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer for amplification from breast tumor cDNA

<400> 103
cttcaacctc 10

<210> 104
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer for amplification from breast tumor cDNA

<400> 104
gctcctcttg ccttaccaac 20

<210> 105
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer for amplification from breast tumor cDNA

<400> 105
gtaagtcgag cagtgtgatg 20

<210> 106
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer for amplification from breast tumor cDNA

<400> 106
gtaagtcgag cagtctgatg 20

<210> 107
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer for amplification from breast tumor cDNA

<400> 107
gacttagtg aaagaatgta 20

<210> 108
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer for amplification from breast tumor cDNA

<400> 108
gtaattccgc caaccgtagt 20

<210> 109
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer for amplification from breast tumor cDNA

<400> 109
atggttgatc gatagtggaa 20

<210> 110
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer for amplification from breast tumor cDNA

<400> 110
acggggaccc ctgcattgag 20

<210> 111
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer for amplification from breast tumor cDNA

<400> 111
tattctagac cattcgctac 20

<210> 112
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer for amplification from breast tumor cDNA

<400> 112
acataaccac ttttagcggtc 20

<210> 113
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer for amplification from breast tumor cDNA

<400> 113
cgggtgatgc ctctcaggc 20

<210> 114
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer for amplification from breast tumor cDNA

<400> 114
agcatgttga gccagacac 20

<210> 115
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer for amplification from breast tumor cDNA

<400> 115
gacaccttgt ccagcatctg 20

<210> 116
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer for amplification from breast tumor cDNA

<400> 116
tacgctgcaa cactgtggag 20

<210> 117
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer for amplification from breast tumor cDNA

<400> 117
cgttagggtc tctatccact 20

<210> 118
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer for amplification from breast tumor cDNA

<400> 118
agactgactc atgtccccta 20

<210> 119
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer for amplification from breast tumor cDNA

<400> 119
tcacgcgtcg gtgactcaag 20

<210> 120
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer for amplification from breast tumor cDNA

<400> 120
caagattcca taggctgacc 20

<210> 121
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer for amplification from breast tumor cDNA

<400> 121
acgtactggc cttgaagtc 20

<210> 122
<211> 20
<212> DNA

<213> Artificial Sequence

<220>

<223> Primer for amplification from breast tumor cDNA

<400> 122
gacgcttggc cacttgacac 20

<210> 123
<211> 20
<212> DNA
<213> Artificial Sequence

<220>

<223> Primer for amplification from breast tumor cDNA

<400> 123
gtatcgacgt agtggctctcc 20

<210> 124
<211> 20
<212> DNA
<213> Artificial Sequence

<220>

<223> Primer for amplification from breast tumor cDNA

<400> 124
tagtgacatt acgacgctgg 20

<210> 125
<211> 20
<212> DNA
<213> Artificial Sequence

<220>

<223> Primer for amplification from breast tumor cDNA

<400> 125
cgggtgatgc ctcctcaggc 20

<210> 126
<211> 23
<212> DNA
<213> Artificial Sequence

<220>

<223> Primer for amplification from breast tumor cDNA

<400> 126
atggetattt tcgggggctg aca 23

<210> 127
<211> 22
<212> DNA
<213> Artificial Sequence

<220>

<223> Primer for amplification from breast tumor cDNA

<400> 127
ccggtatctc ctcgtgggta tt 22

<210> 128
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer for amplification from breast tumor cDNA

<400> 128
ctgcctgagc cacaaatg 18

<210> 129
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer for amplification from breast tumor cDNA

<400> 129
ccggaggagg aagctagagg aata 24

<210> 130
<211> 14
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer

<400> 130
tttttttttt ttag 14

<210> 131
<211> 18
<212> PRT
<213> Artificial Sequence

<220>
<223> Predicited Th Motifs (B-cell epitopes)

<400> 131
Ser Ser Gly Gly Arg Thr Phe Asp Asp Phe His Arg Tyr Leu Leu Val
1 5 10 15
Gly Ile

<210> 132
<211> 22
<212> PRT
<213> Artificial Sequence

<220>

<223> Predicited Th Motifs (B-cell epitopes)

<221> VARIANT

<222> (1)...(22)

<223> Xaa = Any Amino Acid

<400> 132

Gln	Gly	Ala	Ala	Gln	Lys	Pro	Ile	Asn	Leu	Ser	Lys	Xaa	Ile	Glu	Val
1				5				10					15		
Val	Gln	Gly	His	Asp	Glu										
			20												

<210> 133

<211> 23

<212> PRT

<213> Artificial Sequence

<220>

<223> Predicited Th Motifs (B-cell epitopes)

<400> 133

Ser	Pro	Gly	Val	Phe	Leu	Glu	His	Leu	Gln	Glu	Ala	Tyr	Arg	Ile	Tyr
1				5				10						15	
Thr	Pro	Phe	Asp	Leu	Ser	Ala									
			20												

<210> 134

<211> 9

<212> PRT

<213> Artificial Sequence

<220>

<223> Predicited HLA A2.1 Motifs (T-cell epitopes)

<400> 134

Tyr	Leu	Leu	Val	Gly	Ile	Gln	Gly	Ala
1				5				

<210> 135

<211> 9

<212> PRT

<213> Artificial Sequence

<220>

<223> Predicited HLA A2.1 Motifs (T-cell epitopes)

<400> 135

Gly	Ala	Ala	Gln	Lys	Pro	Ile	Asn	Leu
1				5				

<210> 136

<211> 9

<212> PRT

<213> Artificial Sequence

<220>

<223> Predicited HLA A2.1 Motifs (T-cell epitopes)

<221> VARIANT
<222> (1)...(9)
<223> Xaa = Any Amino Acid

<400> 136
Asn Leu Ser Lys Xaa Ile Glu Val Val
1 5

<210> 137
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Predicited HLA A2.1 Motifs (T-cell epitopes)

<400> 137
Glu Val Val Gln Gly His Asp Glu Ser
1 5

<210> 138
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Predicited HLA A2.1 Motifs (T-cell epitopes)

<400> 138
His Leu Gln Glu Ala Tyr Arg Ile Tyr
1 5

<210> 139
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Predicited HLA A2.1 Motifs (T-cell epitopes)

<400> 139
Asn Leu Ala Phe Val Ala Gln Ala Ala
1 5

<210> 140
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Predicited HLA A2.1 Motifs (T-cell epitopes)

<400> 140
Phe Val Ala Gln Ala Ala Pro Asp Ser
1 5

<210> 141
<211> 9388

<212> DNA

<213> Homo sapien

<400> 141

gctcgcggcc	gcgagctcaa	ttaaccctca	ctaaagggag	tcgactcgat	cagactgtta	60
ctgtgtctat	gtagaaagaa	gtagacataa	gagattccat	tttgttctgt	actaagaaaa	120
attcttctgc	cttgagatgc	tgtaaatctg	taaccctagc	cccaaccctg	tgctcacaga	180
gacatgtgct	gtgttgactc	aaggttcaat	ggatttaggg	ctatgctttg	ttaaaaaagt	240
gcttgaagat	aatatgcttg	ttaaaagtca	tcaccattct	ctaattctcaa	gtacccaggg	300
acacaatata	ctcgggaagg	ccgcagggac	ctctgtctag	gaaagccagg	tattgtccaa	360
gatttctccc	catgtgatag	cctgagatat	ggcctcatgg	gaagggtaag	acctgactgt	420
ccccagccc	gacatccccc	agcccagacat	ccccagccc	gacacccgaa	aaggtctgt	480
gctgaggagg	attagtaaaa	gaggaaggcc	tctttgcagt	tgaggtaaga	ggaaggcatc	540
tgtctcctgc	tcgtccctgg	gcaatagaat	gtcttggtgt	aaaacccgat	tgtatgttct	600
acttactgag	ataggagaaa	acatccttag	ggctggagggt	gagacacgct	ggcggcaata	660
ctgctcttta	atgcaccgag	atgtttgtat	aagtgcacat	caaggcacag	cacctttcct	720
taaaacttatt	tatgacacag	agacctttgt	tcacgttttc	ctgctgacct	tctcccact	780
attaccctat	tggcctgcca	catccccctc	tccgagatgg	tagagataat	gatcaataaa	840
tactgaggga	actcagagac	cagtgtccct	gtaggtccct	cgtgtgctga	gcgccggctc	900
cttgggctca	cttttctttc	tctatacttt	gtctctgtgt	ctctttcttt	tctcagtctc	960
tcgttccacc	tgacgagaaa	taccacacag	tgtggagggg	caggccaccc	cttcaataat	1020
ttactagcct	gttcgctgac	aacaagactg	gtggtgcaga	aggttgggtc	ttggtgttca	1080
ccgggtggca	ggcatgggcc	aggtgggagg	gtctccagcg	cctggtgcaa	atctccaaga	1140
aagtgcagga	aacagcacca	aggttgattg	taaattttga	tttggcgcgg	caggtagcca	1200
ttccagcgca	aaaatgcgca	ggaaagcttt	tgctgtgctt	gtaggcaggt	aggccccaag	1260
cacttcttat	tggctaattg	ggagggaacc	tgacacatcca	ttggctgaaa	tctccgtcta	1320
tttgaggctg	actgagcgcg	ttcctttctt	ctgtgttgcc	tggaaacgga	ctgtctgcct	1380
agtaacatct	gatcacgttt	cccattggcc	gccgtttccg	gaagcccgc	ctcccatttc	1440
cggaagcctg	gcgcaagggt	ggtctgcagg	tggcctccag	gtgcaaagtg	ggaagtgtga	1500
gtcctcagtc	ttgggctatt	cggccacgtg	cctgccggac	atgggacgct	ggagggtcag	1560
cagcgtggag	tcctggcctt	ttgcgtccac	gggtgggaaa	ttggccattg	ccacggcggg	1620
aactgggact	caggctgccc	cccggccggt	tctcatccgt	ccaccggact	cgtgggcgct	1680
cgcactggcg	ctgatgtagt	ttcctgacct	ctgaccgcta	ttgtctccag	attaaaggta	1740
aaaacggggc	tttttcagcc	cactcgggta	aaacgccttt	tgattttctag	gcagggtgtt	1800
tgttgacgc	ctgggagggg	gtgaccgca	ggttgagggt	tattaaaata	cattcctggg	1860
ttatgttatg	tttataataa	agcaccctaa	cctttacaaa	atctcacttt	ttgccagttg	1920
tattatttag	tggactgtct	ctgataagga	cagccagtta	aaatggaatt	ttgttgttgc	1980
taattaaacc	aatttttagt	tttgggtgtt	gtcctaatag	caacaacttc	tcaggcttta	2040
taaaaccata	tttcttgggg	gaaatttctg	tgtaaggcac	agcgagttag	tttgggaattg	2100
ttttaaagga	agtaagttcc	tggttttgat	atcttagtag	tgtaatgccc	aacctgggtt	2160
ttactaacc	tgtttttaga	ctctcccttt	ccttaaatca	cctagccttg	tttccacctg	2220
aattgactct	cccttagcta	agagcgccag	atggactcca	tcttggtctt	ttcactggca	2280
gccccttcc	caaggactta	acttgtgcaa	gctgactccc	agcacatcca	agaatgcaat	2340
taactgttaa	gatactgtgg	caagctatat	cgcgagttcc	gaggaattca	tccgattgat	2400
tatgcccata	agccccgcgt	ctatcacctt	gtaataatct	taaagcccct	gcacctggaa	2460
ctattaactt	tcctgtaacc	atttatcctt	ttactttttt	tgcttacttt	atttctgtaa	2520
aattgtttta	actagacctc	ccctcccttt	tctaaaccaa	agtataaaaag	aagatctagc	2580
cccttcttca	gagcggagag	aattttgagc	attagccatc	tcttggcggc	cagctaaata	2640
aatggacttt	taatttgtct	caaagtgtgg	cgttttctct	aactcgctca	ggtacgacat	2700
ttggaggccc	cagcgagaaa	cgtcaccggg	agaaacgtca	ccgggcgaga	gccgggcccg	2760
ctgtgtgctc	ccccggaagg	acagccagct	tgtagggggg	agtgccacct	gaaaaaaaaa	2820
tttccaggtc	cccaaagggt	gaccgtcttc	cggaggacag	cggatcgact	accatgcggg	2880
tgcccaccaa	aattccacct	ctgagtccct	aactgtgac	cccggggtca	ggtaggtcag	2940
atttgacttt	ggttctggca	gaggggaagc	accctgatga	gggtgtccct	cttttgactc	3000
tgcccatttc	tctaggatgc	tagagggtag	agccctgggt	ttctgttaga	cgcctctgtg	3060
tctctgtctg	ggagggaagt	ggccctgaca	ggggccatcc	cttgagtcag	tccacatccc	3120
aggatgctgg	gggactgagt	cctggtttct	ggcagactgg	tctctctctc	tctctttttc	3180
tatctcta	ctttccttgt	tcaggtttct	tggagaatct	ctgggaaaaga	aaaaagaaaa	3240

actgttataa	actctgtgtg	aatggtgaat	gaatggggga	ggacaagggc	ttgcgcttgt	3300
cctccagttt	gtagctccac	ggcgaagct	acggagtcca	agtgggccct	cacctgcggt	3360
tccgtggcga	cctcataagg	cttaaggcag	catccggcat	agctcgatcc	gagccggggg	3420
tttataccgg	cctgtcaatg	ctaagaggag	cccaagtccc	ctaaggggga	gcggccaggc	3480
gggcatctga	ctgatcccat	cacgggaccc	cctccccttg	tttgtctaaa	aaaaaaaaaa	3540
gaagaaactg	tcataactgt	ttacatgccc	tagggtcaac	tgtttgtttt	atgtttattg	3600
ttctgttcgg	tgtctattgt	cttgttttagt	ggtgtgcaag	gttttgcatg	tcaggacgtc	3660
gatattgccc	aagacgtctg	ggtaagaact	tctgcaaggt	ccttagtgct	gattttttgt	3720
cacaggaggt	taaattttct	atcaatcatt	taggctggcc	accacagtcc	tgtcttttct	3780
gccagaagca	agtcagggtg	tgttacggga	atgagtgtaa	aaaaacattc	gcctgattgg	3840
gattttctgg	accatgatgg	ttgtatttag	attgtcatac	cccacatcca	ggttgattgg	3900
acctcctcta	aactaaactg	gtgggtgggt	caaaacagcc	accctgcaga	tttccttgct	3960
cacctctttg	gtactctctg	aacttttcct	gtgcccttaa	atagcacact	gtgtagggaa	4020
acctaccctc	gtactgcttt	acttcgttta	gattcttact	ctgttcctct	gtggctactc	4080
tcccatctta	aaaacgatcc	aagtggtcct	tttcctcctc	cctgccccct	acccacaca	4140
tctcgttttc	cagtgcgaca	gcaagttcag	cgtctccagg	acttggtctc	gctctcactc	4200
cttgaaccct	taaaagaaaa	agctgggttt	gagctatttg	cctttgagtc	atggagacac	4260
aaaagggtatt	taggggtacag	atctagaaga	agagagagaa	cacctagatc	caactgaccc	4320
aggagatctc	gggctggcct	ctagtccctc	tccctcaatc	ttaaagctac	agtgatgtgg	4380
caagtgggat	ttagctgttg	tggtttttct	gctctttctg	gtcatgttga	ttctgttctt	4440
tcgatactcc	agccccccag	ggagtgtgtt	tctctgtctg	tgctgggttt	gatattctatg	4500
ttcaaattct	attaaattgc	cttcaaaaaa	aaaaaaaaaa	gggaaacact	tcctcccagc	4560
cttgtaaggg	ttggagccct	ctccagtata	ctctgcagaa	tttttctctc	ggtttctcag	4620
aggattatgg	agtcgcgcct	aaaaaaggca	agctctggac	actctgcaaa	gtagaatggc	4680
caaagtttgg	agttgagtgg	ccccttgaag	ggtcactgaa	cctcacaatt	gttcaagctg	4740
tgtggcgggg	tgttactgaa	actcccggcc	tccctgatca	gtttccctac	attgatcaat	4800
ggctgagttt	ggtcaggagc	accccttcca	tggctccact	catgcaccat	tcataatttt	4860
acctccaagg	tctctctgag	ccagaccgtg	ttttgcctc	gacctcagc	cggttcagct	4920
cgccctgtac	tgcctctctc	tgaagaagag	gagagtctcc	ctcaccaggt	cccaccgct	4980
taaaaccagc	ctactccctt	agggtcatcc	catgtctcct	cggctatgtc	cctgtaggc	5040
tcatcaccca	ttgcctcttg	gttgcaaccg	tgggtggagg	aagtagcccc	tctactacca	5100
ctgagagagg	cacaagtccc	tctgggtgat	gagtgtctca	cccccttcc	ggtttatgtc	5160
ccttctttct	acttctgact	tgtataattg	gaaaaaccat	aatcctccct	tctctgaaaa	5220
gccccaggct	ttgacctcac	tgtactctgg	tgtactctgg	acacattggc	ccacctggga	5280
tgactgtcaa	cagctccttt	tgacctttt	cacctctgaa	gagagggaaa	gtatccaaag	5340
agaggccaaa	aagtacaacc	tcacatcaac	caataggccg	gaggagggaag	ctagagggaat	5400
agtgattaga	ga'ccaattg	ggacctaat	gggacccaaa	tttctcaagt	ggaggggagaa	5460
cttttgacga	tttccaccgg	tatctcctcg	tgggtattca	gggagctgct	cagaaacctc	5520
taaaacttgc	taagcgact	gaagtcgtcc	acggcgatga	tgatccacca	ggagtgtttt	5580
tagagcacct	ccaggaggct	tatcggattt	acaccccttt	tgacctggca	gccccgaaa	5640
atagccatgc	tcttaatttg	gcatttgtgg	ctcaggcagc	cccagatagt	aaaaggaaac	5700
tccaaaaact	agagggattt	tgctggaatg	aataccagtc	agcttttaga	gatagcctaa	5760
aaggtttttg	acagtcaaga	ggttgaaaaa	caaaaacaag	cagctcaggc	agctgaaaaa	5820
agccactgat	aaagcatcct	ggagtatcag	agtttactgt	tagatcaccc	tcatttgact	5880
tccccctcca	catggtgttt	aaatccagct	acactacttc	ctgactcaaa	ctccactatt	5940
cctgttcatg	actgtcagga	actgttgga	actactgaaa	ctggccgacc	tgatcttcaa	6000
aatgtgcccc	taggaaagg	ggatgccacc	gtgttcacag	acagtgcag	cttcctcgag	6060
aagggactac	gaaaggccgg	tgcagctgtt	accatggaga	cagatgtgtt	gtgggctcag	6120
gctttaccag	caaaacacct	agcacaaaag	ctgaattga	tcgccctcac	tcaggctctc	6180
cgatggggta	aggatattaa	cgttaacact	gacagcaggt	acgcctttgc	tactgtgcat	6240
gtacgtggag	ccatctacca	ggagcgtggg	ctactcacct	cagcaggtgg	ctgtaatcca	6300
ctgtaaagg	catcaaaagg	aaaacacggc	tgttgccctg	ggtaaccaga	aagctgattc	6360
agcagctcaa	gatgcagtgt	gactttcagt	cacgcctcta	aacttgctgc	ccacagtctc	6420
ctttccacag	ccagatctgc	ctgacaatcc	cgcatactca	acagaagaag	aaaactggcc	6480
tcagaactca	gagccaataa	aaatcaggaa	ggttggtgga	ttcttcctga	ctctagaatc	6540
ttcatacccc	gaactcttgg	gaaaacttta	atcagtcacc	tacagtctac	cacccattta	6600
ggaggagcaa	agctacctca	gctcctccgg	agcggtttta	agatcccca	tcttcaaagc	6660
ctaacagatc	aagcagctct	ccggtgcaca	actgcgccc	aggtaaatgc	caaaaaaggt	6720

cctaaaccca	gccaggcca	ccgtctccaa	gaaaactcac	caggagaaaa	gtgggaaatt	6780
gactttacag	aagtaaaacc	acaccgggct	gggtacaaat	accttctagt	actggtagac	6840
accttctctg	gatggactga	agcattttgct	acaaaaaacg	aaactgtcaa	tatggtagtt	6900
aagtttttac	tcaatgaaat	catccctcga	cgtgggctgc	ctgttgccat	agggctctgat	6960
aatggaccgg	ccttcgcctt	gtctatagtt	tagtcagtca	gtaaggcggt	aaacattcaa	7020
tggaagctcc	attgtgccta	tcgacccag	agctctgggc	aagtagaacg	catgaactgc	7080
accctaaaaa	acactcttac	aaaattaatc	ttagaaaccg	gtgtaaattg	tgtaagtctc	7140
cttcctttag	ccctacttag	agtaagggtgc	accccttact	gggctggggt	cttacctttt	7200
gaaatcatgt	atgggagggc	gctgcctatc	ttgcctaagc	taagagatgc	ccaattggca	7260
aaaatatcac	aaactaat	attacagtac	ctacagtctc	cccaacaggt	acaagatatc	7320
atcctgccac	ttgttcgagg	aaccatccc	aatccaattc	ctgaacagac	agggccctgc	7380
cattcattcc	cgccagggtga	cctgttgttt	gttaaaaagt	tccagagaga	aggactccct	7440
cctgcttgga	agagacctca	caccgtcatc	acgatgccaa	cggctctgaa	ggtggatggc	7500
attcctgcgt	ggattcatca	ctcccgcatc	aaaaaggcca	acggagccca	actagaacaa	7560
tggtgcccc	gggctgggtc	aggccctta	aaactgcacc	taagttgggt	gaagccatta	7620
gattaattct	ttttctta	tttgtaaaac	aatgcatagc	ttctgtcaaa	cttatgtatc	7680
ttaagactca	atataacccc	cttgttataa	ctgaggaatc	aatgatttga	ttccccaaaa	7740
acacaagtgg	ggaatgtagt	gtccaacctg	gtttttacta	accctgtttt	tagactctcc	7800
ctttccttta	atcactcagc	cttgtttcca	cctgaattga	ctctccctta	gctaagagcg	7860
ccagatggac	tccatcttgg	ctctttcact	ggcagccgct	tcctcaagga	cttaacttgt	7920
gcaagctgac	tcccagcaca	tccaagaatg	caattaactg	ataagatact	gtggcaagct	7980
atatccgcag	ttcccaggaa	ttcgtccaat	tgattacacc	caaaagcccc	gcgtctatca	8040
ccttgaata	atcttaaagc	ccctgcacct	ggaactatta	acgttctctg	aaccatttat	8100
ccttttaact	tttttgcccta	ctttatttct	gtaaaattgt	tttaactaga	ccccccctct	8160
cctttctaaa	caaaggtata	aaagcaaatc	tagccccttc	ttcaggccga	gagaatttcg	8220
agcgtaggac	gtctcttggc	caccagctaa	ataaacggat	tcttcatgtg	tctcaaagtg	8280
tggcgttttc	tctaactcgc	tcaggtagca	ccgtggtagt	atthtcccca	acgtcttatt	8340
tttagggcac	gtatgtagag	taacttttat	gaaagaaacc	agttaaggag	gttttgggat	8400
ttcctttatc	aactgttaata	ctggttttga	ttattttatt	attttattat	tttttttgag	8460
aaggagtttc	actcttgttg	cccaggctgg	agtgcattgg	tgcatcttgg	gctcactgca	8520
acttccgctc	cccaggttca	agcgattctc	ctgcctcagc	ctcgagagta	gctgggatta	8580
taggcattgc	ccaccacacc	cagctaattt	tgatttttta	gtaaagatgg	ggtttcttca	8640
tggtgggtcaa	gctgggtctg	aactccccgc	ctcgggtgat	ctgcccgcct	cggcctccga	8700
aagtgcgtgg	attacaggtg	tgatccacca	caccagccg	atttatatgt	atataaatca	8760
cattcctcta	acaaaaatgt	agtgtttcct	tccatcttga	atataggctg	tagaccocgt	8820
gggtatggga	cattgttaac	agtgcagcca	cagcagtttt	tatgtcatct	gacagcatct	8880
ccaaatagcc	ttcatgggtg	tcaactgcttc	ccaagacaat	tccaaataac	acttcccagt	8940
gatgacttgc	tacttgctat	tggtacttaa	tggtgttaag	tggtgtttac	agacactatt	9000
agtatgtcag	gaattacacc	aaaatttagt	ggctcaaaac	atcattttat	tatgtatgtg	9060
gattctcatg	gtcagggtcag	gatttcagac	agggcacaag	ggtagccac	ttgtctctgt	9120
ctatgatgtc	tgccctcagc	acaggagact	caacagctgg	ggtctgggac	catttggagg	9180
cttgttccct	cacatctgat	acctggcctt	ggatgttggg	agaggggggt	agctgagact	9240
gagtgcctat	atgtagtgtt	tccatattgg	cttgacttcc	ttacagcctg	gcagcctcag	9300
ggtagtcaga	attcttagga	ggcacagggc	tccagggcag	atgctgaggg	gtcttttatg	9360
aggtagcaca	gcaaatccac	ccaggatc				9388

<210> 142

<211> 419

<212> DNA

<213> Homo sapien

<400> 142

tgtaagtcca	gcagtgatg	ggaaggaatg	gtctttggag	agagcatatc	catctcctcc	60
tcaactgcctc	ctaattgcat	gaggtacact	gagcagaatt	aaacagggtg	gtcttaacca	120
cactattttt	agctaccttg	tcaagcta	ggttaaagaa	cacttttggg	ttacacttgt	180
tggtcatag	aagttgctt	ccgccatcac	gcaataagtt	tgtgtgta	cagaaggagt	240
taccttatgg	tttcagtgtc	attctttagt	taacttggga	gctgtgta	ttaggctttg	300
cgtattattt	cacttctgtt	ctccacttat	gaagtgtatt	tgtgttcgcg	tgtgtgtgcg	360

tgcgcatgtg cttccggcag ttaacataag caaataccca acatcacact gctcgactt 419

<210> 143

<211> 402

<212> DNA

<213> Homo sapien

<400> 143

tgtaagtcga	gcagtgatgat	gtccactgca	gtgtgttgct	gggaacagtt	aatgagcaaa	60
ttgtatacaa	tggttagtac	attgaccggg	atgtgttgaa	gctgggtgagt	gttatgactt	120
agcctgttag	actagtctat	gcacatggct	ctgggtcaact	accgctctct	cattttctcca	180
gataaatccc	ccatgcttta	tattctcttc	caaacatact	atcctcatca	ccacatagtt	240
cctttgttaa	tgtttgttc	tagactttcc	ctttctgttt	ttcttattca	aacctatatac	300
tctttgcata	gattgtaaat	tcaaatgccc	tcagggtgca	ggcagttcat	gtaagggagg	360
gaggctagcc	agttagatct	gcatcacact	gctcgactta	ca		402

<210> 144

<211> 224

<212> DNA

<213> Homo sapien

<400> 144

tcgggtgatg	cctcctcagg	ccaagaagat	aaagcttcag	acccctaaca	catttccaaa	60
aagggaagaaa	ggagaaaaaa	gggcatcatc	cccgttccga	agggtcaggg	aggaggaaat	120
tgagggtgat	tcacgagttg	cggacaactc	ctttgatgcc	aagcgaggtg	cagccggaga	180
ctggggagag	cgagccaatc	aggttttgaa	gttcctctca	gtgc		224

<210> 145

<211> 111

<212> DNA

<213> Homo sapien

<400> 145

agccattttac	cacccatcca	caaaaaaaaa	aaaaaaaaag	aaaaatatca	aggaataaaa	60
atagactttg	aacaaaaagg	aacatttgct	ggcctgagga	ggcatcaccc	g	111

<210> 146

<211> 585

<212> DNA

<213> Homo sapien

<400> 146

tagcatgttg	agcccagaca	cttgtagaga	gaggaggaca	gttagaagaa	gaagaaaagt	60
ttttaaatgc	tgaaagttag	tataagaaaag	ctttggcttt	ggatgagact	tttaaagatg	120
cagaggatgc	tttgcagaaa	cttcataaat	atatgcaggt	gattccttat	ttcctcctag	180
aaatttagtg	atatttgaaa	taatgcccac	acttaatttt	ctcctgagga	aaactattct	240
acattactta	agtaaggcat	tatgaaaagt	ttctttttag	gtatagtttt	tcctaatttg	300
gtttgacatt	gcttcatagt	gcctctgttt	ttgtccataa	tcgaaagtaa	agatagctgt	360
gagaaaacta	ttacctaaat	ttggtatgtt	gttttgagaa	atgtccttat	aggagctca	420
cctggtggtt	tttaaattat	tgttgctact	ataattgagc	taattataaa	aacctttttg	480
agacatattt	taaattgtct	tttctgtaa	tactgatgat	gatgttttct	catgcatttt	540
cttctgaatt	gggaccattg	ctgctgtgtc	tgggctcaca	tgcta		585

<210> 147

<211> 579

<212> DNA

<213> Homo sapien

<220>
 <221> misc_feature
 <222> (1)...(579)
 <223> n = A,T,C or G

<400> 147
 tagcatgttg agcccagaca ctgggcagcg ggggtggcca cggcagctcc tgccgagccc 60
 aagcgtgttt gtctgtgaag gacctgacg tcacctgcca ggctagggag gggtcfaatgt 120
 ggagtgaatg ttcaccgact ttcgcaggag tgtgcagaag ccagggtgcaa cttggtttgc 180
 ttgtgttcat caccctcaa gatatgcaca ctgctttcca aataaagcat caactgtcat 240
 ctccagatgg ggaagacttt ttctccaacc agcaggcagg tcccatcca ctcacacacc 300
 agcagctcca ccttctcggg cagcaccacg tcctccacct tctgtgtgta cacgggtgatg 360
 atgtcagcaa agccgttctg cangaccagc tgcccgtgtg gctgtgccat ctactggcc 420
 tccaccgctg acaccgctct agcccgcgca tantgtgcac agaanaaatg atgatccagt 480
 cccacagccc acgtccaaga ngactttatc cgtcagggat tctttattct gcaggatgac 540
 ctgtggtatt aattgttcgt gtctgggctc aacatgcta 579

<210> 148
 <211> 249
 <212> DNA
 <213> Homo sapien

<400> 148
 tgacaccttg tccagcatct gcaagccagg aagagagtcc tcaccaagat cccaccccg 60
 ttggcaccag gatcttggac ttccaatctc cagaactgtg agaaataagt atttgtcgct 120
 aaataaatct ttgtggttcc agatatttag ctatagcaga tcagggtgac taagagaaac 180
 cccataagag ttacatactc attaatctcc gtctctatcc ccagggtctca gatgctggac 240
 aaggtgtca 255

<210> 149
 <211> 255
 <212> DNA
 <213> Homo sapien

<400> 149
 tgacaccttg tccagcatct gctatcttgt gactttttta taatagccat tctgactgg 60
 gtgagatggg aactcattgt gggtttgggt tgcatttctc taatgacag tgatattaag 120
 ctttttttaa atatgcttgt tgaccacatg tatatcatct tttgagaagt gtctgttcat 180
 atcctttgcc cactttttta tttttttatc ttgtaaattt gtttaatttc cttacagatg 240
 ctggacaagg tgtca 255

<210> 150
 <211> 318
 <212> DNA
 <213> Homo sapien

<400> 150
 ttacgctgca acactgtgga ggccaagctg ggatcacttc ttcattctaa ctggagagga 60
 gggaagtcca agtccagcag aggggtgggtg ggtagacagt ggcactcaga aatgtcagct 120
 ggacccctgt ccccgcatag gcaggacagc aaggctgtgg ctctccaggg ccagctgaag 180
 aacaggacac tgtctcgcgt gccacaaagc gtcagagact cccatctttg aagcacggcc 240
 ttcttggctc tctgtcactt ccctgttctg ttagagacct ggttatagac aaggcttctc 300
 cacagtgttg cagcgtaa 318

<210> 151
 <211> 323
 <212> DNA
 <213> Homo sapien

<220>
 <221> misc_feature
 <222> (1)...(323)
 <223> n = A,T,C or G

<400> 151
 tnacgcngcn acnntgtaga ganggnaagg cnttccccac attnccccctt catnanagaa 60
 ttattcnacc aagnntgacc natgccnttt atgacttaca tgcnnactnc ntaatctgtn 120
 tcnngcctta aaagcnnntc cactacatgc ntcancactg tntgtgtnac ntcatnaact 180
 gtcngnaata ggggncata actacagaaa tgcanttcac actgcttcca ntgccatcng 240
 cgtgtggcct tncctactct tcttntattc caagtagcat ctctggantg cttccccact 300
 ctccacattg ttgcagcnat aat 323

<210> 152
 <211> 311
 <212> DNA
 <213> Homo sapien

<400> 152
 tcaagattcc ataggctgac cagtccaagg agagttgaaa tcatgaagga gagtctatct 60
 ggagagagct gtagttttga gggttgcaaa gacttaggat ggagtgggtg ggtgtggtta 120
 gtctctaagg ttgattttgt tcataaattt catgccctga atgccttgct tgcctcacc 180
 tgggtccaagc cttagtgaac acctaaaagt ctctgtcttc ttgctctcca aacttctcct 240
 gaggatttcc tcagattgtc tacattcaga tcgaagccag ttggcaaaca agatgcagtc 300
 cagagggtca g 311

<210> 153
 <211> 332
 <212> DNA
 <213> Homo sapien

<400> 153
 caagattcca taggctgacc aggaggtat tcaagatctc tggcagttga ggaagtctct 60
 ttaagaaaat agtttaaaca atttgtaaaa atttttctgt cttacttcat ttctgtagca 120
 gttgatattc ggctgtcctt ttataaatgc agagtgggaa ctttccctac catgtttgat 180
 aaatgttgct caggctccat tgccaataat gtgtgttcca aaatgcctgt ttagttttta 240
 aagacggaac tccacccttt gcttgggtctt aagtatgtat ggaatgttat gataggacat 300
 agtagtagcg gtggtcagcc tatggaatct tg 332

<210> 154
 <211> 345
 <212> DNA
 <213> Homo sapien

<220>
 <221> misc_feature
 <222> (1)...(345)
 <223> n = A,T,C or G

<400> 154
 tcaagattcc ataggctgac ctggacagag atctcctggg tctggcccag gacagcaggc 60
 tcaagctcag tggagaaggt ttccatgacc ctccagattcc cccaaacctt ggattgggtg 120
 acattgcac tcctcagaga gggaggagat gtangtctgg gcttccacag ggacctggta 180
 ttttaggatc agggtaaccgc tggcctgagg cttggatcat tcanagcctg ggggtggaat 240
 ggctggcagc ctgtggcccc attgaaatag gctctggggc actccctctg ttcctanttg 300
 aacttgggta aggaacagga atgtggtcan cctatggaat cttga 345

<210> 155
 <211> 295
 <212> DNA
 <213> Homo sapien

<220>
 <221> misc_feature
 <222> (1)...(295)
 <223> n = A,T,C or G

<400> 155
 gacgcttggc cacttgacac attaaacagt ttgacataat cactancatg tatttctagt 60
 ttgctgtctg ctgtgatgcc ctgcctgat tctctggcgt taatgatggc aagcataatc 120
 aaacgctgtt ctgttaattc caagttataa ctggcattga ttaaagcatt atctttcaca 180
 actaaactgt tcttcatana acagcccata ttattatcaa attaagagac aatgtattcc 240
 aatatccttt anggccaata tatttnatgt cccttaatta agagctactg tccgt 295

<210> 156
 <211> 406
 <212> DNA
 <213> Homo sapien

<220>
 <221> misc_feature
 <222> (1)...(406)
 <223> n = A,T,C or G

<400> 156
 gacgcttggc cacttgacac tgcagtggga aaaccagcat gagccgctgc cccaaggaa 60
 cctcgaagcc caggcagagg accagccatc ccagcctgca ggtaaagtgt gtcacctgtc 120
 aggtgggctt ggggtgagtg ggtgggggaa gtgtgtgtgc aaagggggtg tnaatgtnta 180
 tgcgtgtgag catgagtgat ggctagtgtg actgcatgtc agggagtgtg aacaagcgtg 240
 cgggggtgtg tgtgcaagtg cgtatgcata tgagaatatg tgtctgtgga tgagtgcatt 300
 tgaaaagtctg tgtgtgtgcg tgtggtcatg anggtaantt antgactgcg caggatgtgt 360
 gagtgtgcat ggaacactca ntgtgtgtgt caagtggccn ancgtc 406

<210> 157
 <211> 208
 <212> DNA
 <213> Homo sapien

<220>
 <221> misc_feature
 <222> (1)...(208)
 <223> n = A,T,C or G

<400> 157
 tgacgcttgg ccacttgaca cactaaaggg tgttactcat cactttcttc tctcctcggg 60
 ggcattgtgag tgcattctatt cacttggcac tcatttgttt ggcagtgaact gtaanccana 120
 tctgatgcat acaccagctt gtaaattgaa taaatgtctc taatactatg tgctcacaat 180
 anggtanggg tgaggagaag gggagaga 208

<210> 158
 <211> 547
 <212> DNA
 <213> Homo sapien

<220>

<221> misc_feature
 <222> (1)...(547)
 <223> n = A,T,C or G

<400> 158
 cttcaacctc cttcaacctc cttcaacctc ctggattcaa acaatcatcc cacctcagac 60
 tccttagtag ctgagactac agactcacgc cactacatct ggctaaatit ttgtagagat 120
 agggtttcat catgttgccc tggctggtct caaactcctg acctcaagca atgtgcccac 180
 ctacgcctcc caaagtgtct ggattacagg cataagccac catgcccagt ccatntttaa 240
 tctttcctac cacattctta ccacactttc ttttatgttt agatacataa atgcttacca 300
 ttatgataca attgccaca gtattaagac agtaacatgc tgcacagggt tgtagcctag 360
 gaacagtagg caataccaca tagcttaggt gtgtggtaga ctataaccatc taggtttgtg 420
 taagttacac tttatgctgt ttacacaatg acaaaacccat ctaatgatgc atttctcaga 480
 atgtatcctt gtcagtaagc tatgatgtac aggaacact gcccaaggac acagatatgt 540
 tacctgt 547

<210> 159
 <211> 203
 <212> DNA
 <213> Homo sapien

<400> 159
 gtcctctctg ccttaccac tcacccagta tgtcagcaat tttatcrgct ttacctacga 60
 aacagcctgt atccaaacac ttaacacact cacctgaaaa gttcaggcaa caatcgctt 120
 ctcatgggtc tctctgctcc agttctgaac ctttctcttt tcctagaaca tgcatttarg 180
 tcgatagaag ttcctctcag tgc 203

<210> 160
 <211> 402
 <212> DNA
 <213> Homo sapien

<400> 160
 tgtaagtga gcagtgtgat ggggtgaaca ggggtgtaag cagtaattgc aaactgtatt 60
 taaacaataa taataatatt tagcatttat agagcacttt atatcttcaa agtacttgca 120
 aacattayct aattaaatac cctctctgat tataatctgg atacaaatgc acttaaaactc 180
 aggacagggc catgagaraa gtatgcattt gaaagtgtgt gctagctatg ctttaaaaaac 240
 ctatacaatg atgggraagt tagagttcag attctgttgg actgtttttg tgcatttcag 300
 ttcagcctga tggcagaatt agatcatatc tgcactcgat gactytgctt gataacttat 360
 cactgaaatc tgagtgttga tcatcacact gctcgactta ca 402

<210> 161
 <211> 193
 <212> DNA
 <213> Homo sapien

<400> 161
 agcatgttga gccagacac tgaccaggag aaaaaccaac caatagaaac acgcccagac 60
 actgaccagg agaaaaacca accaataaaa acaggcccgg acataagaca aataataaaa 120
 ttagcggaca aggacatgaa aacagctatt gtaagagcgg atatagtggg gtgtgtctg 180
 gctcaacatg cta 193

<210> 162
 <211> 147
 <212> DNA
 <213> Homo sapien

<400> 162

45

tggtgagccc agacactgac caggagaaaa accaaccaat aaaaacaggc ccggacataa	60
gacaaataat aaaattagcg gacaaggaca tgaaaacagc tattgtaaga gcggatatag	120
tggtgtgtgt ctgggctcaa catgcta	147

<210> 163
 <211> 294
 <212> DNA
 <213> Homo sapien

<400> 163	
tagcatgttg agcccagaca caaatctttc cttaagcaat aaatcatttc tgcatatggt	60
tttaaaacca cagctaagcc atgattattc aaaaggacta ttgtattggg tattttgatt	120
tgggttctta tctccctcac attatcttca tttctatcat tgacctctta tcccagagac	180
tctcaaactt ttatgttata caaatcacat tctgtctcaa aaaatatctc acccacttct	240
cttctgtttc tgcgtgtgta tgtgtgtgtg tgtgtgtctg ggctcaacat gcta	294

<210> 164
 <211> 412
 <212> DNA
 <213> Homo sapien

<220>
 <221> misc_feature
 <222> (1)...(412)
 <223> n = A,T,C or G

<400> 164	
cgggattggc tttgagctgc agatgctgcc tgtgaccgca cccggcgtgg aacagaaagc	60
cacctggctg caagtgcgcc agagccgccc tgactacgtg ctgctgtggg gctggggcgt	120
gatgaactcc accgccctga aggaagccca ggccaccgga tcccccgcg acaagatgta	180
cggcgtgtgg tgggccggtg cggagcccga tgtgctgac gtgggcgaag gcgccaaggg	240
ctacaacgcg ctggctctga acggctacgg cacgcagtcc aaggatgatcc angacatcct	300
gaaacacgtg cacgacaagg gccagggcac ggggcccaca gacgaagtgg gctcgggtgct	360
gtacaccgcg ggcgtgatca tccagatgct ggacaagggt tcaatcacta at	412

<210> 165
 <211> 361
 <212> DNA
 <213> Homo sapien

<400> 165	
ttgacacctt gtccagcatc tgcattctgat gagagcctca gatggctacc actaatggca	60
gaaggcaaaag gagaacaggc attgtatggc aagaaaaggaa gaaagagaga ggggagaaaag	120
gtgctagggt cttttcaaca accagttctt gatggaactg agagtaagag ctcaaggcca	180
ggtgtggtga ctccaaccag taatcccaac attttaggag gctgaggcag gcagatgtct	240
tgaccccatg agtttgtgac cagcctgaac aacatcatga gactccatct ctacaataat	300
tacaaaaatt aatcaggcat tgtggtatgc cctgtagtcc cagatgctgg acaagggtgc	360
a	361

<210> 166
 <211> 427
 <212> DNA
 <213> Homo sapien

<400> 166	
twgactgact catgtcccct acaccaact atcttctcca ggtggccagg catgatagaa	60
tctgatcctg acttagggga atattttctt tttacttccc atcttgattc cctgccgggtg	120
agtttctctg ttcagggtaa gaaaggagct caggccaaag taatgaacaa atccatcctc	180

acagacgtac	agaataagag	aacwtggacw	tagccagcag	aacmcaaktg	aaamcagaac	240
mcttamctag	gatracaamc	mccrraratar	ktgcycmcmc	wtataataga	aaccaaactt	300
gtatctaatt	aaatatattat	ccacygtcag	ggcattagt	gttttgataa	atacgctttg	360
gctaggattc	ctgaggttag	aatggaaraa	caattgcamc	gagggtaggg	gacatgagtc	420
aktctaa						427

<210> 167

<211> 500

<212> DNA

<213> Homo sapien

<220>

<221> misc_feature

<222> (1)...(500)

<223> n = A,T,C or G

<400> 167

aacgtcgcat	gctcccgcc	gccatggccg	cgggatatagac	tgactcatgt	cccctaagat	60
agaggagaca	cctgctaggt	gtaaggagaa	gatggttagg	tctacggagg	ctccagggtg	120
ggagtagttc	cctgctaagg	gagggtagac	tgttcaacct	gttctctgctc	cggcctccac	180
tatagcagat	gcgagcagga	gtaggagaga	gggaggtaag	agtcagaagc	ttatgttgtt	240
tatgcgggga	aacgccttat	cgggggcagc	cragttatta	ggggacantr	tagwyartcw	300
agntagcatc	caaagcgnng	gagttntccc	atatggttgg	acctgcaggc	ggccgcatta	360
gtgattagca	tgtgagcccc	agacacgcat	agcaacaagg	acctaaactc	agatcctgtg	420
ctgattactt	aacatgaatt	attgtattta	tttaacaact	ttgagttatg	aggcatatta	480
ttaggtccat	attacctgga					500

<210> 168

<211> 358

<212> DNA

<213> Homo sapien

<400> 168

ttcatcgctc	ggtgactcaa	gcctgtaatc	ccagaacttt	gggaggccga	ggggagcaga	60
tcacctgagg	ttgggagttt	gagaccagcc	tgccaacat	ggtgacaacc	cgtctctgct	120
aaaaatacaa	aaattagcca	agcatggtgg	catgcacttg	taatcccagc	tactcgggag	180
gctgaggcag	gagaatcact	tgaggccagg	aggcagaggt	tgcatgagg	cagagggtga	240
gatcatgcca	ctgcactcca	gcctgggcaa	cagagtaaga	ctccatctca	aaaaaaaaaa	300
aaaaaaaaagaa	tgatcagagc	cacaaataca	gaaaaccttg	agtcaccgag	cgatgaaa	358

<210> 169

<211> 1265

<212> DNA

<213> Homo sapien

<400> 169

ttctgtccac	accaatctta	gagctctgaa	agaatttgtc	tttaaataac	ttttaatagt	60
aacatgtatt	ttatggacca	aattgacatt	ttcgactatt	ttttcccaaa	aaaagtcagg	120
tgaatttcag	cacactgagt	tggaatttc	ttatcccaga	agwccgcacg	agcaatttca	180
tattttattta	agattgattc	catactccgt	tttcaaggag	aatccctgca	gtctccttaa	240
aggtagaaca	aatactttct	attttttttt	caccattgtg	ggattggact	ttaagagggtg	300
actctaaaaa	aacagagaac	aaatatgtct	cagttgtatt	aagcacggac	ccatattatc	360
atattcactt	aaaaaaatga	tttctgtg	accttttggc	aacttctctt	ttcaatgtag	420
ggaaaaactt	agtcaccctg	aaaaccaca	aaataaataa	aacttgtaga	tgtgggcaga	480
argtttgggg	gtggacattg	tatgtgttta	aattaaaccc	tgtatcactg	agaagctgtt	540
gtatgggtca	gagaaaatga	atgcttagaa	gctgttcaca	tcttcaagag	cagaagcaaa	600
ccacatgtct	cagctatatt	attatttatt	ttttatgcat	aaagtgaatc	atttcttctg	660
tattaatttc	caaagggttt	taccctctat	ttaaatgctt	tgaaaaacag	tgcattgaca	720

```

atggggttgat atttttcttt aaaagaaaaa tataattatg aaagccaaga taatctgaag      780
cctgttttat tttaaaactt tttatgttct gtgggttgatg ttgtttgttt gtttgtttct      840
attttggttg ttttttactt tgttttttgt tttgttttgt tttggtttdg catactacat      900
gcagtttctt taaccaatgt ctgtttggct aatgtaatta aagttgttaa tttatatgag      960
tgcatttcaa ctatgtcaat ggtttcttaa tatttattgt gtagaagtac tggtaatttt     1020
tttatttaca atatgtttaa agagataaca gtttgatatg ttttcatgtg tttatagcag     1080
aagttattta tttctatggc attccagcgg atatttttgt gtttgcgagg catgcagtca     1140
atattttgta cagtttagtg acagtattca gcaacgcctg atagcttctt tggccttatg     1200
ttaaataaaa agacctgttt gggatgtaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa     1260
aaaaa                                           1265

```

<210> 170

<211> 383

<212> DNA

<213> Homo sapien

<400> 170

```

tgtaagtcga gcagtgatg gacgatattc ttcttattaa tgtggttaatt gaacaaatga      60
tctgtgatac tgatcctgag ctaggaggcg ctgttcagtt aatgggactt cttcgtactc     120
taattgatcc agagaacatg ctggctacaa ctaataaaaac cgaaaaaagt gaatttctaa     180
attttttcta caaccattgt atgcatgttc tcacagcacc acttttgacc aatacttcag     240
aagacaaaatg tgaaaaaggat aatatagttg gatcaaacaa aaacaacaca atttgcctcg     300
ataattatca aacagcacag ctacttgcct taattttaga gttactcaca ttttgtgtgg     360
aacatcacac tgctcgactt aca                                           383

```

<210> 171

<211> 383

<212> DNA

<213> Homo sapien

<400> 171

```

tgggcacctt caatatcgca agttaaaaat aatgttgagt ttattatact tttgacctgt      60
ttagctcaac aggggtgaagg catgtaaaga atgtggactt ctgaggaatt ttctttttaa     120
aagaacataa tgaagtaaca ttttaattac tcaaggacta cttttggttg aagtttataa     180
tctagatacc tctacttttt gtttttgctg ttcgacagtt cacaaagacc ttcagcaatt     240
tacagggtaa aatcgttgaa gtagtggagg tgaaactgaa atttaaaatt attctgtaaa     300
tactataggg aaagaggctg agcttagaat cttttggttg ttcagtgtgt ctgtgctctt     360
atcatcacac tgctcgactt aca                                           383

```

<210> 172

<211> 699

<212> DNA

<213> Homo sapien

<220>

<221> misc_feature

<222> (1)...(699)

<223> n = A,T,C or G

<400> 172

```

tcgggtgatg cctcctcagg cttgtcgtta gtgtacacag agctgctcat gaagcgacag      60
cggtgcccc tggaacttca gaacctcttc ctctacactt ttggtgcgct tctgaatcta     120
ggtctgcatg ctggcgggcg ctctggccca ggctcctggt aaagtttctc aggatgggca     180
gcaactcgtg tgctgagcca ggcactaaat ggactgctca tgtctgctgt catggagcat     240
ggcagcagca tcacacgcct ctttgtggtg tcctgctcgc tgggtgtcaa cgccgtgctc     300
tcagcagtcg tgctacggct gcagctcaca gccgccttct tcctggccac attgctcatt     360
ggcctggcca tgcgcctgta ctatggcagc cgctagtccc tgacaacttc caccctgatt     420
ccggaccctg tagattgggc gccaccacca gatccccctc ccaggccttc ctccctctcc     480

```

catcagcggc	cctgtaacaa	gtgccttggtg	agaaaagctg	gagaagtgag	ggcagccagg	540
ttattctctg	gaggttggtg	gatgaagggg	tacccttagg	agatgtgaag	tgtgggtttg	600
gttaaggaaa	tgcttaccat	ccccacccc	caaccaagtt	nttcagact	aaagaattaa	660
ggtaacatca	atacctaggc	ctgaggaggc	atcacccga			699

<210> 173
 <211> 701
 <212> DNA
 <213> Homo sapien

<400> 173						
tcgggtgatg	cctcctcagg	ccagatcaaa	cttgggggttg	aaaactgtgc	aaagaaatca	60
atgtcggaga	aagaattttg	caaaaagaaa	atgcctaatac	agtactaatt	taataggcca	120
cattagcagt	ggaagaagaa	atgttgatat	tttatgtcag	ctattttata	atcaccagag	180
tgcttagctt	catgtaagcc	atctcgtatt	cattagaaat	aagaacaatt	ttattcgtcg	240
gaaagaactt	ttcaattttat	agcatcttaa	ttgctcagga	ttttaaattt	tgataaagaa	300
agctccactt	ttggcaggag	tagggggcag	ggagagagga	ggctccatcc	acaaggacag	360
agacaccagg	gccagtaggg	tagctggtgg	ctggatcagt	cacaacggac	tgacttatgc	420
catgagaaga	aacaacctcc	aaatctcagt	tgcttaatac	aacacaagct	catttccttc	480
tcacgttaca	tgctctatgt	agatcaacag	caggtgactc	agggaccacg	gctccatctc	540
catatgagct	tccatagcca	ccaggacacg	ggctctgaaa	gtgtcctcca	tgacgggaca	600
catgcctctt	cctttcattg	ggcagagcaa	gtcacttatg	gccagaagtc	acactgcagg	660
gcagtgccat	cctgctgtat	gcctgaggag	gcacacccga	a		701

<210> 174
 <211> 700
 <212> DNA
 <213> Homo sapien

<220>
 <221> misc_feature
 <222> (1)...(700)
 <223> n = A,T,C or G

<400> 174						
tcgggtgatg	cctcctcang	cccctaaatc	agagtccagg	gtcagagcca	caggagacag	60
ggaaagacat	agatttttaac	cggcccctt	caggagattc	tgaggctcag	ttcactttgt	120
tgagtttga	acagaggcag	caaggctagt	ggttaggggc	acggtctcta	aagctgcact	180
gcctggatct	gcctcccagc	tctgccagga	accagctgcg	tgcccttgag	ctgctgacac	240
gcagaaagcc	ccctgtggac	ccagtctcct	cgtctgtaag	atgaggacag	gactctagga	300
accctttccc	ttggtttggc	ctcactttca	caggctccca	tcttgaactc	tatctactct	360
tttcttgaaa	ccttgtaaaa	gaaaaaagt	ctagcctggg	caacatggca	aaaccctgtc	420
tctacaaaaa	atacaaaaat	tagttgggtg	tggtggcatg	tgctgtagt	cccagccact	480
tggtgaggtg	tgaggtggga	ggatcacttg	agcccgggag	gtggaggttg	cagtgcagcca	540
agatcatgcc	actgcactcc	agcctgagta	atagagtaag	actctgtctc	aaaaacaaca	600
acaacaacag	tgagtgtgcc	tctgtttccg	ggttggtagg	ggcaccacat	ttatgcatct	660
ctcagatttg	gacgctgcag	cctgaggagg	catcacccga			700

<210> 175
 <211> 484
 <212> DNA
 <213> Homo sapien

<220>
 <221> misc_feature
 <222> (1)...(484)
 <223> n = A,T,C or G

<400> 175

tatagggcga	attgggcccg	agttgcatgn	tcccggccgc	catggccgcg	ggattcgggt	60
gatgcctcct	caggcttgct	tgccacaagc	tacttctctg	agctcagaaa	gtgccccctg	120
atgagggaaa	atgtcctact	gcactgcgaa	tttctcagtt	ccattttacc	tcccagtcct	180
ccttctaatac	cagttaataa	attcattcca	caagtattta	ctgattacct	gcttgtgcca	240
gggactattc	tcaggctgaa	gaagggtgga	ggggaggggc	gaacctgagg	agccacctga	300
gccagcttta	tatttcaacc	atggctggcc	catctgagag	catctcccca	ctctcgccaa	360
cctatcgggg	catagcccag	ggatgcccc	aggcggccca	ggttagatgc	gtcccctttg	420
cttgtcagtg	atgacataca	ccttagctgc	ttagctggtg	ctggcctgag	gaggcatcac	480
ccga						484

<210> 176

<211> 432

<212> DNA

<213> Homo sapien

<400> 176

tcgggtgatg	cctcctcagg	gctcaaggga	tgagaagtga	cttctttctg	gagggaccgt	60
tcatgccacc	caggatgaaa	atggataggg	accacttg	aggacttgct	gatatgtttg	120
gacaaatgcc	aggtagcgga	attggtactg	gtccaggagt	tatccaggat	agattttcac	180
ccaccatggg	acgtcatcgt	tcaaatcaac	tcttcaatgg	ccatggggga	cacatcatgc	240
ctcccacaca	atcgagttt	ggagagatgg	gaggcaagtt	tatgaaaagc	caggggctaa	300
gccagctcta	ccataaccag	agtcagggac	tcttatccca	gctgcaagga	cagtcgaagg	360
atatgccacc	tcggttttct	aagaaaggac	agcttaatgc	agatgagatt	agcctgagga	420
ggcatcaccc	ga					432

<210> 177

<211> 788

<212> DNA

<213> Homo sapien

<400> 177

tagcatgttg	agcccagaca	cagtagcatt	tgtgccaat	tctggttga	atggtgacaa	60
catgctggag	ccaagtgcta	acatgccttg	gttcaaggga	tggaaagtca	ccogtaaggga	120
tggcaatgcc	agtgaacca	cgctgcttga	ggctetggac	tgcacccctac	caccaactcg	180
cccaactgac	aagcccttgc	gcctgcctct	ccaggatgtc	tacaaaattg	gtgggtattgg	240
tactgttcct	gttgcccgag	tggagactgg	tgttctcaaa	cccggtatgg	tggtcacctt	300
tgtctcagtc	aacgttacaa	cggaaagtaa	atctgtcgaa	atgcaccatg	aagctttgag	360
tgaagctctt	cctggggaca	atgtgggctt	caatgtcaag	aatgtgtctg	tcaaggatgt	420
tcgtcgtggc	aacgttgctg	gtgacagcaa	aaatgaccca	ccaatggaag	cagctggctt	480
cactgctcag	gtgattatcc	tgaaccatcc	aggccaaata	agtgccggct	atgccccctgt	540
attggattgc	cacacggctc	acattgcatg	caagtttgct	gagctgaagg	aaaagattga	600
tcgccgttct	ggtaaaaagc	tggaaagatgg	ccctaaattc	ttgaagtctg	gtgatgctgc	660
cattgttgat	atggttcctg	gcaagcccat	gtgtgttgag	agcttctcag	actatccacc	720
tttgggtcgc	tttgcgtgtc	gtgatatgag	acagacagtt	gcggtgggtg	tctgggctca	780
acatgcta						788

<210> 178

<211> 786

<212> DNA

<213> Homo sapien

<400> 178

tagcatgttg	agcccagaca	cctgtgtttc	tgggagctct	ggcagtggtg	gattcatagg	60
cacttggtgt	gcactttgaa	tgacacactt	ggctttatta	gattcactag	tttttaaaaa	120
attgtgtgtc	gtttcttttc	attaaagggt	taatcagaca	gatcagacag	cataattttg	180
tatttaataga	cagaaacgtt	ggtacatttc	ttcatgaatg	agcttgcatc	ctgaagcaag	240
agcctacaaa	aggcacttgt	tataaatgaa	agttctggct	ctagaggcca	gtactctgga	300

gtttcagagc	agccagtgat	tggtccagtc	agtgatgcct	agttatatag	aggaggagta	360
cactgtgcac	tcttctaggt	gtaagggat	gcaacttttg	atcttaaaat	tctgtacaca	420
tacacacttt	atatatatgt	atgtatgtat	gaaaacatga	aattagtttg	tcaaatatgt	480
gtgtgtttag	tatttttagct	tagtgcaact	atttcacat	tatttattaa	attgatctaa	540
gacactttct	tggtgacacc	ttgaatatta	atgttcaagg	gtgcaatgtg	tattccttta	600
gattgttaaa	gcttaattac	tatgatttgt	agtaaattaa	cttttaaaat	gtatttgagc	660
ccttctgtag	tgtcgtagg	ctcttacagg	gtgggaaaga	ttttaatttt	ccagttgcta	720
attgaacagt	atggcctcat	tatatatttt	gatttatagg	agtttgtgtc	tgggctcaac	780
atgcta						786

<210> 179

<211> 796

<212> DNA

<213> Homo sapien

<400> 179

tagcatgttg	agcccagaca	ctgggtacaa	gaccagacct	gcttcctcca	tatgtaaaca	60
gcttttaaaa	agccagtgaa	cctttttaat	actttggcaa	ccttctttca	caggcaaaga	120
acacccccat	ccgccccttg	tttggagtgc	agagtttggc	tttggttcct	tgccctgcct	180
ggagtatact	tctaattcct	gttgtcctgc	acaagctgaa	taccgagcta	cccaccgcca	240
cccaggccag	gtttccactc	atttattact	ttatgtttct	gttccattgc	tggtccacag	300
aaataagttt	tcctttggag	gaatgtgatt	ataccctttt	aatttcctcc	ttttgctttt	360
ttttaatatc	attggtatgt	gtttggccca	gaggaaactg	aaattcacca	tcattctgac	420
tggcaatccc	attaccatgc	tttttttaaa	aaacgtaatt	tttcttgcc	tacattggca	480
gagtagccct	tcctggctac	tggtttaatg	tagtcaactca	gtttctaggt	ggcattaggc	540
atgagacctg	aagcacagac	tgtcttacca	caaagggtga	caagatctca	aaccttagcc	600
aaagggtat	gtcagggttc	aatgctatct	gcttctgttc	ctgctcactg	ttctggattt	660
tgtccttctt	catccctagc	accagaattt	cccagtctcc	ctccctacct	tcccttgttt	720
taattcta	ctatcagcaa	aataactttt	caaatgtttt	aaccggtatc	tccatgtgtc	780
tgggctcaac	atgcta					796

<210> 180

<211> 488

<212> DNA

<213> Homo sapien

<400> 180

ggatgtgctg	caaggcgatt	aagttgggta	acgccagggt	tttcccagtc	acgacgttgt	60
aaaacgacgg	ccagtggaatt	gtaatacgac	tcactatagg	gcgaattggg	ccgacgtcg	120
catgctcccg	gccgccatgg	ccgcgggata	gcatgttgag	cccagacacc	tgacagtcac	180
ttggagagat	ttttcacgtt	accagcttga	tggtcttttt	caggaggaga	gacactgagc	240
actcccaagg	tgaggttgaa	gatttcctct	agatagccgg	ataagaagac	taggagggat	300
gcctagaaaa	tgattagcat	gcaaatttct	acctgccatt	tcagaactgt	gtgtcagccc	360
acattcagct	gcttcttggt	aactgaaaag	agagagggtat	tgagactttt	ctgatggccg	420
ctctaacatt	gtaacacagt	aatctgtgtg	tgtgtgggtg	tgtgtgtgtg	tctgggctca	480
acatgcta						488

<210> 181

<211> 317

<212> DNA

<213> Homo sapien

<400> 181

tagcatgttg	agcccagaca	cggcgacgg	acctgatgag	tgggggtgatg	gcacctgtga	60
aaaggaggaa	cgatcatccc	catgatattg	gggacccaga	tgatgaacca	tggctcccg	120
tcaatgcata	tttaatccat	gatactgctg	attggaagga	cctgaacctg	aagtttgtgc	180
tgacagttta	tcgggactat	tacctcacgg	gtgatcaaaa	cttcctgaag	gacatgtggc	240
ctgtgtgtct	agtaagggat	gcacatgcag	tggccagtgt	gccaggggta	tggttggtgt	300

ctgggctcaa catgcta

317

<210> 182
 <211> 507
 <212> DNA
 <213> Homo sapien

 <220>
 <221> misc_feature
 <222> (1)...(507)
 <223> n = A,T,C or G

<400> 182
 tagcatgttg agcccagaca ctggctgtta gccaaatcct ctctcagctg ctccctgtgg 60
 tttggtgact caggattaca gaggcacccct gtttcaggga acaaaaagat tttagctgcc 120
 agcagagagc accacataca ttagaatggt aaggactgcc acctccttca agaacaggag 180
 tgagggtggt ggtgaatggg aatggaagcc tgcattccct gatgcatttg tgctctctca 240
 aatcctgtct tagtcttagg aaaggaagta aagtttcaag gacggttccg aactgctttt 300
 tgtgtctggg ctcaacatgc tatcccgcg ccatggcggc cgggagcatg cgacgtcggg 360
 cccaattcgc cctatagtga gtcgtattac aattcactgg ccgtcgtttt acaacgctgt 420
 gactgggaaa accctggcgt tacccaactt aatcgccctg cagcacatcc ccctttccca 480
 gctggcgtaa tancgaaaag gcccgca 507

<210> 183
 <211> 227
 <212> DNA
 <213> Homo sapien

<400> 183
 gattttacgct gcaacactgt ggaggtagcc ctggagcaag gcaggcatgg atgcttctgc 60
 aatccccaaa tggagcctgg tatttcagcc aggaatctga gcagagcccc ctctaattgt 120
 agcaatgata agttattctc tttgttcttc aaccttccaa tagccttgag cttccagggg 180
 agtgtcgtta atcattacag cctgggtctcc acagtgttgc agcgtaa 227

<210> 184
 <211> 225
 <212> DNA
 <213> Homo sapien

<400> 184
 ttacgtcgca aactgtgga gcagattaac atcagacttt tctatcaaca tgactggggg 60
 tactaaaaag acaacaaatc aatggcttca aaagtctaag gaataatttc gatacttcaa 120
 ctttataaaa cctgacaaaa ctatcaatca agcataaaga cagatgaaga acatttccag 180
 attttggcca atcagatatt ttacctccac agtgttgcag cgtaa 225

<210> 185
 <211> 597
 <212> DNA
 <213> Homo sapien

<400> 185
 ggcccgcagt cgcgtgtcc cggccgcat ggccgcggga ttcgttaggg tctctatcca 60
 ctgggaccca taggctagtc agagtattta gagttgagtt cttttctgct tcccagaatt 120
 tgaaagaaaa ggagttaggt gatagagctg agagatcaga tttgcctctg aagcctgttc 180
 aagatgtatg tgctcagacc ccaccactgg ggcctgtggg tgaggctcctg ggcacttatt 240
 tgaatgaatt gctgaagggg agcactatgc caagggaagg gaacccatcc tggcactggc 300
 acaggggtca ccttatccag tgctcagtgc ttctttgctg ctacctggtt ttctctcata 360
 tgtgaggggc aggtagaag aagtgcctrg tgttgtgcga gttttagaac atctaccagt 420

aagtggggaa	gtttcacaaa	gcagcagctt	tgttttgtgt	attttcacct	tcagttagaa	480
gaggaaggct	gtgagatgaa	tgttagttag	gtggaaaaga	cgggtaagct	tagtgatag	540
agaccctaac	gaatcactag	tgcgggccgc	ttgcaggctg	accatatggg	agagctc	597

<210> 186

<211> 597

<212> DNA

<213> Homo sapien

<400> 186

ggcccgaagt	tgcatgttcc	cggccgccat	ggccgcgga	ttcgttagg	tctctatcca	60
ctacctaaaa	aatcccaaac	atataactga	actcctcaca	cccaattgga	ccaatccatc	120
acccagaggg	cctacagatc	ctcctttgat	acataagaaa	atttcccaa	actacctaac	180
tatatcattt	tgcaagattt	gttttaccaa	attttgatgg	cctttctgag	cttgctcagt	240
tgaaccacta	ttacgaacga	tcggatatta	actgccctc	accgtccagg	tgtagctggc	300
aacatcaagt	gcagtaaata	ttcattaaat	ttcacctac	taagggtgct	aaacacccta	360
gggtgccatg	tcggtagcag	atcttttgat	ttgtttttat	ttcccataag	ggtcctgttc	420
aaggtcaatc	atacatgtag	tgtgagcagc	tagtcactat	cgcatgactt	ggaggggtat	480
aatagaggcc	tcctttgctg	ttaaagaact	cttgctccag	cctgtcaaag	tgatagaga	540
ccctaacgaa	tcactagtgc	ggccgcctgc	aggtcgacca	tatgggagag	ctcccaa	597

<210> 187

<211> 324

<212> DNA

<213> Homo sapien

<400> 187

tcgttaggg	ctctatccac	ttgcaggtaa	aatccaatcc	tgtgtatata	ttatagtctt	60
ccatatgtag	tggttcaaga	gactgcagtt	ccagaaagac	tagccgagcc	catccatgtc	120
ttccacttaa	ccctgctttg	ggttacacat	cttaactttt	ctgttcaagt	ttctctgtgt	180
agtttatagc	atgagtattg	ggawaatgcc	ctgaaacctg	acatgagatc	tggaacacac	240
aaacttactc	aataagaatt	tctcccatat	ttttatgatg	gaaaaatttc	acatgcacag	300
aggagtggat	agagacccta	acga				324

<210> 188

<211> 178

<212> DNA

<213> Homo sapien

<220>

<221> misc_feature

<222> (1)...(178)

<223> n = A,T,C or G

<400> 188

gcgcggggat	tcgggggtgat	acctcctcat	gccaaaatac	aacgtntaat	ttcacaactt	60
gccttccaat	ttacgcattt	tcaatttgct	ctccccattt	gttgagtcac	aacaaacacc	120
attgcccaga	aacatgtatt	acctaacatg	cacatactct	taaaactact	catccctt	178

<210> 189

<211> 367

<212> DNA

<213> Homo sapien

<400> 189

tgacaccttg	tccagcatct	gacacagtct	tggtcttgg	aaaatattgg	ataaatgaaa	60
atgaatttct	ttagcaagtg	gtataagctg	agaatatacg	tatcacatat	cctcattcta	120
agacacattc	agtgctccctg	aaattagaat	aggacttaca	ataagtgtgt	tcactttctc	180

53

```

aatagctgtt attcaattga tggtaggcct taaaagtcaa agaaatgaga gggcatgtga      240
aaaaaagctc aacatcactg atcattagaa aacttccatt caaaccacca atgagatacc      300
atctcatacc agtcagaatg gctattatta aaaagtcaaa aaataacaga tgctggacaa      360
ggtgtca                                           367

```

```

<210> 190
<211> 369
<212> DNA
<213> Homo sapien

```

```

<220>
<221> misc_feature
<222> (1)...(369)
<223> n = A,T,C or G

```

```

<400> 190
gacaccttgt ccagcatctg acaacgctaa cagcctgagg agatctttat ttatttattt      60
agtttttact ctggctaggc agatggtggc taaaacattc attta'cccat ttattcattt      120
aattgttcct gcaaggccta tggatagagt attgtccagc actgctctgg aagctaggag      180
catggggatg aacaagatag gctacatcct gtccccacag aacttccact ttagtctggg      240
aaacagatga tatatacaaa tatataaatg aattcaggta gttttaagta cgaaaagaat      300
aagaaagcag agtcatgatt tanaatgctg gaaacagggg ctattgcttg agatattgaa      360
ggtgcccaa                                           369

```

```

<210> 191
<211> 369
<212> DNA
<213> Homo sapien

```

```

<400> 191
tgacaccttg tccagcatct gcacagggaa aagaaactat tatcagagtg aacaggcaac      60
ctacagaatg ggagaaaatt ttgcaatct atccatctga caaagggcta atatccagaa      120
tctacaaaga acttatacaa atttacaaga aacaaacaaa caaacaactc ctcaaaaagt      180
gggtgaagga tgtgaacaga cacttctcaa aagaagacat ttatggggcc aacaaacata      240
tgaaaaaaag ctcacatca ctggtcacta gataaatgca aatcaaaacc acaatgagat      300
accatctcat tccagttaga atggcaatca ttaaaaagtc aggaacaac agatgctgga      360
caaggtgtc                                           369

```

```

<210> 192
<211> 449
<212> DNA
<213> Homo sapien

```

```

<400> 192
tgacgcttgg ccacttgaca cttcatcttt gcacagaaaa acttctttac agatttaatt      60
caagactggc ctagtacag tcctccagac attttttcat ttgttccata tacgtggaat      120
tttaaaatca tgtttcatca gtttgaaatg atttgggctg ctaatcaaca caattggatc      180
gactgttcta ctaaacaca ggaaaatgtg tatctggcag cctgtggaga aacactaaac      240
attgatTTTT ctttgccttt tacggacttt gttccageta catgtaatac caagtctctt      300
ttaagaggag aagatgttga tcttcatttg tttctaccag actgccaccc tagtaaatat      360
tctttattta tgctgtgtaa aaattgccat ccaaataaga tgattcatga tactgggtatt      420
cctgctgagt gtcaagtggc caagcgtca                                           449

```

```

<210> 193
<211> 372
<212> DNA
<213> Homo sapien

```



```

<400> 193
tgacgcttgg ccacttgaca ccagggatgt akcagttgaa tataatcctg caattgtaca      60
tattggcaat ttcccatcaa acattctaga aagagacaac caggattgct aggccataaa    120
agctgcaata aataactggt aattgcagta atcatttcag gccaatcaaa tccagtttgg    180
ctcagagggtg ccttttgctg agagaagagg tgagatataa tgtgttttct tgcaacttct    240
tggagaataa actccacaat agtctgagga ctagatacaa acctatttgc cattaaagca    300
ccagagtctg ttaattccag tactgataag tgttgagat tagactccag tgtgtcaagt    360
ggccaagcgt ca                                                           372

```

```

<210> 194
<211> 309
<212> DNA
<213> Homo sapien

<220>
<221> misc_feature
<222> (1)...(309)
<223> n = A,T,C or G

```

```

<400> 194
tgacgcttgg ccacttgaca cttatgtaga atccatcgtg ggctgatgca agccctttat      60
ttaggcttag tgttgtgggc accttcaata tcacactaga gacaaacgcc acaagatctg    120
cagaaacatt cagttctgan cactcgaatg gcaggataac tttttgtgtt gtaatccttc    180
acatatataa aaacaaactc tgcantctca cgttacaaaa aaacgtactg ctgtaaaata    240
ttaagaaggg gtaaaggata ccatctataa caaagtaact tacaactagt gtcaagtggc    300
caagcgtca                                                           309

```

```

<210> 195
<211> 312
<212> DNA
<213> Homo sapien

<220>
<221> misc_feature
<222> (1)...(312)
<223> n = A,T,C or G

```

```

<400> 195
tgacgcttgg ccacttgaca cccaatctcg cacttcatcc tcccagcacc tgatgaagta      60
ggactgcaac tatcccact tcccagatga ggggaccaan gtacacatta ggacccggat    120
gggagcacag atttgtccga tcccagactc caagcactca gcgtcactcc aggacagcgg    180
ctttcagata aggtcacaaa catgaatggc tccgacaacc ggagtcagtc cgtgctgagt    240
taaggcaatg gtgacacgga tgcacgtgtn acctgtaatg gttcatcgta agtgtcaagt    300
ggccaagcgt ca                                                           312

```

```

<210> 196
<211> 288
<212> DNA
<213> Homo sapien

```

```

<400> 196
tgtatcgacg tagtgggtctc ctcagccatg cagaactgtg actcaattaa acctctttcc      60
tttatgaatt acccaatctc gggtagtgtc tttatagtag tgtgagaatg gactaataca    120
agtacatttt acttagtaat aataataaac aaatatatta catttttgtg tatttactac    180
accatatttt ttattgttat tgtagtgtac accttctact tattaaaaga aataggccccg    240
aggcgggcag atcacgaggt caggagatgg agaccactac gtcgatac                 288

```

```

<210> 197

```

<211> 289
 <212> DNA
 <213> Homo sapien

<400> 197
 ttgggcacct tcaatatcat gacaggtgat gtgataacca agaaggctac taagtgatta 60
 atgggtgggt aatgtataca gagtaggtac actggacaga ggggtaattc atagccaagg 120
 caggagaagc agaatggcaa aacatttcat cactactctc aggatagcat gcagtttaaa 180
 acctataagt agtttatatt tggaattttc cacttaatat ttccagactg caggtaacta 240
 aactgtggaa cacaagaaca tagataaggg gagaccacta cgtcgatac 289

<210> 198
 <211> 288
 <212> DNA
 <213> Homo sapien

<400> 198
 gtatcgacgt agtgggtctcc caagcagtggt gaagaaaacg tgaaccaatt aaaatgtatc 60
 agatacccca aagaaaggcg cttgagtaaa gattccaagt gggtcacaat ctccagatctt 120
 aaaattcagg ctgtcaaaga gatttgctat gaggttgctc tcaatgactt caggcacagt 180
 cggcaggaga ttgaagccct ggccattgtc aagatgaagg agctttgtgc catgtatggc 240
 aagaaagacc ccaatgagcg ggactcctgg agaccactac gtcgatac 288

<210> 199
 <211> 1027
 <212> DNA
 <213> Homo sapien

<220>
 <221> misc_feature
 <222> (1)...(1027)
 <223> n = A,T,C or G

<400> 199
 gctttttggg aaaaacncaa ntgggggaaa gggggnttnn tngcaagggg ataaaagggg 60
 aancccaggg tttcccccatt caggggaggtg taaaaaagncg gccaggggat tgtaanagga 120
 ttcaataata gggggaatgg gccnngaagt tgcaagggtc cngcccgcga tgnccgaggg 180
 atttagtgac attacgacgs tggtataaaa gtgggsccaa waaatatttg tgatgtgatt 240
 tttsgaccag tgaaccatt gwacaggacc tcatttctcy tgagatgrta gccataatca 300
 gataaaagrt tagaagtytt tctgcacgtt aacagcatca ttaaattggg tggcatcacc 360
 aatttcaccc tttgttagcc gatacccttc ccttgaaggc attcaattaa gtgaccaatc 420
 gtcatacgag aggggatggc atggggattg atgatgatat caggggtgat accttcacag 480
 gtgaaaggca taccctcttg tctatactga ataccacaag tacccttttg accatgtcga 540
 ctacgaaatt tgtctccaat ctgtgtwatc cctaacagag cgtaccctta ttttacaaaa 600
 tttatatact tctgtattga gagttaccat aacctgatcc acaatgcccg tctcgtwtgt 660
 tctgagaaaa gtgctacagt ctctcttggt atagcgtcta ttgggtgctc ccaattcatc 720
 ttcatTTTTT aggcaagggt aactgttttg cctataataa cmtcatctcc tgatacmcga 780
 aacccckgga rctatcaaac catcatcatc cagcgttckt watgtymcta aatccctatt 840
 gggccgcct gcagggtcaac atatnggaaa acccccacc ccttnggagc ntaccttgaa 900
 tttccatat gtcccnTaaa ttanctngnc ttanctggc cntaacctnt tccggtttaa 960
 attgtttccg ccccncttcc cnccttnna accggaacc ttaattttna accnggggtt 1020
 cctatcc 1027

<210> 200
 <211> 207
 <212> DNA
 <213> Homo sapien

56

```

<400> 200
agtgacatta cgacgctggc catcttgaat cctagggcat gaagttgcc caaagttcag    60
cacttggtta agcctgatcc ctctggttta tcacaaagaa taggatggga taaagaaagt   120
ggacacttaa ataagctata aattatatgg tccttgtcta gcaggagaca actgcacagg   180
tatactacca gcgtcgtaat gtcacta                                     207

```

```

<210> 201
<211> 209
<212> DNA
<213> Homo sapien

```

```

<400> 201
tgggcacctt caatatctat taaaagcaca aatactgaag aacacaccaa gactatcaat    60
gaggttacat ctggagtcct cgatatatca ggaaaaaatg aagtgaacat tcacagagtt   120
ttacttcttt gggaactcaa atgctagaaa agaaaagggt gccctctttc tctggcttcc   180
tggtcctatc cagcgtcgta atgtcacta                                     209

```

```

<210> 202
<211> 349
<212> DNA
<213> Homo sapien

```

```

<220>
<221> misc_feature
<222> (1)...(349)
<223> n = A,T,C or G

```

```

<400> 202
ntacgctgca acactgtgga gccactgggt tttattcccg gcaggttatc cagcaaacag    60
tcactgaaca caccgaagac cgtggtatgg taaccgttca cagtaatcgt tcagtcgctc   120
tgccgggacc cgacgagcgt cactgggtac agaccagatt cagccggaag agaaagcgcc   180
gcaggagag actcgaactc cactccgctg gtgagcagcc ccatgttttc aactcgaagt   240
tcaaacggca ttgggttata taccatcagc tgaacttcac acacatctcc ttgaaccac    300
tggaatcta ttttcttggt ccgctcttct ccacagtgtt gcagcgtaa               349

```

```

<210> 203
<211> 241
<212> DNA
<213> Homo sapien

```

```

<400> 203
tgctctcttt gccttaccaa cccaaagccc actgtgaaat atgaagtga tgacaaaatt    60
cagttttcaa cgcaatatag tatagtttat ctgattcttt tgatctccag gacactttaa   120
acaactgcta ccaccaccac caacctaggg atttaggatt ctccacagac cagaaattat   180
ttctcctttg agtttcaggc tcctctggga ctctgttca tcaatgggtg gtaaatggct   240
a                                                    241

```

```

<210> 204
<211> 248
<212> DNA
<213> Homo sapien

```

```

<400> 204
tagccattta ccacccatct gcaaaccswg acmwwcargr cywgwackya ggcgatttga    60
agtactggta atgctctgat catgttagtt acataagtgt ggtcagttta caaaaattca   120
cagaactaaa tactcaatgc tatgtgttca tgtctgtgtt tatgtgtgtg taatgtttca   180
attaagtttt tttaaaaaaa agagatgatt tccaaataag aaagccgtgt tggttaaggca   240
agaggagc                                     248

```

<210> 205
 <211> 505
 <212> DNA
 <213> Homo sapien

<220>
 <221> misc_feature
 <222> (1)...(505)
 <223> n = A,T,C or G

<400> 205
 tacgctgcaa cactgtggag ccattcatac aggtccctaa ttaaggaaca agtgattatg 60
 ctacctttgc acggttaggg taccgcggcc gttaaacaatg tgctactggg caggcgggtgc 120
 ctctaatact ggtgatgcta gaggtgatgt ttttggtaaa caggcgggggt aagatttgcc 180
 gagttccttt tacttttttt aacctttcct tatgagcatg cctgtgttgg gttgacagtg 240
 ggggtaataa tgacttggtg gttgattgta gatattgggc tgtaattgt cagttcagtg 300
 ttttaatctg acgcaggctt atgcggagga gaatgttttc atgttactta tactaacatt 360
 agttcttcta tagggtgata gattgggtcca attgggtgtg aggagttcag ttatatgttt 420
 gggatttttt aggtagtggg tgttgancct gaacgctttc ttaattggtg gctgctttta 480
 rgcctactat ggggtgtaaa tggct 505

<210> 206
 <211> 179
 <212> DNA
 <213> Homo sapien

<400> 206
 tagactgact catgtcccct accaaagccc atgtaaggag ctgagttctt aaagactgaa 60
 gacagactat tctctggaga aaaataaaat ggaaattgta ctttaaaaaa aaaaaaatc 120
 ggccgggcat ggtagcacac acctgtaatc ccagctacta ggggacatga gtcagtcta 179

<210> 207
 <211> 176
 <212> DNA
 <213> Homo sapien

<400> 207
 agactgactc atgtccccta cccacacctc tgctgtgctg ccgtgttcct aacagggtcac 60
 agactggtac tggtcagtgg cctgggggtt ggggacctct attatatggg atacaaattt 120
 aggagtggga attgacacga tttagtact gatgggatat ggggtgtaaa tggcta 176

<210> 208
 <211> 196
 <212> DNA
 <213> Homo sapien

<400> 208
 agactgactc atgtccccta tttacaggg tctctagtgc tgtgaaaaaa aaaaatgctg 60
 aacattgcat ataacttata ttgtaagaaa tactgtacaa tgactttatt gcatctgggt 120
 agctgtaagg catgaaggat gccagaagt ttaaggaata tgggtggtaa atggctaggg 180
 gacatgagtc agtcta 196

<210> 209
 <211> 345
 <212> DNA
 <213> Homo sapien

<220>
 <221> misc_feature
 <222> (1)...(345)
 <223> n = A,T,C or G

<400> 209
 gacgcttggc cacttgacac cttttatattt ttaaggattc ttaagtcatt tangtnactt 60
 tgtaagtttt tcctgtgccc ccataagaat gatagcttta aaaattatgc tggggtagca 120
 aagaagatac ttctagcttt agaattgtga ggtatagcca ggattcttgt gaggaggggt 180
 gatttagagc aaatttctta ttctccttgc ctcatctgta acatggggat aataatagaa 240
 ctggcctgac aaggttggaa ttagtattac atggtaaata catgtaaaat gtttagaatg 300
 gtgccaagta tctaggaagt acttgggcat gggtggtaaa tggct 345

<210> 210
 <211> 178
 <212> DNA
 <213> Homo sapien

<400> 210
 gacgcttggc cacttgacac tagagtaggg ttggccaac tttttctata aaggaccaga 60
 gagtaaatat ttcaggcttt gtgggttggt cagtctctct tgcaactact cagctctgcc 120
 attgtagcat agaatcagc catagacagg acagaaatga atgggtggta aatggcta 178

<210> 211
 <211> 454
 <212> DNA
 <213> Homo sapien

<400> 211
 tgggcacctt caatatctat ccagcgcac taaattcgct tttttcttga ttaaaaattt 60
 caccacttgc tgtttttgct catgtatacc aagtagcagt ggtgtgaggc catgcttggt 120
 ttttgattcg atatcagcac cgtataagag cagtgccttg gccattaatt tatcttcatt 180
 gtagacagca tagtgtagag tggatctctc atactcatct ggaatatattg gatcagtgcc 240
 atgttcacgc aacattaacg cacattcatc ttcttggcat tgtacggcct ttgtcagagc 300
 tgtcctcttt ttgttgtaaa ggacattaag ttgacatcgt ctgtccagca cgagttttac 360
 tacttctgaa ttcccattgg cagaggccag atgtagagca gtcctctttt gcttgtccct 420
 cttgttcaca tcagtgtccc tgagcataac ggaa 454

<210> 212
 <211> 337
 <212> DNA
 <213> Homo sapien

<400> 212
 tccgttatgc caccagaaa acctactgga gttacttatt aacatcaagg ctggaaccta 60
 tttgcctcag tcctatctga ttcattgagca catggttatt actgatcgca ttgaaaacat 120
 tgatcacctg ggtttcttta ttatcgact gtgtcatgac aaggaaactt acaaactgca 180
 acgcagagaa actattaaaag gtattcagaa acgtgaagcc agcaattgtt tcgcaattcg 240
 gcattttgaa aacaaatttg ccgtggaaac tttaatttgt tcttgaacag tcaagaaaaa 300
 cattattgag gaaaattaat atcacagcat aacggaa 337

<210> 213
 <211> 715
 <212> DNA
 <213> Homo sapien

<220>
 <221> misc_feature

59

<222> (1)...(715)

<223> n = A,T,C or G

<400> 213

tcgggtgatg	cctcctcagg	catcttccat	ccatctcttc	aagattagct	gtcccaaagt	60
tttttccttc	tcttctttac	tgataaat	ggactccttc	ttgacactga	tgacagcttt	120
agtatccttc	ttgtcacctt	gcagacttta	aacataaaaa	tactcattgg	ttttaaaagg	180
aaaaaagtat	acattagcac	tattaagctt	ggccttgaaa	cattttctat	cttttattaa	240
atgtcggtta	gctgaacaga	attcatttta	caatgcagag	tgagaaaaga	aggagagctat	300
atgcatttga	gaatgcaagc	attgtcaaat	aaacatttta	aatgctttct	taaagtgagc	360
acatacagaa	atacattaag	atattagaaa	gtgtttttgc	ttgtgtacta	ctaattaggg	420
aagcaccttg	tatagttcct	cttctaaaa	tgaagtagat	tttaaaaacc	catgtaattt	480
aattgagctc	tcagttcaga	ttttaggaga	attttaacag	ggatttggtt	ttgtctaaat	540
tttgtcaatt	tntttagtta	atctgtataa	ttttataaat	gtcaaacgtg	atttagtccg	600
ttttcatgct	gctatgaaag	aaatacccan	gacagggtta	tttataaang	gaaagangtt	660
aatttgactc	ccagttcaca	ggcctgagga	ngnatcnccc	gaaatcctta	ttgcg	715

<210> 214

<211> 345

<212> DNA

<213> Homo sapien

<220>

<221> misc_feature

<222> (1)...(345)

<223> n = A,T,C or G

<400> 214

ggtaangngc	atacntcggt	gctccggccg	ccggagtcgg	gggattcggg	tgatgcctcc	60
tcaggcccac	ttgggcctgc	ttttcccaaa	tggcagctcc	tctggacatg	ccattccttc	120
tcccacctgc	ctgattcttc	atatgttggg	tgccctgtt	tttctggtgc	tatttcctga	180
ctgctgttca	gctgccactg	tctgcaaag	cctgcctttt	taaatgcctc	accattcctt	240
catttgtttc	ttaaatatgg	gaagtgaag	tgccacctga	ggccgggcac	agtggctcac	300
gcctgtaatc	ccagcacttt	gggagcctga	ggaggcatca	cccga		345

<210> 215

<211> 429

<212> DNA

<213> Homo sapien

<400> 215

ggtgatgcct	cctcaggcga	agctcagggg	ggacagaaac	ctcccgtgga	gcagaagggc	60
aaaagctcgc	ttgatcttga	ttttcagtac	gaatacagac	cgtgaaagcg	gggcctcacg	120
atccttctga	ccttttgggt	tttaagcagg	aggtgtcaga	aaagttacca	cagggataac	180
tggttgttgg	cggccaagcg	ttcatagcga	cgtcgtttt	tgatccttcg	atgtcggctc	240
ttcctatcat	tgtgaagcag	aattcaccaa	gcgttggtt	gttcacccac	taatagggaa	300
cgtgagctgg	gttttagaccg	tcgtgagaca	ggttagtttt	accctactga	tgatgtgtkg	360
ttgccatggg	aatcctgctc	agtacgagag	gaaccgcagg	ttcasacatt	tggtgtatgt	420
gcttgccctt						429

<210> 216

<211> 593

<212> DNA

<213> Homo sapien

<220>

<221> misc_feature

<222> (1)...(593)

<223> n = A,T,C or G

<400> 216

tgacacctat	gtcncgcatc	tgttcacagt	ttccacaaat	agccagcctt	tgccacctc	60
tctgtcctga	ggtatacaag	tatatcagga	ggtgtatacc	ttctcttctc	ttccccacca	120
aagagaacat	gcaggctctg	gaagctgtct	taggagcctt	tggtctcaga	atttcagagt	180
cttgggtacc	ttggtgtgtg	tctggaagga	gaaacatttg	ctctggataa	ggagtacagc	240
cggaggagg	tcacagagcc	ctcagctcaa	gccctgtgc	cttagtctaa	aagcagcttt	300
ggatgaggaa	gcaggttaag	taacatacgt	aagcgtacac	aggtagaaag	tgctgggagt	360
cagaattgca	cagtgtgtag	gagtagtacc	tcaatcaatg	agggcaaadc	aactgaaaga	420
agaagaccna	ttaatgaatt	gcttangggg	aaggatcaag	gctatcatgg	agatctttct	480
aggaagatta	ttgtttanaa	ttatgaaagg	antagggcag	ggacagggcc	agaagtanaa	540
ganaacattg	cctatanccc	ttgtcttgca	cccagatgct	ggacaagggtg	tca	593

<210> 217

<211> 335

<212> DNA

<213> Homo sapien

<400> 217

tgacaccttg	tccagcatct	gacgtgaaga	tgagcagctc	agaggaggtg	tcctggattt	60
cctggttctg	tggtctccgt	ggcaatgaat	tcttctgtga	agtggatgaa	gactacatcc	120
aggacaaatt	taatcttact	ggactcaatg	agcaggtccc	tcactatcga	caagctctag	180
acatgatctt	ggacctggag	cctgatgaag	aactggaaga	caacccaac	cagagtgaac	240
tgattgagca	ggcagccgag	atgctttatg	gattgatcca	cgcccgctac	atccttacca	300
accgtggcat	cgcccagatg	ctggacaagg	tgtea			335

<210> 218

<211> 248

<212> DNA

<213> Homo sapien

<400> 218

tacgtactgg	tcttgaagg	cttaggtaga	gaaaaaatgt	gaatatttaa	tcaaagacta	60
tgtatgaaat	gggactgtaa	gtacagaggg	aagggtggcc	cttatcgcca	gaagttggta	120
gatgcgtccc	cgatcatgaa	tgttgtgtca	ctgcccagaca	tttgccgaat	tactgaaatt	180
ccgtagaatt	agtgcgaatt	ctaacgttgt	tcatctaaga	ttatggttcc	atgtttctag	240
tactttta						248

<210> 219

<211> 530

<212> DNA

<213> Homo sapien

<220>

<221> misc_feature

<222> (1)...(530)

<223> n = A,T,C or G

<400> 219

tgacgcttgg	ccacttgaca	caagtagggg	ataaggacaa	agacccatna	ggtggcctgt	60
cagccttttg	ttactgttgc	ttccctgtca	ccacggcccc	ctctgtaggg	gtgtgctgtg	120
ctctgtggac	attggtgcat	tttcacacat	accattctct	ttctgtttca	cagcagtcct	180
gaggcgggag	cacacaggac	taccttgtca	gatgangata	atgatgtctg	gccaaactcac	240
ccccaaacct	tctcactagt	tatangaaga	gccangccta	naaccttcta	tctgncccc	300
ttgccctatg	acctcatccc	tgttccatgc	cctattctga	tttctggtga	actttggagc	360
agcctggttt	ttcctcctca	ctccagcctc	tctccatacc	atggtanggg	ggtgctgttc	420
cacncaaang	gtcaggtgtg	tctggggaat	cctnananct	gccnggagtt	tccnangcat	480

61

tcttaaaaac cttcttgcc t aatcanatng tgtccagtgg ccaacntcn 530

<210> 220

<211> 531

<212> DNA

<213> Homo sapien

<400> 220

tgacgcttg	ccacttgaca	ctaaatagca	tcttctaaag	gcctgattca	gagttgtgga	60
aaattctccc	agtgtcaggg	attgtcagga	acagggtgc	tcctgtgtc	actttacctg	120
ctgtgtttct	gctggaaaag	gagggaagag	gaatggctga	tttttaccta	atgtctccca	180
gtttttcata	ttcttcttg	atcctcttct	ctgacaactg	ttcccttttg	gtcttcttct	240
tcttgctcag	agagcaggtc	tctttaaaac	tgagaaggga	gaatgagcaa	atgattaaag	300
aaaacacact	tctgaggccc	agagatcaaa	tattaggtaa	atactaaacc	gcttgccctg	360
tgtggctact	tttctcctct	ttcacatgct	ctatccctct	atccccacc	tattcatatg	420
gcttttatct	gccaaagttat	ccggcctctc	atcaaccttc	tcccctagcc	tactggggga	480
tatccatctg	ggtctgtctc	tggtgtattg	gtgtcaagtg	gccaaagcgc	a	531

<210> 221

<211> 530

<212> DNA

<213> Homo sapien

<400> 221

attgacgctt	ggccacttga	cacccgcctg	cctgcaatac	tggggcaagg	gccttcaactg	60
ctttcctgcc	accagctgcc	actgcacaca	gagatcagaa	atgctacca	ccaagactgt	120
tggtcctcag	cctctctgag	gagaaagagc	agaagcctgg	aagtcagaag	agaagctaga	180
tcggctacgg	ccttggcagc	cagcttcccc	acctgtggca	ataaagtcgt	gcatggctta	240
acaatggggg	cacctcctga	gaaacacatt	gttaggcaat	tcggcgtgtg	ttcatcagag	300
catatttaca	caaacctcga	tagtgagccc	tactatccac	tattgtcct	acgctgcaaa	360
cctgaacagc	atgggactgt	actgaatact	ggaagcagct	ggtgatggta	cttatttgtg	420
tatctaaaca	cagagaaggt	acagtaagaa	tatggatatca	taaacttaca	gggaccgcca	480
tcctatatgc	agtctgttgt	gacccaaatg	tgtcaagtgg	ccaagcgtca		530

<210> 222

<211> 578

<212> DNA

<213> Homo sapien

<220>

<221> misc_feature

<222> (1)...(578)

<223> n = A,T,C or G

<400> 222

tgtatcgacg	tagtggctc	cgggctacta	ggccgttgtg	tgtgtgtagt	acctggttca	60
ctgaaaggcg	catctccctc	cccgcgtcgc	cctgaagcag	ggggaggact	tcgcccagcc	120
aaggcagttg	tatgagtttt	agctgcggca	cttcgagacc	tctgagccca	cctccttcag	180
gagccttccc	cgattaagga	agccagggtg	aggattcctt	cctccccag	acaccacgaa	240
caaaccacca	ccccccctat	tctggcagcc	catatacatc	agaacgaaac	aaaaataaca	300
aataaacnaa	aacccaaaaa	aaaagagaag	gggaatgta	tatgtctgtc	catcctgttg	360
ctttagcctg	tcagctccta	nagggcaggg	accgtgtctt	ccgaatggtc	tgtgcagcgc	420
cgaactgagg	aagtatcgga	ggaggaagca	gagtcagcag	aagttgaacg	gtgggcccg	480
cggctcttgg	gggctgtgtg	tgtacttcga	gaccgctttc	gctttttgtc	ttagatttac	540
gtttgctctt	tggagtggga	naccactacn	tcnatata			578

<210> 223

<211> 578

<212> DNA

<213> Homo sapien

<400> 223

tgtatcgacg	tagtgggtctc	ctcttgcaaa	ggactggctg	gtgaatgggt	tcctgaatt	60
atggacttac	cctaaacata	tcttatcatc	attaccagtt	gcaaaatatt	agaatgtgtt	120
gtcactgttt	catttgattc	ctagaagggt	agtccttagat	atgttacttt	aacctgtatg	180
ctgtagtgct	ttgaatgcat	tttttgtttg	catttttggt	tgcccaacct	gtcaattata	240
gctgcttagg	tctggactgt	cctggataaa	gctgttaaaa	tattcaccag	tccagccatc	300
ttacaagcta	attaagtcaa	ctaaatgctt	ccttgttttg	ccagacttgt	tatgtcaatc	360
ctcaatttct	gggttcattt	tgggtgccct	aaatccttag	gtgtgacttt	cttagcatcc	420
tgtaacatcc	attccaagc	aagcacaact	tcacataata	ctttccagaa	gttcattgct	480
gaagcccttc	cttcacccag	cggagcaact	tgattttcta	caacttcct	catcagagcc	540
acaagagtat	gggatatgga	gaccactacg	tcgatata			578

<210> 224

<211> 345

<212> DNA

<213> Homo sapien

<220>

<221> misc_feature

<222> (1)...(345)

<223> n = A,T,C or G

<400> 224

tgtatcgacg	tantgggtctc	ccaagggtgct	gggattgcag	gcatgagcca	ccactcccag	60
gtggatcttt	ttctttatac	ttacttcatt	aggtttctgt	tattcaagaa	gtgtagtggt	120
aaaagtcttt	tcaatctaca	tggttaaata	atgatagcct	gggaaataaa	tagaaatttt	180
ttctttcatc	tttaggttga	ataaagaaac	agaaaaata	gaacatactg	aaaataatct	240
aagttccaac	catagaagaa	ctgcagaaga	aatgaagaaa	gtgatgatga	tttagatttt	300
gatattgatt	tagaagacac	aggaggagac	cactacgtcg	ataca		345

<210> 225

<211> 347

<212> DNA

<213> Homo sapien

<400> 225

tgtatcgacg	tagtgggtctc	caaactgagg	tatgtgtgcc	actagcacac	aaagccttcc	60
aacagggacg	caggcacagg	cagttttaaag	ggaatctgtt	tctaaattaa	tttccacctt	120
ctctaagtat	tctttcctaa	aactgatcaa	ggtgtgaagc	ctgtgctctt	tcccaactcc	180
cctttgacaa	cagccttcaa	ctaacacaag	aaaaggcatg	tctgacactc	ttcctgagtc	240
tgactctgat	acgttgttct	gatgtctaaa	gagctccaga	acaccaaagg	gacaattcag	300
aatgctggtg	tataacagac	tccaatggag	accactacgt	cgatata		347

<210> 226

<211> 281

<212> DNA

<213> Homo sapien

<220>

<221> misc_feature

<222> (1)...(281)

<223> n = A,T,C or G

<400> 226

aggngnggga	ntgtatcgac	gtagtgggtct	cccaacagtc	tgtcattcag	tctgcagggtg	60
------------	------------	-------------	------------	------------	-------------	----

tcagtgtttt	ggacaatgag	gcaccattgt	cacttattga	ctcctcagct	ctaaatgctg	120
aaattaaatc	ttgtcatgac	aagtctggaa	ttcctgatga	ggttttacaa	agtatttttg	180
atcaatactc	caacaaatca	gaaagccaga	aagaggatcc	tttcaatatt	gcagaaccac	240
gagtggattt	acacacctca	ggagaccact	acgtcgatac	a		281

<210> 227
 <211> 3646
 <212> DNA
 <213> Homo sapien

<400> 227

gggaaacact	tcctcccagc	cttgtaaggg	ttggagccct	ctccagtata	tgctgcagaa	60
tttttctctc	ggtttctcag	aggattatgg	agtcgcgcct	aaaaaaggca	agctctggac	120
actctgcaaa	gtagaatggc	caaagtttgg	agttgagtg	ccccttgaag	ggtcactgaa	180
cctcacaatt	gttcaagctg	tgtggcgggt	tgttactgaa	actcccggcc	tccctgatca	240
gtttccctac	attgatcaat	ggctgagttt	ggtcaggagc	acccttccg	tggtccact	300
catgcaccat	tcataatttt	acctccaagg	tcctcctgag	ccagaccgtg	ttttcgccctc	360
gaccctcagc	cggttcggct	cgccctgtac	tgctctctc	tgaagaagag	gagagtctcc	420
ctcaccagct	cccaccgcct	taaaaccagc	ctactccctt	agggtcatcc	catgtctcct	480
cggctatgtc	ccctgtaggc	tcacaccca	ttgcctcttg	gttgcaaccg	tggtgggagg	540
aagtagcccc	tctactacca	ctgagagagg	cacaagtccc	tctgggtgat	gagtgtctca	600
ccccttccct	ggtttatgtc	ccttctttct	acttctgact	tgtataattg	gaaaacccat	660
aatcctccct	tctctgaaaa	gccccaggct	ttgacctcac	tgatggagtc	tgtactctgg	720
acacattggc	ccacctggga	tgactgtcaa	cagtcctttt	tgacctttt	cacctctgaa	780
gagagggaaa	gtatccaaag	agaggccaaa	aagtacaacc	tcacatcaac	caataggccg	840
gaggaggaag	ctagaggaat	agtgattaga	gacccaattg	ggacctaat	gggacccaaa	900
tttctcaagt	ggaggagaaa	cttttgacga	tttccaccgg	tatctcctcg	tggttattca	960
gggagctgct	cagaaacctc	taaacttgct	taaggcgact	gaagtcgtcc	aggggcatga	1020
tgagtcacca	ggagtgtttt	tagagcacct	ccaggaggct	tatcagattt	acaccccttt	1080
tgacctggca	gccccgaaa	atagccatgc	tcttaatttg	gcatttgtgg	ctcaggcagc	1140
cccagatagt	aaaaggaaac	tccaaaaact	agagggattt	tgctggaatg	aataccagtc	1200
agctttttaga	gatagcctaa	aagggttttg	acagtcgaag	ggttgaaaaa	caaaaacaag	1260
cagctcaggc	agctgaaaaa	agccactgat	aaagcatcct	ggagtatcag	agtttactgt	1320
tagatcagcc	tcatttgact	tccctcccca	catggtgttt	aaatccagct	acactacttc	1380
ctgactcaaa	ctccactatt	cctgttcatg	actgtcagga	actgttgga	actactgaaa	1440
ctggccgacc	tgatcttcaa	aatgtgcccc	taggaaagg	ggatgccacc	atgttcacag	1500
acagttagcag	cttctcagag	aagggaactc	gaaaggccgg	tgcaagctgt	accatggaga	1560
cagatgtgtt	gtgggtcag	gctttaccag	caaacacctc	agcacaaaag	gctgaattga	1620
tcgccctcac	tcaggctctc	cgatggggta	aggatattaa	cgtaaacact	gacagcaggt	1680
acgcctttgc	tactgtgcac	gtacgtggag	ccatctacca	ggagcgtggg	ctactcacct	1740
cagcaggtgg	ctgtaatcca	ctgtaaagga	catcaaaagg	aaaacacggc	tgttgcccgt	1800
ggtaaccaga	aagctgattc	agcagctcaa	gatgcagtg	gactttcagt	cacgcctcta	1860
aacttgctgc	ccacagtctc	ctttccacag	ccagatctgc	ctgacaatcc	cgcatactca	1920
acagaagaag	aaaactggcc	tcagaactca	gagccaataa	aaatcaggaa	ggttggtgga	1980
ttcttctctga	ctctagaatc	ttcatacccc	gaactcttgg	gaaaacttta	atcagtcacc	2040
tacagtctac	cacccattta	ggaggagcaa	agctacctca	gctcctccgg	agccgtttta	2100
agatcccca	tcttcaaagc	ctaacagatc	aagcagctct	ccggtgcaca	acctgcgccc	2160
aggtaaatgc	caaaaaaggt	cctaaaccga	gccaggccca	ccgtctccaa	gaaaactcac	2220
caggagaaaa	gtgggaaatt	gactttacag	aagtaaaacc	acaccgggct	gggtacaaat	2280
accttctagt	actggtagac	accttctctg	gatggactga	agcatttget	accaaaaacg	2340
aaactgtcaa	tatggtagtt	aagtttttac	tcaatgaaat	catccctcga	catgggctgc	2400
ctgtttgcca	tagggtctga	taatggaccg	gccttcgcct	tgtctatagt	ttagtcagtc	2460
agtaaggcgt	taaacattca	atggaagctc	catttgtcct	atcgacccca	gagctctggg	2520
caagtagaac	gcatgaactg	caccctaaaa	aacactctta	caaaattaat	cttagaaaacc	2580
ggtgtaaat	gtgtaagtct	ccttccttta	gccctactta	gagtaagggtg	caccccttac	2640
tgggctgggt	tcttaccttt	tgaaatcatg	tatgggaggg	tgctgcctat	cttgccctaag	2700
ctaagagatg	cccaattggc	aaaaatatca	caaactaatt	tattacagta	cctacagctc	2760
ccccaacag	tacaagatat	catcctgcca	cttgtctcag	gaacccatcc	caatccaatt	2820

cctgaacaga	cagggccctg	ccattcattc	ccgccagggt	acctgttggt	tgtaaaaaag	2880
ttccagagag	aaggactccc	tctgtcttgg	aagagacctc	acaccgtcat	cacgatgcca	2940
acggctctga	aggtggatgg	cattcctgcg	tggtatcatc	actcccgcac	caaaaaggcc	3000
aacagagccc	aactagaaac	atgggtcccc	agggctgggt	cagggccctt	aaaactgcac	3060
ctaagttggg	tgaagccatt	agattaattc	tttttcttaa	ttttgtaaaa	caatgcatag	3120
cttctgtcaa	acttatgtat	cttaagactc	aatataaccc	ccttgttata	actgaggaat	3180
caatgatttg	attcccccaa	aaacacaagt	ggggaatgta	gtgtccaacc	tggtttttac	3240
taaccctggt	tttagactct	ccctttcctt	taatcactca	gcttgtttcc	acctgaattg	3300
actctccctt	agctaagagc	gccagatgga	ctccatcttg	gctctttcac	tggcagccgc	3360
ttcctcaagg	acttaacttg	tgcaagctga	ctcccagcac	atccaagaat	gcaattaact	3420
gataagatac	tgtggcaagc	tatatccgca	gttcccagga	attcgtccaa	ttgatcacag	3480
cccctctacc	cttcagcaac	caccaccctg	atcagtcagc	agccatcagc	accgaggcaa	3540
ggccctccac	cagcaaaaag	attctgactc	actgaagact	tggtatgatca	ttagtatttt	3600
tagcagtaaa	gttttttttt	ctttttcttt	ctttttttct	cgtgcc		3646

<210> 228

<211> 419

<212> DNA

<213> Homo sapien

<220>

<221> misc_feature

<222> (1)...(419)

<223> n = A,T,C or G

<400> 228

taagagggtg	caagatctaa	gcacagccgt	caatgcagaa	cacagaacgt	agcctggtaa	60
gtgtgttaag	agtgggaatt	tttgagtag	agagtaaggc	acctaaccct	agctgggggt	120
tggtgacggg	ccagatggc	ttacagaaga	aagtgtcctg	agatgagttt	ttaagaatga	180
ataaggatag	acacaagtga	ggactgactt	ggcagtgggt	aatggtgggt	ggcaaaaaac	240
ttcgcatgta	tggaaactgc	acgtacagga	atgaagaatg	agactgtgtg	gtgtttaatg	300
agctgcaaat	actaatttta	tcctgaaagt	tttgaagagt	taactaaaaa	gtatttttta	360
gtaaggaaat	aaccctacat	ttcagggtta	ttgtttgttt	anatattgaa	ggtgcccaa	419

<210> 229

<211> 148

<212> DNA

<213> Homo sapien

<400> 229

aagagggtac	ctgtatgtag	ccatggtggc	aatgagagac	tgattactac	ctgctggaga	60
ttgtttaagt	gagttaatat	attaaggata	aagggagcca	ggttttttga	ctgttgagga	120
aggaaattac	agatattgaa	ggtcccaa				148

<210> 230

<211> 257

<212> DNA

<213> Homo sapien

<400> 230

taagagggtg	cmaaaaaaaa	aaaatagaac	gaatgagtaa	gacctactat	ttgatagtag	60
aacagggtga	ctatagtcaa	tgataactta	attatacatt	taacatagag	tgtaattgga	120
ttgtttgtaa	ctcgaaggat	aaatgcttga	gaggatggat	acccattctt	ccatgatgta	180
cttatttcac	attacatgcc	tgtatcaaag	catctcatat	accctataaa	tatgtacacc	240
tactatgtac	cctcttta					257

<210> 231

<211> 260

<212> DNA

<213> Homo sapien

<400> 231

taagagggta	cgggtatttg	ctgatgggat	ttttttttct	ttctttttct	ttggaaaaca	60
aaatgaaagc	cagaacaaaa	ttattgaaca	aaagacaggg	actaaatctg	gagaaatgaa	120
gtcccctcac	ctgactgcca	tttcattcta	tctgaccttc	cagtctaggt	taggagaata	180
gggggtggag	gggattaatc	tgatacaggt	atattttaaag	caactctgca	tgtgtgccag	240
aagtccatgg	taccctctta					260

<210> 232

<211> 596

<212> DNA

<213> Homo sapien

<220>

<221> misc_feature

<222> (1)...(596)

<223> n = A,T,C or G

<400> 232

tgctcctctt	gccttaccaa	ccacaaatta	gaaccataat	gagatgtcac	ctcatacctg	60
gtgggattaa	cattatttaa	aaaatcagaa	gtattgacaa	ggatgtgaag	aaattagaac	120
atctgtgcac	tggttggtgg	aatgtaaaaa	aggtgtggcc	actatgggta	acagcatgaa	180
ggttcctcaa	aaaaaatttt	ttttaatcta	ctctatgata	gatcttgagg	ttgtttatgc	240
aaaagaactg	aatcaggat	tttgaggaaa	tattcacatt	cccacatcca	tttctgcttt	300
attcataata	ctcaagagat	ggaacaacc	taaatgtcca	tcccgggatg	aatggataaa	360
cacagtgtgg	tatatgcata	caatggaata	ttatttagtc	tttaaaaaga	aaaattctat	420
catatactac	aacttanatn	aaccttgagg	acacaatgct	nagtgaata	agccacggaa	480
ggacgaatac	tgcattatc	ccttatatga	agtatctaaa	gtggtcaaac	tcttanagca	540
naaaagtaaaa	atgggtgggt	gccanacagt	tggttaggcn	agaaganaaa	cctant	596

<210> 233

<211> 96

<212> DNA

<213> Homo sapien

<400> 233

tcttctgaag	acctttcgcg	actcttaagc	tcgtggttgg	taaggcaaga	ggagcgttgg	60
taaggcaaga	ggagcgttgg	taaggcaaga	ggagca			96

<210> 234

<211> 313

<212> DNA

<213> Homo sapien

<400> 234

tgtaagtcca	gcagtgtgat	gataaaactt	gaatggatca	atagttgctt	cttatggatg	60
agcaaagaaa	gtagtttctt	gtgatggaat	ctgctcctgg	caaaaatgct	gtgaacgttg	120
ttgaaaagac	aacaaagagt	ttagagtagt	acataaat	agaatagtag	ataaacttag	180
aatagtacat	aaacttagta	cataaataat	gcacgaagca	ggggcagggc	ttgagagaat	240
tgacttcaat	ttggaaagag	tatctactgt	aggtttagatg	ctctcaaaaca	gcacacact	300
gctcgactta	caa					313

<210> 235

<211> 550

<212> DNA

<213> Homo sapien

<400> 235
 aacgaggaca gatccttaaa aagaatgttg agtgaaaaaa gtagaaaaata agataatctc 60
 caaagtccag tagcattatt taaacatttt taaaaaatac actgataaaa attttgtaca 120
 tttcccaaaa atacatatgg aagcacagca gcatgaatgc ctatgggrtt gaggataggg 180
 gttgggagta gggatgggga taaaggggga aaataaaacc agagaggagt cttacacatt 240
 tcatgaacca aggagtataa ttatttcaac tattttgtacc wgaagtccag aaagagtgga 300
 ggcagaaggg ggagaagagg gcgaagaaac gtttttggga gaggggtccc asaagagaga 360
 ttttcgcat gtggcgctac atacgttttt ccaggatgcc ttaagctctg caccctattt 420
 ttctcatcac taatattaga ttaaaccctt tgaagacagc gtctgtggtt tctctacttc 480
 agctttccct ccggtgtctt cacacagtag ctgttttaca agggttgaac tgactgaagt 540
 gagattattc 550

<210> 236
 <211> 325
 <212> DNA
 <213> Homo sapien

<400> 236
 tagactgact catgtcccct accagagtag ctagaattaa tagcacaagc ctctacaccc 60
 aggaactcac tattgaatac ataaatggaa tttattcagc cttaaaaagt ttggaaggaa 120
 attctgacat atgctaaaac atggatgaac cttgaagact ttatgataag taaaagaagc 180
 cagtcataaa aggaaaaata ttgcatgatt ccacttatat gaggtaccta gagtagtcaa 240
 tttcatagaa acacaaaata gaatggtgtt tgccagggtt tttgaggaaa agggaatgac 300
 aagttagggg acatgagtca gtcta 325

<210> 237
 <211> 373
 <212> DNA
 <213> Homo sapien

<220>
 <221> misc_feature
 <222> (1)...(373)
 <223> n = A,T,C or G

<400> 237
 tagactgact catgtcccct atctactcaa catttccact tgaagtctga taggcatctc 60
 agacttatct tgtcccaaag caaactcttt atttcttttc atcctagtct ttatttcttg 120
 tgctgtctta cccatctcaa aagagtgccaa aaatccacca agttgctgaa acagaaatct 180
 aagaaatata cttgattctt ctttttccca tctacttcac ttctaattca ttagtaaata 240
 atctgtttca gaaaaccaa cactcatgt tctcactcat aagggggagt tgaacaatga 300
 gaacacacag acacagggag gggaacatca cacaccacgg cccgtcaggg agtangggac 360
 atgagtcagt cta 373

<210> 238
 <211> 492
 <212> DNA
 <213> Homo sapien

<220>
 <221> misc_feature
 <222> (1)...(492)
 <223> n = A,T,C or G

<400> 238
 tagactgact catgtcccct ataatgctcc caggcatcag aaagcatctc aaactggagc 60
 tgacaccatg gcagaggttt caggtaagtc acaaaagggg tcctaaagaa tttgccctca 120

67

```

ataticagagt gattagaaga agtggacaga gctacccaag ttaaaccatat gcgagataaa 180
aaaaatatgg cacttgtaga cacacactac aggaggaaaa taaggaaacat aatagcatat 240
tgtgctatta tgatgatgaa gaacctctct anaagaaaaac ataaccaaaag aaacaaagaa 300
aattcctgcn aatgtttaat gctatagaag aaattaacaa aaacatatat tcaatgaatt 360
cagaaaagtt agcaggtcan aagaaaacaa atcaaagacc agaataatcc catttttagat 420
tgtcgagtaa actanaacag aaagaatacc actggaaatt gaattcctac gtangggaca 480
tgantcantic ta 492

```

<210> 239

<211> 482

<212> DNA

<213> Homo sapien

<220>

<221> misc_feature

<222> (1)...(482)

<223> n = A,T,C or G

<400> 239

```

tggaagtat ttaatgatgg gcaacttgct gtttacttcc tacatatccc atcatcttct 60
gtattttttt aaataacttt tttttggatt tttaaagtaa ccttattctg agaggtaaca 120
tggtattacat acttctaagc cattaggaga ctctatgtta aaccaaaagg aaatgttact 180
agatcttcat ttgatcaata ggatgtgata atcatcatct ttctgctcta atggaaaagt 240
actanaaaca tggaaccata atcttagatg aacaacgtta gaatttgcac taattctacg 300
gaatttcagt aattcggcaa atgtcgggca gtgacacaac atttcatgac ggggacgcac 360
ctaccaactt ctggcgataa gggccaccct tccctctgta cttacagtcc catttcatac 420
acagtctttg attaaatatt cacatttttt ctctacctaa agaccttcaa gaccagtacg 480
ta 482

```

<210> 240

<211> 519

<212> DNA

<213> Homo sapien

<220>

<221> misc_feature

<222> (1)...(519)

<223> n = A,T,C or G

<400> 240

```

tgtatcgacg tagtggtctc cccatgtgat agtctgaaat atagcctcat gggatgagag 60
gctgtgccc agcccgacac ccgtaaaggg tctgtgctga ggtggattag taaaagagga 120
aagccttgca gttgagatag aggaagggca ctgtctcctg cctgccctg ggaactgaat 180
gtctcggtat aaaacccgat tgtacatttg ttcaattctg agataggaga aaaaccaccc 240
tatggcgagg ggcgagacat gttggcagca atgctgcctt gttatgcttt actccacaga 300
tgtttgggcg gagggaaaca taaatctggc ctacgtgcac atccaggcat agtacctccc 360
tttgaactta attatgacac agattccttt gctcacatgt ttttttgctg accttctcct 420
tattatcacc ctgtctcct accgcattcc ttgtgctgag ataataaaaa taatatcaat 480
aaaaacttga nggaactcgg agaccactac gtcgataca 519

```

<210> 241

<211> 771

<212> DNA

<213> Homo sapien

<220>

<221> misc_feature

<222> (1)...(771)

<223> n = A,T,C or G

<400> 241

tgtatcgacg	tagtgggtctc	cactccccgcc	ttgacggggc	tgctatctgc	cttccagggc	60
actgtcacgg	ctccccgggta	gaagtcactt	atgagacaca	ccagtgtggc	cttgtttggct	120
tgaagctcct	cagaggaggg	tggaacaga	gtgaccgagg	gggcagcctt	gggctgacct	180
aggacggta	gcttgggtccc	tccgccaaac	acgagagtgc	tgctgcttgt	atatgagctg	240
cagtaataat	cagcctcgtc	ctcagcctgg	agcccagaga	tggtcagggg	ggccgtgttg	300
ccanacttgg	agccagagaa	gcgattagaa	acccctgagg	gccgattacc	gacctcataa	360
atcatgaatt	tgggggcttt	gcctgggtgc	tgttggtacc	angagacatt	attataacca	420
ccaacgtcac	tgctgggtcc	antgcaggga	aaatgggtga	tcnaactgtc	caagaaaacc	480
actacgtcca	taccaatcca	ctaattgccn	gccgcctgca	ggttcaacca	tattggggaa	540
naactcccn	ccgccgtttg	ggattgncat	naacctttga	aattttttcc	tattanttgt	600
ccccctaaaa	taaaccnttg	ggcnttaatc	cattgggtcc	atancttntt	tncccggttt	660
ttaaaanttg	tttatccgc	cncccnattt	ccccccaac	tttccaaaac	ccgaaacnt	720
tnaaatttnt	tnaaacctg	gggggttccc	nnaattnnan	ttnaancnnc	c	771

<210> 242

<211> 167

<212> DNA

<213> Homo sapien

<400> 242

tgggcacctt	caatatcggg	ctcatcgata	acatcacgct	gctgatgctg	ctgttgctgg	60
tcctctctag	gaacctctgg	attttcaaat	tctttgagga	attcatccaa	attatctgcc	120
tctcctcctt	tcctcctttt	tctaaggtct	tctggtacaa	gcggta		167

<210> 243

<211> 338

<212> DNA

<213> Homo sapien

<400> 243

ttgggcacct	tcaatatcta	ctgatctaaa	tagtgtgggt	tgaggcctct	tgttctggc	60
taaaaatcct	tggcaagagt	caatctccac	tttacaatag	aggtaaaaa	cttacaatgg	120
atattcttga	caaagctagc	atagagacag	caattttaca	caaggatatt	ttcacctgtt	180
taataacagt	ggttttccta	cacccatagg	gtgccaccaa	gggaggagtg	cacagttgca	240
gaaacaaatt	aagatactga	agacaacact	acttaccatt	tcccgtatag	ctaaccacca	300
gttcaactgt	acatgtatgt	tcttatgggc	aatcaaga			338

<210> 244

<211> 346

<212> DNA

<213> Homo sapien

<400> 244

tttttggtc	ccatacagca	cactctcatg	ggaaatgtct	gttctaagg	caaccataa	60
tgcaaaaatc	atcaatatac	ttgaagatcc	ccgtgtaagg	tacaatgtat	ttaatattat	120
cactgataca	attgatccaa	taccagtttt	agctcggcat	tgaatcaaat	cactgttttt	180
gttgtataaa	aagagaaata	tttagcttat	atttaagtac	catattgtaa	gaaaaaagat	240
gcttatcttt	acatgctaaa	atcatgatct	gtacattgg	gcagtgaata	ttactgtaaa	300
aggggaaga	gaatgaagac	gagctaagga	tattgaagg	gcccaa		346

<210> 245

<211> 521

<212> DNA

<213> Homo sapien

<220>
 <221> misc_feature
 <222> (1)...(521)
 <223> n = A,T,C or G

<400> 245
 accaatccca cacggatact gagggacaag tatatcatcc catttcatcc ctacagcagc 60
 aacttcatga ggcaggagtt attagtccca ttttacagaa gaggaaactg agacttaggg 120
 agatcaagta atttgcccag gtcgcacaat tagtgataga gccagggtt gaagcgacgt 180
 ctgtcttaag ccaatgaccc ctgcagatta ttagagcaac tgttctccac aacagtgtaa 240
 gcctcttgct anaagctcag gtccacaagg gcagagattt ttgtctgttt tgctcattgc 300
 tccttcccca ttgcttagag cagggtctgc cacgaancag gttctcaatg catagtatt 360
 aaatgtatat aagagcaaac atatgttaca gagaactttc tgtatgcttg tcacttacat 420
 gaatcacctg tganatgggt atgcttggtc cccantggtg cagatnaaga tattgaangt 480
 gcccaaata ctaantgcgg gcgcctgcan gtccancata t 521

<210> 246
 <211> 482
 <212> DNA
 <213> Homo sapien

<220>
 <221> misc_feature
 <222> (1)...(482)
 <223> n = A,T,C or G

<400> 246
 tggaaccaat ccaaataccc atcaatgata gactggataa agaaaatttg gcacatgttc 60
 accatgaaat actatgcagc cataaaaaag gatgagttca tatcctttgc agggacatgg 120
 atgaagctgg agaccatcat tctcagcaaa ctaacaaggg aacagaaaac caaacactgc 180
 atgttctcac tcttaagtgg gagctgaaca atgagaacac atggacacag ggaggggaac 240
 atcacacagt ggggcctgct ggtgggtagg ggtctagggg agggatagca ttaggagaaa 300
 tacctaagt agatgacggg ttgatgggtg cagcaaacca ccatgacacg tgtataccta 360
 tgtaacaaac ctgcatgttc tgcacatgta cccagaact taaagtgtta ataaaaaat 420
 taagaaaaaa gttaagtatg tcatagatac ataaaatatt gtanatattg aaggtgccca 480
 aa 482

<210> 247
 <211> 474
 <212> DNA
 <213> Homo sapien

<220>
 <221> misc_feature
 <222> (1)...(474)
 <223> n = A,T,C or G

<400> 247
 ttcgatacag gcacagagta agcagaaaaa tggctgtggt ttaaccaagt gagtacagtt 60
 aagtgagaga ggggcagaga agacaaggc atatgcaggg ggtgattata acagggtggt 120
 gtgctgggaa gtgagggtac tcggggatga ggaacagtga aaaagtggca aaaagtggta 180
 agatcagtga attgtacttc tccagaattt gatttctggn ggagtcaaat aactatccag 240
 tttggggat catangcaaa cagttgaggt ataggaggta gaagtcncag tgggataatt 300
 gaggttatga anggtttggt actgactggt actgacaang tctgggttat gaccatggga 360
 atgaatgact gtanaagcgt anaggatgaa actattccac ganaaagggg tccnaaaact 420
 aaaaannnaa gnnnnngggg aatattattt atgtggatat tgaangtgcc caaa 474

<210> 248

<211> 355
 <212> DNA
 <213> Homo sapien

<220>
 <221> misc_feature
 <222> (1)...(355)
 <223> n = A,T,C or G

<400> 248
 ttcgatacag gcaaacaatga actgcaggag ggtggtgacg atcatgatgt tgccgatggt 60
 ccgatggnc acgaagacgc actggancac gtgcttacgt ccttttgctc tgttgatggc 120
 cctgagggga cgcaggaccc ttatgaccct cagaatcttc acaacgggag atggcactgg 180
 attgantccc antgacacca gagacacccc aaccaccagn atatcantat attgatgtag 240
 ttctgttaga nggccccctt gtggaggaaa gctccatnag ttggtcatct tcaacaggat 300
 ctcaacagtt tccgatggct gtgatgggca tagtcatant taaccntgtn tcgaa 355

<210> 249
 <211> 434
 <212> DNA
 <213> Homo sapien

<400> 249
 ttggattggt cctccaggag aacaagggga aaaagggtgac cgagggctcc ctggaactca 60
 aggatctcca ggagcaaaaag gggatggggg aattcctggt cctgctggtc ccttaggtcc 120
 acctgggtcct ccaggcttac caggctcctca aggcccaaag ggtaacaaag gctctactgg 180
 acccgctggc cagaaagggtg acagtgggtct tccagggcct cctgggcctc cagggtccacc 240
 tgggtgaagtc attcagcctt taccaatctt gtctccaaa aaaacgagaa gacatactga 300
 aggcattgcaa gcagatgcag atgataatat tcttgattac tcggatggaa tggagaagaat 360
 atttggttcc ctcaattccc tgaaacaaga catcgagcat atgaaatttc caatgggtac 420
 tcagaccaat ccaa 434

<210> 250
 <211> 430
 <212> DNA
 <213> Homo sapien

<220>
 <221> misc_feature
 <222> (1)...(430)
 <223> n = A,T,C or G

<400> 250
 tggattggtc acatggcaga gacaggatcc caaggcagtg agaggaggat acaatgcttc 60
 tcactagtta ttattattta ttttattttt gagatgaagt ctgctttgt ctcccaggct 120
 ggagagcggg ggtgcgatct tggctctctg caacccccgc ctcaagcaat tctcctgtct 180
 tagcctcgcg ggtagatgga attacaggcg cccaccgcca tgcccaacta atttttttgt 240
 gtcttcagta gagacagggt ttcgccatgt tgggcaggct ggtcttgaac tcctgacctc 300
 nagtgatctg ccctcctcgg cctcacaag tgctggaatt acaggcatgg gctgctgcac 360
 ccagtcaact tctcactagt tatggcctta tcattttcac cacattctat tggcccaaaa 420
 aaaaaaaaaa 430

<210> 251
 <211> 329
 <212> DNA
 <213> Homo sapien

<400> 251

71

tggtactcca	ccatyatggg	gtcaaccgcc	atcctcgccc	tcctcctggc	tgttctccaa	60
ggagtctgtg	ccgaggtgca	gctgrtgag	tctggagcag	aggtgaaaaa	gtccggggag	120
tctctgaaga	tctcctgtaa	gggttctgga	tacaccttta	agatctactg	gatcgcttgg	180
gtgcgccagt	tgccccggaa	aggcctggag	tggatggggc	tcattctttcc	tgatgactct	240
gataccagat	acagcccgtc	cttccaaggc	caggtcacca	tctcagtcga	taagtccttc	300
agcaccgcct	atctgcagtg	gagtaccaa				329

<210> 252

<211> 536

<212> DNA

<213> Homo sapien

<400> 252

tggtactcca	ctcagcccaa	ccttaattaa	gaattaagag	ggaacctatt	actattctcc	60
caggtctctc	tgctctaacc	aggttctctg	gacagtatta	gaaaaggatg	tctcaacaag	120
tatgtagatc	ctgtactggc	ctaagaagtt	aaactgagaa	tagcataaat	cagaccaaac	180
ttaatgggtc	ttgagacttg	tgtcctggag	cagctgggat	aggaaaaactt	ttgggcagca	240
agaggaagaa	ctgcctggaa	gggggcatca	tgttaaaaaat	tacaagggga	acccacacca	300
ggcccccttc	ccagctctca	gcctagagta	ttagcatttc	tcagctagag	actcacaact	360
tccttgctta	gaatgtgcc	ccggggggag	tcctctgtgg	tgatgaggct	ctcaagagtg	420
agagtggcat	cctatcttct	gtgtgcccac	aggagcctgg	cccagagactt	agcagggtgaa	480
gtttctgtgc	caggctttgc	ccttgactca	ctatgtgacc	tctggtggag	taccaa	536

<210> 253

<211> 507

<212> DNA

<213> Homo sapien

<220>

<221> misc_feature

<222> (1)...(507)

<223> n = A,T,C or G

<400> 253

ntgttgcgat	cccagtaact	cggaagctg	aggcgggag	atcacctgag	ctcaggaggt	60
tgaggccgca	gtgagccggg	accacgccac	tacactccag	cctggggcat	agagtggagc	120
cctccaagac	agaaaaagaa	agaaaggaag	ggaaagggaa	agggaagagg	aaaaggaaaa	180
ggaaaaggaa	aaggaaaaga	caagacaaaa	caagacttga	atttggatct	cctgacttca	240
attttatgtt	ctttctacac	cacaattcct	ctgcttacta	agatgataat	ttagaaaccc	300
ctcgttccat	tctttacagc	aagctggaag	tttgggtcaag	taattacaat	aatagtaaca	360
aatttgaata	ttatatgcc	ggtgtttttc	attcctgtct	tcacttaatt	ctcaccactc	420
tgatataaat	acaattgctg	ccgggtgtgg	tggctcatgc	ctgtaatccc	ggcacttttg	480
gagaccgagg	tgggcggats	gcaacaa				507

<210> 254

<211> 222

<212> DNA

<213> Homo sapien

<220>

<221> misc_feature

<222> (1)...(222)

<223> n = A,T,C or G

<400> 254

ttggattggt	cactgtgagg	aagccaaatc	ggatccgaga	gtctttttct	aaaggccagt	60
actggccaca	ctttctcctg	ccgccttcct	caaagctgaa	gacacacaga	gcaaggcgct	120
tctgttttac	tccccaatgg	taactccaaa	ccatagatgg	ttagctnccc	tgctcatctt	180

tccacatccc tgcatttcag tatagtcggt ggaccaatcc aa 222

<210> 255
 <211> 463
 <212> DNA
 <213> Homo sapien

<400> 255
 tgttgcgac cataaatgct gaaatggaaa taaacaacat gatgaggag gattaagttg 60
 gggaggagc acattaaggt ggccatgaag tttgttgaa gaagtgaact ttgaacaagg 120
 ccttggtgtt aagagctgat gagagtgtcc cagacagagg gccactggt acaatagacg 180
 agatgggaga gggcttgaa ggtgtgcgaa ataggaggga gttgttctg gtatgagctt 240
 agtgaacaca gaggcgagag gccctggtg gtgcagctgg agagtattgc agaataacat 300
 taggccctgt gggggactgt agactgtcag caataatcca cagtttggat ttatttctaa 360
 gagtgatggg aagcgtgga aaggggtta agcaaggagt gaaattatca gatttacagt 420
 gataaaaata aattggtctg gctactgggg aaaaaaaaaa aaa 463

<210> 256
 <211> 262
 <212> DNA
 <213> Homo sapien

<400> 256
 ttggattggt caacctgtct aactctacyt ttctctcttc ttcttaaaaa attaatgaat 60
 ccaatacatt aatgcaaaaa cccttggtgt ttatcaatat ttctgttaaa aagtattatc 120
 cagaactgga cataatacta cataataata cataacaacc ccttcattctg gatgcaaaca 180
 tctattaata tagcttaaga tcactttcac ttacagaag caacatcctg ttgatgttat 240
 ttgatgttt ggaccaatcc aa 262

<210> 257
 <211> 461
 <212> DNA
 <213> Homo sapien

<220>
 <221> misc_feature
 <222> (1)...(461)
 <223> n = A,T,C or G

<400> 257
 gnggnnnnnn nnncaattcg actcngttcc cntggtancc ggtcgacatg gccgcgggat 60
 taccgcttgt nnctgggggt gtatggggga ctatgaccgc ttgtagctgg ggtgtatgg 120
 gggactatga ccgcttgtag mtggkgtgt atgggggact atgaccgctt gtcgggtggt 180
 cgataaaacc gacgcaaggg acgtgatcga agctgcgttc ccgctcttgc gcatcggtag 240
 ggatcatgga cagcaatata cgcattcgyt tgaaggcggt cgaccatcgc gtgctcgatc 300
 aggcgaccgg cgacatcgcc gacaccgcac gccgtaccgg cgcgctcatc cgcggtccga 360
 tcccgcttcc cagcgcacg gagaagttca cggtaaccg tggcccgcac gtcgacaaga 420
 agtcgcgcga gcagttcgag gtgcgtacct acaagcggtc a 461

<210> 258
 <211> 332
 <212> DNA
 <213> Homo sapien

<220>
 <221> misc_feature
 <222> (1)...(332)
 <223> n = A,T,C or G

```

<400> 258
tgaccgcttg tagctggggg tgtatggggg actacgaccg cttgtagctg ggggtgtatg      60
ggggactatg accgcttgta gctgggggtg tatgggggac tatgaccgct tgtagctggg    120
ggtgtatggg ggactaggac cgcttgtagc tgggggtgta tgggggacta tgaccgcttg    180
tagctggggg tgtatggggg actacgaccg cttgtagctg ggggtgtatg ggggactatg    240
accgcttgta nctgggggtg tatgggggac tatgaccgct tgtgctgcct gggggatggg    300
aggagagttg tggttgggga aaaaaaaaaa aa                                  332

```

```

<210> 259
<211> 291
<212> DNA
<213> Homo sapien

<220>
<221> misc_feature
<222> (1)...(291)
<223> n = A,T,C or G

```

```

<400> 259
taccgcttgt gaccgcttgt gaccgcttgt gaccgcttgt gaccgcttgt gaccgcttgt      60
gaccgcttgt gaccgcttgt gaccgcttgt gaccgcttgt gaccgcttgt gaccgcttgt    120
gaccgcttgt gaccgcttgt nacngggggt gtctggggga ctatgannga ntgtactgg      180
gggtgtctgg gggncatga nngantgtna cnggggggtg ctgggggact atganngact    240
gtgcnnctg ggggatcnga ggagantngn ggntagngat ggttngggan a              291

```

```

<210> 260
<211> 238
<212> DNA
<213> Homo sapien

```

```

<400> 260
taagagggtg ctggttaaaa tacaggaaat ctggggtaat gaggcagaga accaggatac      60
tttgagggtc gggatgaaaa ctagaatgtt tttctttttt tttgcctgag aaacttgctg    120
ctctgaagag gccatgtat taattgcttt gatcttcctt ttcttacagc cctttcaagg    180
gcagagccct ccttatcctg aaggaatcct atccttagct atagtatgta ccctctta     238

```

```

<210> 261
<211> 746
<212> DNA
<213> Homo sapien

```

```

<220>
<221> misc_feature
<222> (1)...(746)
<223> n = A,T,C or G

```

```

<400> 261
ttgggcacct tcaatatcaa tagctaacat ttattgagtg tttatcgtat cataaaacac      60
tgttctaagc ctttaaacgt actaattcat ttaatgctca taatcacttt agaaggtggg    120
tactagtatt agtctcattt acagatgcaa catgcaggca cagagagggt aattaacttg    180
ccaagggtaa cacagctaag aaatagaaaa aatattgaat ctggaaagtt gggcttctgg    240
gtaacccaca gagtcttcaa tgagcctggg gcctcactca gtttgctttt acaaagcgaa    300
tgagtaacat cacttaattc agtgagtagg ccaaatggag gtcagctacg agtttctgct    360
gttcttgtag tggactgaca gatgtttaca acgtctggcc atcagtwaat ggactgatta    420
tcattgggaw gtgggtgggc tgaatgttgg ccagtgaagt ttattcawgc catattttta    480
tgtttaggat gacttttggc tggtcctagg gcaagctctg tctgscacgg aacacagaat    540
wacacaggga cccctcaat ttctgggtg gctagaacca tgaaccactg gttgggggaa      600

```

caagcgggtca	aaacctaagt	gcggccggct	ggcaggggtcc	acccatatgg	ggaaaactcc	660
cnacgcgttt	ggaatgcctn	agctngaatt	attctaana	ttgtccnct	aaaattagcc	720
tgggcggttaa	tcangggctn	naagcc				746

<210> 262
 <211> 588
 <212> DNA
 <213> Homo sapien

<220>
 <221> misc_feature
 <222> (1)...(588)
 <223> n = A,T,C or G

<400> 262						
tgaccgcttg	tcatctcaca	tggggtcctg	cacgcttttg	cctttgtagg	aaacctgaca	60
tttgtctgtt	tcttctttct	cttttccttc	ccatatactc	ctaatttacg	tttgacttgt	120
ttgctgagga	ggcaggagct	agagactgct	gtgagctcat	aggggtggga	agtttatcct	180
tcaagtcccg	cccactcatc	actgcttctc	accttcccct	gaccaggctt	acaagtgggt	240
tcttgctgc	tttccctttg	gacccaacaa	gcccctgtaa	tgagtgtgca	tgactctgac	300
agctgtggac	tcagggctct	tggtacagc	tgccatgtaa	aatatctcat	ccagttctcg	360
caaattgtta	aaataaccac	atttcttaga	ttccagtacc	caaatcatgt	ctttacgaac	420
tgctcctcac	acccagaagt	ggcacaataa	ttcttgggga	attattactt	ttttttttct	480
ctctnttnc	gnnnngnnng	gnnnngnccag	gaattaccac	nttggaagac	ctggccngaa	540
tttattatan	aggggagccg	attntttttc	ctaacacaaa	gcgggtca		588

<210> 263
 <211> 730
 <212> DNA
 <213> Homo sapien

<220>
 <221> misc_feature
 <222> (1)...(730)
 <223> n = A,T,C or G

<400> 263						
tttttttttt	tttggcctga	gcaactgaaa	ttatgaaatt	tccatatact	caaaagagta	60
agactgcaaa	aagattaaat	gtaaaagttg	tcttgatatac	agtaatgttt	aagataccta	120
ttanatttat	aaatggaaaa	ttagggcatt	tgatataca	agttgaaaat	tcaggagtga	180
ggttgggctg	gctgggtata	tactgaaaac	tgctagtaca	cagatgacat	ctaaaaccac	240
aaatctggtt	ttattttagc	agtgatatgt	gtcactccca	caaaagcctt	cccaattggc	300
ctcagcatat	acaacaagtc	acctccccac	agccctctac	acataaacia	attccttagt	360
ttagttcagg	aggaaatgcg	cccttttcct	tccgctctag	gtgaccgcaa	ggcccagttc	420
tcgtcaccaa	gatgttaagg	gaagtctgcc	aaagaggcat	ctgaaaggaa	ataaggggaa	480
tggtgagtgc	cacaaaggaa	agccaaggan	aaactttgga	gaccgtttct	aganccctgg	540
catttcacaa	caaaactcng	gaacaaacct	tgtctcatca	atcatttaag	cccttcgttt	600
ggannagact	ttctgaactg	ggcgctgaac	ataancctca	ttgaatgtct	tcacagtctc	660
ccagctgaag	gcacaccttg	ggccagaagg	ggaatcttcc	aggtcctcaa	nacagggtctc	720
gccctttgnc						730

<210> 264
 <211> 715
 <212> DNA
 <213> Homo sapien

<220>
 <221> misc_feature

75

<222> (1)...(715)

<223> n = A,T,C or G

<400> 264

tttttttttt	tttggccagt	atgatagtct	ctaccactat	attgaagctc	ttaggtcatt	60
tacacttaat	gtggttatag	atgctgttga	gcttacttct	accaccttgc	tatttctccc	120
gtctcttttt	tgttcccttt	ctcttctttt	cctcccttat	tttataattg	aatttttttag	180
gattctattt	tatatagatt	tatcagctat	aacactttgt	attcttttgt	tttgtgggtc	240
ttctgtcatt	tcaatgtgca	tcttaaaactc	atcacaaatct	attttcaaat	aatatcataat	300
aaccttacat	ataatgtaag	aatctaccac	catatatttc	catttctccc	ttccatccta	360
tgntgtcat	attttttctt	ttatatatgt	tttaaagaca	taatagtata	tgggagggtt	420
ttgcttaaaa	tgtgatcaat	attccttcaa	ngaaacgtaa	aaattcaaaa	taaatntctg	480
tttattctca	aatnnaccta	atatttctta	ccatntctna	tacntttcaa	gaatctgaag	540
gcattgggtt	tttccggctt	aagaacctcc	tctaagcac	tctaagcaga	attaagtctt	600
ctgggagagg	aattctccca	agcttggggc	ttnanmtgta	ctccntnang	gttaaanntt	660
ggccgggaaa	tagaaattcc	aagttaacag	gntanttttt	nttttnttn	tcncc	715

<210> 265

<211> 152

<212> DNA

<213> Homo sapien

<400> 265

tttttttttt	tttcccaaca	caaagcacca	ttatctttcc	tcacaatfff	caacatagtt	60
tgattcccat	gaagaggtta	tgatttctaa	agaaaacatg	gctactatac	tatcaatcag	120
ggttaaatct	tttttttttg	agacggagtt	ta			152

<210> 266

<211> 193

<212> DNA

<213> Homo sapien

<220>

<221> misc_feature

<222> (1)...(193)

<223> n = A,T,C or G

<400> 266

taaactccgt	ccccttctta	atcaatatgg	aggctaccca	ctccacatta	ccttcttttc	60
aagggactgt	ttccgtaact	gttgtgggta	ttcacgacca	ggcttctaaa	cctcttaaaa	120
ctccccaatt	ctggtgccaa	cttggaacaac	atgctttttt	tttttttttt	tttttttttn	180
gagacggagt	tta					193

<210> 267

<211> 460

<212> DNA

<213> Homo sapien

<400> 267

tggtgcatc	ccttaagcat	gggtgctatt	aaaaaaatgg	tggagaagaa	aatacctgga	60
atttacgtct	tatcttttaga	gattgggaag	accctgatgg	aggacgtgga	gaacagcttc	120
ttcttgaatg	tcaattccca	agtaacaaca	gtgtgtcagg	cacttgctaa	ggatcctaaa	180
ttgcagcaag	gctacaatgc	tatgggatcc	tcccaggagg	gccaatctct	gagggcagtg	240
gctcagagat	gcccttcacc	tcccatgatc	aatctgatct	cggttggggg	acaacatcaa	300
gggtgttttg	gactccctcg	atgccagga	gagagctctc	acatctgtga	cttcatccga	360
aaaacactga	atgctggggc	gtactccaaa	gttggtcagg	aacgcctcgt	gcaagccgaa	420
tactggcatg	accataaaa	ggaggatgtg	gatcgcaaca			460

76

<210> 268
 <211> 533
 <212> DNA
 <213> Homo sapien

<220>
 <221> misc_feature
 <222> (1)...(533)
 <223> n = A,T,C or G

<400> 268
 tgttgcatc cgttgataga atagcgacgt ggtaatgagt gcatggcacg cctccgactt 60
 accttcgcc gtggggaacc cgagtacgtc tacggcgctc tcaactagag taccctctgg 120
 acgcccgggc gcgttcgatt taccggaagc gcgagctgca gtgggcttgc gccccggcc 180
 aaattctttg ggggggttaa ggcgcgggg aatttgaggt atctctatca gtatgtagcc 240
 aagttggaac agtcgccatt cccgaaatcg ctttctttga atccgcaccg cctccagcat 300
 tgcctcattc atcaacctga aggcacgcat aagtgcgggt tgtgtcttca gcagctccac 360
 tccataacta gcgcgctcga cctcgtcttc gtacgcgcca ggtccgtgcg tgcgaattcc 420
 caactccggt gagttgcgca tttcaagtn cgaaactgtt cgctccacn atttggcatg 480
 ttcacgcatg acacggaata aactcgtcca gtaccgggaa tgggatcgca aca 533

<210> 269
 <211> 50
 <212> DNA
 <213> Homo sapien

<400> 269
 tttttttttt ttgcctgaa ttagctacag atcctcctca caagcgttca 50

<210> 270
 <211> 519
 <212> DNA
 <213> Homo sapien

<400> 270
 tgttgcatc caaataaacc accagcttct tgcacacttc gcagaagcca cgtcctttg 60
 gctgagtcac gtgaacgggtc agtgcaagca gccgcgtgcc agagcagagg tgcagcatgc 120
 tgcacaccag ctacagggtg acctcctcca gcaggatgga caggatggag ctgccgtacg 180
 tgtccaccac ctctggcac tcttccgaca gggacttcgg cagcttcgag cacattttgt 240
 caaaagcgtc gagtatttct ttctcagtct tgttggtgtc aatcagcttg gtcacctcct 300
 tcaccaggaa ttacacaccc tcacagtaaa catcagactt tgctgggacc tctgtcttct 360
 taatgggctc caccagttcc agggcagggg tgacattctt ggaggccact ttggcgggga 420
 ccagagtctg catgggcacg tctttcacct catcacagaa cccaaccagc gcacagatct 480
 ccttgggttg catgtgcacg atcatctggg atcgcaaca 519

<210> 271
 <211> 457
 <212> DNA
 <213> Homo sapien

<400> 271
 tttttttttt ttccggcggc gaccggacgt gcactcctcc agtagcggct gcacgtcgtg 60
 ccaatggccc gctatgagga ggtgagcgtg tccggcttcg aggagtcca ccgggccgtg 120
 gaacagcaca atggcaagac cattttcgcc tactttacgg gttctaagga cggcggggg 180
 aaaagctggt gccccgactg cgtgcaggct gaaccagtcg tacgagagg gctgaagcac 240
 attagtgaag gatgtgtgtt catctactgc caagtaggag aagagcctta ttggaaagat 300
 ccaaataatg acttcagaaa aaacttgaaa gtaacagcag tgcctacact acttaagtat 360
 ggaacacctc aaaaactggt agaactcgag tgtcttcagg ccaacctggt ggaaatgttg 420

ttctctgaag attaagattt taggatggca atcaaga 457

<210> 272
<211> 102
<212> DNA
<213> Homo sapien

<400> 272

tttttttttt ttgggcaaca acctgaatac cttttcaagg ctctggcttg ggctcaagcc 60
cgcaggggaa atgcaactgg ccaggtcaca gggcaatcaa ga 102

<210> 273
<211> 455
<212> DNA
<213> Homo sapien

<220>
<221> misc_feature
<222> (1)...(455)
<223> n = A,T,C or G

<400> 273

tttttttttt ttggcaatca acagggtttaa gtcttcggcc gaagttaatc tcgtgttttt 60
ggcaatcaac aggttttaagt cttcggccga agttaatctc gtgttttttg caatcaacag 120
gtttaagtct tcggccgaag ttaatctcgt gtttttggca atcaacagggt ttaagtcttc 180
ggccgaagtt aatctcgtgt ttttggcaat caacagggtt aagtcttcgg ccgaagttaa 240
tctcgtgttt ttggcaatca acagggtttaa gtcttcggcc gaagttaatc tcgtgttttt 300
ggcaatcaag aggttttaagt cttcggccga agttaatctc gtgttttttg caatcaacag 360
gtttaagtct tcggccgaan ttaatctcgt gtttttggca atcaacagggt ttaantcttc 420
ggccgaagtt aatctcgtgt ttttggcaat caana 455

<210> 274
<211> 461
<212> DNA
<213> Homo sapien

<400> 274

tttttttttt ttggccaata ccttgatga acatcaatgt gaaaatcctc ggtaaaatac 60
tggcaaacca aatccagcag cacatcaaaa agcttatcca ccatgatcaa gtgggcttca 120
tccctgggat gcaaggctgg ttcaacataa gaaaatcaat aaatgtaatc catcacataa 180
acagaaccaa agacaaaaac cacatgatta tctcaataga tgcagaaaag gccttggaca 240
aattcaacag cccttcatgc taaacactct taataaacta gatattgatg gaatgtatct 300
caaaataata agagctatct atgacaaacc cacagccaat atcatactga atgggcaaag 360
actggaagca ttccctttga aaactggcac aagacaagga tgccctctct caccgctcct 420
attcaacata gtattggaag ttctggccag ggcaatcaag a 461

<210> 275
<211> 729
<212> DNA
<213> Homo sapien

<220>
<221> misc_feature
<222> (1)...(729)
<223> n = A,T,C or G

<400> 275

tttttttttt ttggccaaca ccaagtcttc cacgtgggag gttttattat gttttacaac 60

catgaaaaca	taggaaggtg	gctgttacag	caaacatttc	agatagacga	atcggccaaag	120
ctccccaac	cccaccttca	cagcctcttc	cacacgtctc	ccanagattg	ttgtccttca	180
cttgcaaatt	canggatgtt	ggaagtngac	attnnagtn	gcnggaaccc	catcagtga	240
ncantaagca	gaantacgat	gactttgana	nacanctgat	gaagaacacn	ctacnganaa	300
ccctttctnt	cgtgttanga	tctcnngtcc	ntcactaatg	cggccccctg	cnggtccacc	360
atttgggaga	actcccccn	cgttggatcc	cccccttgagt	ntcccattct	ngtccccan	420
accngncttg	ngngncantn	cnnectcnca	cctgttttcc	ctgnngtnaa	aatnngtttt	480
nccgcncccc	naattccac	ccnaatcaca	gcgaanceng	aaggccttcn	naagtgttta	540
angcccnng	gtttcctcnt	ntanttgag	cctaccctcc	cncttnnnmt	tncgngttgg	600
tgcgcacctg	gncncgectn	gttctctctt	nnggnnacia	cctngntcnn	nggcncntcn	660
nnctnttcc	tnnnactagc	tngectntcc	ncnccngngn	ncanngcaca	ttncncnnac	720
tntgtnncc						729

<210> 276
 <211> 339
 <212> DNA
 <213> Homo sapien

<400> 276						
tgacctgaca	tgtagtagat	acttaataaa	tatttgtgga	atgaatggat	gaagtggagt	60
tacagagaaa	aatagaaaag	tacaaattgt	tgtagtggtt	ttgaaggaaa	attatgatct	120
ttcccaaagt	tctgacttca	ttctaagaca	gggttagtat	ctccatacat	aattttactt	180
gcttttgaaa	atcaaatgag	ataatctatt	tagattgata	atttatttag	actggctata	240
aactattaag	tgctagcaaa	tatacatctt	aatctcattt	tccacctctt	gtgatatagc	300
tatgtagggtg	ttgactttaa	tgtagtgcag	gtcaatccc			339

<210> 277
 <211> 664
 <212> DNA
 <213> Homo sapien

<220>						
<221> misc_feature						
<222> (1)...(664)						
<223> n = A,T,C or G						
<400> 277						
tgacctgaca	tcataaacia	aatctttctc	catttatattc	ttctagggga	atttcttgaa	60
aagcatccaa	aggaaaciaa	tgatggtaag	accgtgccaa	gtggggagca	gacaccaaa	120
taagaccaca	gattttacat	tcaacaggta	gtcacagta	ctttgcccg	caactgtggc	180
agaaatagcc	tcctaattga	agccctggct	cagtattgcc	atccaaatgc	gccatgctga	240
aagagggttt	tgcatcctgg	tcagatnaag	aagcaatggt	gtgctgagga	aatcccatac	300
gaataagtga	gcattcagaa	cttgagctag	caggaggagg	actaatatga	tgtgtgagca	360
actctttgta	atggctttca	tctaaaataa	catggtacgt	gccaccagtt	tcacgagcaa	420
gtacagtga	aacgcgaact	tctgcagaca	atccaataac	agatactcta	atttttagctg	480
ccttttaggt	cttgattaaa	tcataaatat	tagatggatc	gcaagttgta	aggntgctaa	540
aagatgatta	gtacttctcg	acttgtatgt	ccaggcatgt	tgttttaaan	tctgccttag	600
nccctgctta	ggggaatttt	taaagaagat	ggctctccat	gttcanggtc	aatcacnaat	660
tgcc						664

<210> 278
 <211> 452
 <212> DNA
 <213> Homo sapien

<220>
 <221> misc_feature
 <222> (1)...(452)

<223> n = A,T,C or G

<400> 278

tgacctgaca	ttgaggaaga	gcacacacct	ctgaaattcc	ttagggttcag	aagggcattt	60
gacacagagt	gggcctctga	taattcatga	aatgcattct	gaagtcaccc	agaatggagg	120
ctgcaatctg	ctgtgctttg	ggggttgcc	cactgtgctc	ctggatatca	cacaaaagct	180
gcaatccttc	ttcttcaact	aacattttgc	agtatttgct	gggattttta	ctgcagacat	240
gatacatagc	ccatagtgcc	cagagctgaa	cctctggttg	agagaagttg	ccaaggagcg	300
ggaaaaatgt	cttgaaagat	ctataggtca	ccaatgctgt	catcttacia	cttgaacttg	360
gccaatctctg	tatggttgca	tgcagatctt	ggagaagagt	acgcctctgg	aagtcacggg	420
atatccaaan	ctgtctgtca	gatgtcaggt	ca			452

<210> 279

<211> 274

<212> DNA

<213> Homo sapien

<400> 279

tttttttttt	ttcggcaagg	caaatttact	tctgcaaaag	ggtgctgctt	gcacttttgg	60
ccactgcgag	agcacaccaa	acaaagtagg	gaaggggttt	ttatccctaa	cgcggttatt	120
ccctggttct	gtgtcgtgtc	cccattggct	ggagtcagac	tgcacaatct	acactgaccc	180
aactggctac	tgtttaaaat	tgaatatgaa	taattaggta	ggaaggggga	ggctgtttgt	240
tacgggtacaa	gacgtgtttg	ggcatgtcag	gtca			274

<210> 280

<211> 272

<212> DNA

<213> Homo sapien

<400> 280

tacctgacat	ggagaaataa	cttgtagtat	tttgcgtgca	atggaatact	atatgagggg	60
gaaaatgaat	gaactagcaa	tgcgtgtatc	aacatgaata	aatcccaaaa	acataataat	120
gttgaatgga	aaaggtgagt	ttcagaagga	tatatatgcc	ctctaaatcc	atttatgtaa	180
acctttaaaa	aactacatta	tttatgggtca	taagtcctac	cagaaaatat	ttaaaaacct	240
acatgggatt	gataactact	gatgtcaggt	ca			272

<210> 281

<211> 431

<212> DNA

<213> Homo sapien

<220>

<221> misc_feature

<222> (1)...(431)

<223> n = A,T,C or G

<400> 281

tttttttttt	ttggccaata	gcatgattta	aacattggaa	aaagtcaa	gagcaatgcg	60
aatttttatg	ttctcttgaa	taatcaaaag	agtaggcaac	attggttcc	cattcttgaa	120
tagcattaat	cagaaaatat	tgcatagcct	ctagcctcct	tagagtaggt	gtgtctcttc	180
aaatatatca	tagtcccaca	gtttatttca	tgtatatatt	ctgcctgaat	cacatagaca	240
tttgaatttg	caacgcctga	tgtaaaatata	taaattctta	ccaatcagaa	acatagcaag	300
aaattcaggg	acttggatca	yatcagggtg	tgacagcana	tccctgtara	aacactgata	360
cacactcaca	cacgtatgca	acgtggagat	gtcgcyyttw	kkktwywcm	rmrycrwcm	420
aatcacttan	n					431

<210> 282

<211> 98

80

<212> DNA

<213> Homo sapien

<400> 282

attcgattcg	atgcttgagc	ccaggagttc	aagactgcag	tgagccactg	cacttcaggc	60
tggaacaacag	agcgaggtccc	tgtgccaaaa	aaaaaaaa			98

<210> 283

<211> 764

<212> DNA

<213> Homo sapien

<220>

<221> misc_feature

<222> (1)...(764)

<223> n = A,T,C or G

<400> 283

tttttttttt	ttcgcaagca	cgtgcacttt	attgaatgac	actgtagaca	ggtgtgtggg	60
tataaactgc	tgtatctagg	ggcaggacca	agggggcagg	ggcaacagcc	ccagcgtgca	120
gggccascac	tgcacagtgg	astgcaaagg	ttgcaggcta	tgggcggcta	ctavtaacc	180
cgtttttcct	gtattatctg	taacataata	tggtagactg	tcacagagcc	gaatwccart	240
hacasgatga	atccaawggg	caygaggatg	cccasaatca	gggcccasat	sttcaggcac	300
ttggcgggtg	gggcataagc	ctgkgccccc	gtcacgtcsc	caaccwtcty	cctgtcccta	360
cmcttgawtc	cncncctttn	nntncctna	tntgcccgcc	cncctcctng	ngtcaaccng	420
natctgcact	anctccctcn	ccccttntgg	antctentcc	ttcaantaan	nttatccctn	480
acnccccctc	cncctttccc	ctnccncccn	tnatcccnng	nccnctatca	ntcntnccct	540
cncntnctcn	cnnatcggtc	cncctnntaa	ctacnctttn	nacnannccct	cactnatncc	600
ngnnantttc	ttccttccct	cccnaecgcn	tgcgtgcgcc	cgtctngcct	nnnctnecna	660
cccnactttt	atttaccctt	ncaccctagc	nctctacttn	acccancnc	tcctacctcc	720
nggnccaccc	nnccttnatc	nctnnctctn	tcnnctentt	cccc		764

<210> 284

<211> 157

<212> DNA

<213> Homo sapien

<400> 284

caagtgtagg	cacagtgatg	aaagcctgga	gcaaacacaa	tctgtgggta	attaacgttt	60
atttctcccc	ttccaggaac	gtcttgcag	gatgatcaaa	gatcagctcc	tggtcaacat	120
aaataagcta	gtttaagata	cgttccccta	cacttga			157

<210> 285

<211> 150

<212> DNA

<213> Homo sapien

<400> 285

attcgattgt	actcagacaa	caatatgcta	agtggaagaa	gtcagtcaca	aaagaccaca	60
tactgtatga	cttcatctac	attaagtgtc	cagaataggc	aaatccgtag	agacagaaag	120
tagatgagca	gctgcctagg	tctgagtaca				150

<210> 286

<211> 219

<212> DNA

<213> Homo sapien

<400> 286

attcgatttt	tttttttttg	gccatgatga	aattcttact	ccctcagatt	ttttgtctgg	60
ataaatgcaa	gtctcaccac	cagatgtgaa	attacagtaa	actttgaagg	aatctcctga	120
gcaaccttgg	ttaggatcaa	tccaatattc	accatctggg	aagtcaggat	ggctgagttg	180
caggtcttta	caagttcggg	ctggattggg	ctgagtaca			219

<210> 287
 <211> 196
 <212> DNA
 <213> Homo sapien

<400> 287						
attcgattct	tgaggctacc	aggagctagg	agaagaggca	tggaacaaat	tttccctcat	60
atccatactc	agaaggaacc	aaccctgctg	acaccttaat	ttcagcttct	ggcctctaga	120
actgtgagag	agtacatttc	tcttggttta	agccaagaga	atctgtcttt	tggtacttta	180
tatcatagcc	tcaaga					196

<210> 288
 <211> 199
 <212> DNA
 <213> Homo sapien

<400> 288						
attcgatttc	agtccagtcc	cagaacccac	attgtcaatt	actactctgt	araagattca	60
tttgttgaaa	ttcattgagt	aaaacattta	tgatccctta	atatatgcca	attaccatgc	120
taggtactga	agattcaagt	gaccgagatg	ctagcccttg	ggttcaagtg	atccctctcc	180
cagagtgcac	tggaactgaa					199

<210> 289
 <211> 182
 <212> DNA
 <213> Homo sapien

<400> 289						
attcgattct	tgaggctaca	aacctgtaca	gtatgttact	ctactgaata	ctgtaggcaa	60
tagtaataca	gaagcaagta	tctgtatatg	taaacattaa	aaaggtagag	tgaaaacttca	120
gtattataat	cttagggacc	accattatat	atgtggtcca	tcattggcca	aaaaaaaaaa	180
aa						182

<210> 290
 <211> 1646
 <212> DNA
 <213> Homo sapien

<400> 290						
ggcacgagga	gaaatgtaat	tccatatttt	atttgaaact	tattccatat	tttaattgga	60
tattgagtga	ttgggttatc	aaacacccac	aaactttaat	tttgtaaata	ttatatggct	120
ttgaaataga	agtataagtt	gctaccattt	tttgataaca	ttgaaagata	gtattttacc	180
atctttaatc	atcttggaat	atacaagtcc	tgtgaacaac	cactctttca	cctagcagca	240
tgaggccaaa	agtaaaggct	ttaaattata	acatatggga	ttcttagtag	tatgtttttt	300
tcttgaaact	cagtggctct	atctaacctt	actatctcct	cactctttct	ctaagactaa	360
actctaggct	cttaaaaaatc	tgcccacacc	aatcttagaa	gctctgaaaa	gaattttgtct	420
ttaaatatct	tttaatatga	acatgtattt	tatggacca	attgacattt	tcgactattt	480
tttccaaaaa	agtcagggtga	atttcagcac	actgagttgg	gaatttctta	tcccagaaga	540
ccaaccaatt	tcataatttat	ttaaagattga	ttccatactc	cgttttcaag	gagaatccct	600
gcagctcctc	ttaaaggtaga	acaaataactt	tctatttttt	tttcaccatt	gtgggatttg	660
actttaagag	gtgactctaa	aaaaacagag	aacaaatatg	tctcagttgt	attaagcacg	720
gacccatatt	atcatattca	cttaaaaaaa	tgatttcctg	tgacaccttt	ggcaacttct	780
cttttcaatg	tagggaaaaa	cttagtcacc	ctgaaaaacc	acaaaataaa	taaaacttgt	840

agatgtgggc	agaaggtttg	ggggtggaca	ttgtatgtgt	ttaaattaaa	ccctgtatca	900
ctgagaagct	gttgtatggg	tcagagaaaa	tgaatgctta	gaagctgttc	acatcttcaa	960
gagcagaagc	aaaccacatg	tctcagctat	attattatgt	atgttttatg	cataaagtga	1020
atcatttctt	ctgtattaat	ttccaaaggg	ttttaccctc	tatttaaatg	ctttgaaaaa	1080
cagtgcattg	acaatgggtt	gatatatttc	tttaaaagaa	aaatataatt	atgaaagcca	1140
agataatctg	aagcctgttt	tattttaaaa	ctttttatgt	tctgtggttg	atgttggttg	1200
tttgtttgtt	tctattttgt	tggtttttta	ctttgttttt	tgttttgttt	tgttttgttt	1260
kgcatactac	atgcagttct	ttaaccaatg	tctgtttggc	taatgtaatt	aaagttgtta	1320
atttatatga	gtgcatttca	actatgtcaa	tggtttctta	atatttatgt	tgtagaagta	1380
ctggtaattt	ttttattttac	aatatgttta	aagagataac	agtttgatat	gttttcatgt	1440
gttttatagca	gaagttatgt	atttctatgg	cattccagcg	gatattttgg	gttttgcgag	1500
gcatgcagtc	aatattttgt	acagtttagtg	gacagtattc	agcaacgcct	gatagcttct	1560
ttggccttat	gttaataaaa	aagacctgtt	tgggatgtat	tttttatttt	taaaaaaaaa	1620
aaaaaaaaaa	aaaaaaaaaa	aaaaaa				1646

<210> 291

<211> 1851

<212> DNA

<213> Homo sapien

<400> 291

tcatcaccat	tgccagcagc	ggcaccgtta	gtcagggtttt	ctgggaatcc	cacatgagta	60
cttcctgtgt	cttcattctt	cttcaatagc	cataaatctt	ctagctctgg	ctggctgttt	120
tcaacttctt	taagcctttg	tgactcttcc	tctgatgtca	gcttttaagtc	ttgttctgga	180
ttgtctgttt	cagaagagat	ttttaacatc	tgtttttctt	tgtagtcaga	aagtaactgg	240
caaattacat	gatgatgact	agaaacagca	tactctctgg	ccgtcttctcc	agatcttgag	300
aagatacatc	aacatttttg	tcaagtagag	ggctgactat	acttgctgat	ccacaacata	360
cagcaagtat	gagagcagtt	cttccatata	tatccagcgc	atttaaatcc	gcttttttct	420
tgattaaaaa	tttcaccact	tgctgttttt	gctcatgtat	accaagtagc	agtgggtgtga	480
ggccatgctt	gttttttgat	tcgatatacag	caccgtataa	gagcagtgtc	ttggccatta	540
atttatcttc	attgtagaca	gcatagtgta	gagtgggtatt	tccatactca	tctggaatat	600
ttggatcagt	gccatgttcc	agcaacatta	acgcacattc	atcttctctg	cattgtacgg	660
cctttgtcag	agctgtcttc	tttttgttgt	caaggacatt	aagttgacat	cgtctgtcca	720
gcacgagttt	tactacttct	gaattcccat	tgccagaggc	cagatgtaga	gcagtcctct	780
tttgcttgtc	cctctgttcc	acatccgtgt	ccctgagcat	gacgatgaga	tcctttcttg	840
ggactttacc	ccaccaggca	gctctgtgga	gcttgtccag	atcttctcca	tggaagctgt	900
acctgggata	catgaaggcg	ctgtcatcgt	agtctcccca	agcgaccacg	ttgtctttgc	960
cgtctccctg	cagcagggga	agcagtggca	gcaccacttg	cacctcttgc	tccaagcgt	1020
cttcacagag	gagtcgttgt	ggtctccaga	agtgccacg	ttgtctttgc	cgtctccctt	1080
gtccatccag	ggaggaagaa	atgcaggaaa	tgaagatgc	atgcacgatg	gtatactcct	1140
cagccatcaa	acttctggac	agcaggtcac	ttccagcaag	gtggagaaag	ctgtccaccc	1200
acagaggatg	agatccagaa	accacaatat	ccattcacaa	acaaacactt	ttcagccaga	1260
cacaggtact	gaaatcatgt	catctgcggc	aacatgggtg	aacctaccca	atcacacatc	1320
aagagatgaa	gacactgcag	tatatctgca	caacgtaata	ctcttcatcc	ataacaaaat	1380
aatataattt	tcctctggag	ccatatggat	gaactatgaa	ggaagaactc	cccgaagaag	1440
ccagtcgcag	agaagccaca	ctgaagctct	gtcctcagcc	atcagcgcca	cggacaggag	1500
tgtgtttctt	ccccagtgat	gcagcctcaa	gttatcccga	agctgccgca	gcacacgggt	1560
gctcctgaga	aacaccccag	ctcttccggt	ctaacacagg	caagtcaata	aatgtgataa	1620
tcacataaac	agaattaaaa	gcaaagtcac	ataagcatct	caacagacac	agaaaaggca	1680
tttgacaaaa	tccagcatcc	ttgtatttat	tgttgacagt	ctcagaggaa	atgcttctaa	1740
cttttcccca	tttagtatta	tgttggctgt	gggcttgtca	taggtgggtt	ttattacttt	1800
aaggtatgtc	ccttctatgc	ctgttttgc	gagggtttta	attctcgtgc	c	1851

<210> 292

<211> 1851

<212> DNA

<213> Homo sapien

<400> 292
 tcatcaccat tgccagcagc ggcaccgtta gtcagggttt ctgggaatcc cacatgagta 60
 cttccgtgtt cttcattctt cttcaatagc cataaatctt ctagctctgg ctggctgttt 120
 tcaacttctt taagcctttg tgactcttcc tctgatgtca gctttaagtc ttgttctgga 180
 ttgctgtttt cagaagagat ttttaacatc tgtttttctt tgtagtcaga aagtaactgg 240
 caaattacat gatgatgact agaaacagca tactctctgg ccgtctttcc agatcttgag 300
 aagatacatc aacattttgc tcaagtagag ggctgactat acttgctgat ccacaacata 360
 cagcaagtat gagagcagtt cttccatatac tatccagcgc atttaaattc gcttttttct 420
 tgattaaaaa tttcaccact tgctgttttt gctcatgtat accaagtagc agtgggtgta 480
 ggccatgctt gttttttgat tcgatatcag caccgtataa gagcagtgtt ttggccatta 540
 atttatcttc attgtagaca gcatagtgtt gagggttatt tccatactca tctggaatat 600
 ttggatcagt gccatgttcc agcaacatta acgcacattc atcttctgg cattgtacgg 660
 cctttgtcag agctgtcttc tttttgttgt caaggacatt aagttgacat cgtctgtcca 720
 gcacgagttt tactacttct gaattcccat tggcagaggc cagatgtaga gcagtcctct 780
 tttgcttgct cctctgttcc acatccgtgt ccttgagcat gacgatgaga tcctttctgg 840
 ggactttacc ccaccaggca gctctgttga gcttgtccag atcttctcca tggacgtggt 900
 acctgggagt catgaaggcg ctgtcatcgt agtctccca agcgaccaag ttgctcttgc 960
 cgctcccctg cagcagggga agcagtgcca gcaccacttg cacctcttgc tcccaagcgt 1020
 cttcacagag gagtctgtgt ggtctccaga agtgccacg ttgctcttgc cgctcccct 1080
 gtccatccag ggaggaagaa atgcaggaaa tgaagatgc atgcacgatg gtatactcct 1140
 cagccatcaa acttctggac agcaggtcac ttccagcaag gtggagaaag ctgtccaccc 1200
 acagaggatg agatccagaa accacaatat ccattcacaa acaaacactt ttcagccaga 1260
 cacaggtact gaaatcatgt catctgcggc aacatggtgg aacctacca atcacacatc 1320
 aagagatgaa gacactgcag tatactctga caacgtaata ctcttcatcc ataacaaaat 1380
 aatataattt tcctctggag ccatatggat gaactatgaa ggaagaactc cccgaagaag 1440
 ccagtgcag agaagccaca ctgaagctct gtctcagcc atcagcgcca cggacaggar 1500
 tgtgtttctt cccagtgat gcagctcaa gttatcccga agctgccgca gcacacgggt 1560
 gctcctgaga aacaccccag ctcttccggt ctaacacagg caagtcaata aatgtgataa 1620
 tcacataaac agaattaaaa gcaaagtcac ataagcatct caacagacac agaaaaggca 1680
 tttgacaaaa tccagcatcc ttgtatttat tgttgagtt ctcagaggaa atgcttctaa 1740
 cttttcccca tttagtatta tgttggtgtt gggcttgtca taggtggtt ttattacttt 1800
 aaggtatgtc ccttctatgc ctgttttgc gagggtttta attctcgtgc c 1851

<210> 293
 <211> 668
 <212> DNA
 <213> Homo sapien

<400> 293
 cttgagcttc caaataygga agactggccc ttacacasgt caatgttaaa atgaatgcat 60
 ttcagtattt tgaagataaa attrgtagat ctataccttg ttttttgatt cgatatcagc 120
 accrtataag agcagtgtt tggtcattaa tttatcttcc attrtagaca gcrtagtgya 180
 gagtgttatt tccatactca tctggaatat ttggatcagt gccatgttcc agcaacatta 240
 acgcacattc atcttcttgg cattgtacgg cctgtcagta ttagacccaa aaacaaatta 300
 catatcttag gaattcaaaa taacattcca cagctttcac caactagtta tatttaaagg 360
 agaaaactca tttttatgcc atgtattgaa atcaaaccca cctcatgctg atatagttgg 420
 ctactgcata cttttatcag agctgtcctc tttttgttgt caaggacatt aagttgacat 480
 cgtctgtcca gcaggagttt tactacttct gaattcccat tggcagaggc cagatgtaga 540
 gcagtcctat gcaggtgaga agacttttta ggaattgta gtgcactagc tacagccata 600
 gcaatgatcc atgtaactgc aaacactgaa tagcctgcta ttactctgcc ttcaaaaaaa 660
 aaaaaaa 668

<210> 294
 <211> 1512
 <212> DNA
 <213> Homo sapien

<400> 294

gggtcgccca	gggggsgcgt	gggctttcct	cggttggttg	tgggttttcc	ctgggtgggg	60
tgggctgggc	trgaatcccc	tgctgggggt	ggcagggttt	ggctgggatt	gacttttytc	120
ttcaaacaga	ttggaacccc	ggagttacct	gctagttggt	gaaactgggt	ggtagacgcg	180
atctgttggc	tactactggc	ttctcctggc	tggtaaaagc	agatggtggt	tgaggttgat	240
tccatgccgg	ctgcttcttc	tgtgaagaag	ccatttggtc	tcaggagcaa	gatgggcaag	300
tggtgctgcc	gttgcttccc	ctgctgcagg	gagagcggca	agagcaacgt	gggcacttct	360
ggagaccacg	acgactctgc	tatgaagaca	ctcaggagca	agatgggcaa	gtggtgccgc	420
cactgcttcc	cctgctgcag	ggggagtggc	aagagcaacg	tgggcgcttc	tggagaccac	480
gacgaytctg	ctatgaagac	actcaggaac	aagatgggca	agtgggtgctg	ccactgcttc	540
ccctgctgca	gggggagcrg	caagagcaag	gtgggcgctt	ggggagacta	cgatgacagt	600
gccttcatgg	agcccaggta	ccacgtccgt	ggagaagatc	tggacaagct	ccacagagct	660
gcctgggtgg	gtaaaagccc	cagaaaggat	ctcatcgtca	tgctcaggga	cactgacgtg	720
aacaagaagg	acaagcaaaa	gaggactgct	ctacatctgg	cctctgccaa	tgggaattca	780
gaagtagtaa	aactcstgct	ggacagacga	tgtcaactta	atgtccttga	caacaaaaag	840
aggacagctc	tgayaaaggc	cgtacaatgc	caggaagatg	aatgtgcgtt	aatgttgctg	900
gaacatggca	ctgatccaaa	tattccagat	gagtatggaa	ataccactct	rcactaygct	960
rtctayaatg	aagataaatt	aatggccaaa	gcactgctct	tatayggtgc	tgatatcgaa	1020
tcaaaaaaca	aggtatagat	ctactaattt	tatcttcaaa	atactgaaat	gcattcattt	1080
taacattgac	gtgtgtaagg	gccagtcttc	cgtatttgga	agctcaagca	taacttgaat	1140
gaaaatattt	tgaatgacc	taattatctm	agactttatt	ttaaataattg	ttattttcaa	1200
agaagcatta	gagggtagac	tttttttttt	ttaaatagcac	ttctggtaaa	tacttttggt	1260
gaaaacactg	aatttgtaaa	aggtataact	tactattttt	caatttttcc	ctcctaggat	1320
ttttttcccc	taatgaatgt	aagatggcaa	aatttgccct	gaaataggtt	ttacatgaaa	1380
actccaagaa	aagttaaaca	tgtttcagtg	aatagagatc	ctgctccttt	ggcaagttcc	1440
taaaaaacag	taatagatac	gaggtgatgc	gcctgtcagt	ggcaagggtt	aagatatttc	1500
tgatctcgtg	cc					1512

<210> 295

<211> 1853

<212> DNA

<213> Homo sapien

<400> 295

gggtcgccca	gggggsgcgt	gggctttcct	cggttggttg	tgggttttcc	ctgggtgggg	60
tgggctgggc	trgaatcccc	tgctgggggt	ggcagggttt	ggctgggatt	gacttttytc	120
ttcaaacaga	ttggaacccc	ggagttacct	gctagttggt	gaaactgggt	ggtagacgcg	180
atctgttggc	tactactggc	ttctcctggc	tggtaaaagc	agatggtggt	tgaggttgat	240
tccatgccgg	ctgcttcttc	tgtgaagaag	ccatttggtc	tcaggagcaa	gatgggcaag	300
tggtgctgcc	gttgcttccc	ctgctgcagg	gagagcggca	agagcaacgt	gggcacttct	360
ggagaccacg	acgactctgc	tatgaagaca	ctcaggagca	agatgggcaa	gtggtgccgc	420
cactgcttcc	cctgctgcag	ggggagtggc	aagagcaacg	tgggcgcttc	tggagaccac	480
gacgaytctg	ctatgaagac	actcaggaac	aagatgggca	agtgggtgctg	ccactgcttc	540
ccctgctgca	gggggagcrg	caagagcaag	gtgggcgctt	ggggagacta	cgatgacagy	600
gccttcatgg	akcccaggta	ccacgtccrt	ggagaagatc	tggacaagct	ccacagagct	660
gcctgggtgg	gtaaaagccc	cagaaaggat	ctcatcgtca	tgctcaggga	cackgaygtg	720
aacaagargg	acaagcaaaa	gaggactgct	ctacatctgg	cctctgccaa	tgggaattca	780
gaagtagtaa	aactcstgct	ggacagacga	tgtcaactta	atgtccttga	caacaaaaag	840
aggacagctc	tgayaaaggc	cgtacaatgc	caggaagatg	aatgtgcgtt	aatgttgctg	900
gaacatggca	ctgatccaaa	tattccagat	gagtatggaa	ataccactct	rcactaygct	960
rtctayaatg	aagataaatt	aatggccaaa	gcactgctct	tatayggtgc	tgatatcgaa	1020
tcaaaaaaca	agcatggcct	cacaccactg	ytacttggtg	tacatgagca	aaaacagcaa	1080
gtsgtgaaat	ttttaatyaa	gaaaaaagcg	aattttaaatt	gcrctggata	gatatggaag	1140
ractgctctc	atacttgctg	tatgttggtg	atcagcaagt	atagtacgoc	ytctacttga	1200
gcaaaatrtr	gatgtatctt	ctcaagatct	ggaaagacgg	ccagagagta	tgctgtttct	1260
agtcatcatc	atgtaatttg	ccagttactt	tctgactaca	aagaaaaaca	gatgttaaaa	1320
atctcttctg	aaaacagcaa	tccagaacaa	gacttaaaagc	tgacatcaga	ggaagagtca	1380
caaaggctta	aagggaagtga	aaacagccag	ccagaggcat	ggaaactttt	aaatttaaac	1440
ttttggttta	atgttttttt	tttttgcctt	aataatatta	gatatgcccc	aatgaaatwa	1500

cctatgagac	taggctttga	gaatcaatag	attctttttt	taagaatctt	ttggctagga	1560
gcggtgtctc	acgcctgtaa	ttccagcacc	ttgagaggct	gaggtgggca	gatcacgaga	1620
tcaggagatc	gagaccatcc	tggctaacac	ggtgaaaccc	catctctact	aaaaatacaa	1680
aaacttagct	gggtgtggtg	gcgggtgcct	gtagtcccag	ctactcagga	rgctgaggca	1740
ggagaatggc	atgaaccggg	gaggtggagg	ttgcagtgag	ccgagatccg	ccactacact	1800
ccagcctggg	tgacagagca	agactctgtc	tcaaaaaaaaa	aaaaaaaaaaa	aaa	1853

<210> 296

<211> 2184

<212> DNA

<213> Homo sapien

<400> 296

ggcacgagaa	ttaaaaccct	cagcaaaaca	ggcatagaag	ggacatacct	taaagtaata	60
aaaaccacct	atgacaagcc	cacagccaac	ataatactaa	atggggaaaa	gttagaagca	120
tttcctctga	gaactgcaac	aataaataca	aggatgctgg	attttgtcaa	atgccttttc	180
tgtgtctggt	gagatgctta	tgtgactttg	cttttaattc	tgtttatgtg	attatcacat	240
ttattgactt	gcctgtgtta	gaccggaaga	gctggggtgt	ttctcaggag	ccaccgtgtg	300
ctgcgcgagc	ttcgggataa	cttgaggctg	catcactggg	gaagaaacac	aytccctgtcc	360
gtggcgctga	tggctgagga	cagagcttca	gtgtggcttc	tctgcgactg	gcttcttcgg	420
ggagtctctc	cttcatagtt	catccatatg	gctccagagg	aaaattatat	tattttgtta	480
tggatgaaga	gtattacgtt	gtgcagatat	actgcagtgt	cttcactctc	tgatgtgtga	540
ttgggtaggt	tccaccatgt	tgccgcagat	gacatgattt	cagtacctgt	gtctggctga	600
aaagtgtttg	tttgtgaatg	gataattgtg	tttctggatc	tcacccctctg	tgggtggaca	660
gctttctcca	ccttgctgga	agtgcactgc	tgtccagaag	tttgatggct	gaggagtata	720
ccatcgtgca	tgcactcttc	atttccctgca	tttcttcctc	cctggatgga	cagggggagc	780
ggcaagagca	acgtgggcac	ttctggagac	cacaacgact	cctctgtgaa	gacgcttggg	840
agcaagaggt	gcaagtgggtg	ctgccactgc	ttcccctgct	gcaggggagc	ggcaagagca	900
acgtggtcgc	ttggggagac	tacgatgaca	gcgccttcac	ggatcccagg	taccacgtcc	960
atggagaaga	tctggacaag	ctccacagag	ctgcctgggtg	gggtaaagtc	cccagaaagg	1020
atctcatcgt	catgctcagg	gacacggatg	tgaacaagag	ggacaagcaa	aagaggactg	1080
ctctacatct	ggcctctgcc	aatgggaatt	cagaagtagt	aaaactcgtg	ctggacagac	1140
gatgtcaact	taatgtcctt	gacaacaaaa	agaggacagc	tctgacaaag	gccgtacaat	1200
gccaggaaga	tgaatgtgcg	ttaatgttgc	tggaaacatg	cactgatcca	aatattccag	1260
atgagtatgg	aaataccact	ctacactatg	ctgtctacaa	tgaagataaa	ttaatggcca	1320
aagcactgct	cttatacggg	gctgatatcg	aatcaaaaaa	caagcatggc	ctcacaccac	1380
tgctacttgg	tatacatgag	caaaaacagc	aagtgtgtaa	atttttaatc	aagaaaaaag	1440
cgaatttaaa	tgcgctggat	agatatggaa	gaactgctct	catacttgct	gtatgttgtg	1500
gatcagcaag	tatagtcagc	cctctacttg	agcaaaatgt	tgatgtatct	tctcaagatc	1560
tggaaagacg	gccagagagt	atgctgtttc	tagtcatcat	catgtaattt	gccagttact	1620
ttctgactac	aaagaaaaac	agatgttaaa	aatctcttct	gaaaacagca	atccagaaca	1680
agacttaaa	ctgacatcag	aggaagagtc	acaaaggctt	aaaggaagtg	aaaacagcca	1740
gccagaggca	tggaaacttt	taaatttaaa	cttttgggtt	aatgtttttt	ttttttgcct	1800
taataaatatt	agatagtccc	aaatgaaatw	acctatgaga	ctaggctttg	agaatcaata	1860
gattcttttt	ttaagaatct	tttggctagg	agcgggtgtc	cacgcctgta	attccagcac	1920
cttgagaggc	tgaggtgggc	agatcacgag	atcaggagat	cgagaccatc	ctggctaaca	1980
cggtgaaacc	ccatctctac	taaaaataca	aaaacttagc	tgggtgtggg	ggcgggtgcc	2040
tgtagtccca	gctactcagg	argctgaggc	aggagaatgg	catgaaccgg	ggaggtggag	2100
gttgagtgga	gccgagatcc	gccactacac	tccagcctgg	gtgacagagc	aagactctgt	2160
ctcaaaaaaa	aaaaaaaaaa	aaaa				2184

<210> 297

<211> 1855

<212> DNA

<213> Homo sapien

<220>

<221> misc_feature

<222> (1)... (1855)

<223> n = A, T, C or G

<400> 297

tgacgcac	ggccagtgtc	tgtgccacgt	acactgacgc	cccctgagat	gtgcacgccg	60
cacgcgcacg	ttgcacgcgc	ggcagcggct	tggctggctt	gtaacggctt	gcacgcgcac	120
gccgcccccg	cataaccgtc	agactggcct	gtaacggctt	gcaggcgcac	gccgcacgcg	180
cgtaacggct	tggctgccct	gtaacggctt	gcacgtgcat	gctgcacgcg	cgtaacggc	240
ttggctggca	tgtagccgct	tggcttggct	ttgcattytt	tgctkggctk	ggcgttgkty	300
tcttgattg	acgcttcctc	cttggatkga	cgtttcctcc	ttggatkgac	gtttcytyty	360
tcgcgttctc	ttgctggact	tgacctttty	tctgctgggt	ttggcattcc	tttgggtgg	420
gctgggtgtt	ttctccgggg	gggkktgccc	ttcctggggg	ggcgtgggk	cgccccagg	480
ggcgtgggc	tttccccggg	tgggtgtggg	ttttcctggg	gtgggtggg	ctgtgctggg	540
atccccctgc	tgggttggc	agggattgac	tttttcttcc	aaacagattg	gaaacccgga	600
gtaacntgct	agttgtgaa	actggttgg	agacgcgcatc	tgctggtact	actgtttctc	660
ctggctgtta	aaagcagatg	gtggctgagg	ttgattcaat	gccggctgct	tcttctgtga	720
agaagccatt	tggctcagg	agcaagatgg	gcaagtggg	cgccactgct	tccccgtctg	780
cagggggagc	ggcaagagca	acgtgggcac	ttctggagac	cacaacgact	cctctgtgaa	840
gacgcttggg	agcaagaggt	gcaagtggg	ctgcccactg	cttccccctgc	tgacggggag	900
cggaagagc	aacgtggkcg	cttggggaga	ctacgatgac	agcgccttca	tggakcccag	960
gtaccacgtc	crtggagaag	atctggacaa	gctccacaga	gctgcctgg	ggggtaaagt	1020
ccccagaaag	gatctcatcg	tcatgctcag	ggacactgag	gtgaacaaga	rggacaagca	1080
aaagaggact	gctctacatc	tggcctctgc	caatgggaat	tcagaagtag	taaaactcgt	1140
gctggacaga	cgatgtcaac	ttaatgtcct	tgacaacaaa	aagaggacag	ctctgacaaa	1200
ggcgttaca	tgccaggaag	atgaatgtgc	gttaatgttg	ctggaacatg	gcactgatcc	1260
aaatattcca	gatgagtatg	gaaataccac	tctacactat	gctgtctaca	atgaagataa	1320
attaatggcc	aaagcactgc	tcttatacgg	tgctgatatac	gaatcaaaaa	acaaggtata	1380
gatctactaa	ttttatcttc	aaaatactga	aatgcattca	ttttaacatt	gacgtgtgta	1440
agggccagtc	ttccgtattt	ggaagctcaa	gcataacttg	aatgaaaata	ttttgaaatg	1500
acctaattat	ctaagacttt	attttaaata	ttgttatttt	caaagaagca	ttagagggtta	1560
cagttttttt	tttttaaattg	cacttctggt	aaatactttt	gttgaaaaca	ctgaatttgt	1620
aaaaggtaat	acttactatt	tttcaatttt	tccctcctag	gatttttttc	ccctaattgaa	1680
tgtgaagatg	ctgaatttgc	cctgaaatag	gttttaccatg	aaaactccaa	gaaaagttaa	1740
acatgtttca	gtgaatagag	atcctgctcc	tttggaaggt	tcctaaaaaa	cagtaataga	1800
tacgaggtga	tgccgctgtc	agtggcaagg	tttaagatat	ttctgatctc	gtgcc	1855

<210> 298

<211> 1059

<212> DNA

<213> Homo sapien

<400> 298

gcaacgtggg	cacttctgga	gaccacaacg	actcctctgt	gaagacgctt	gggagcaaga	60
ggtgcaagtg	gtgctgcccc	ctgcttcccc	tgctgcaggg	gagcggcaag	agcaacgtgg	120
gcgcttgrrg	agactmcgat	gacagygcct	tcatggagcc	caggtaccac	gtccgtggag	180
aagatctgga	caagctccac	agagctgccc	tgggtgggta	aagtccccag	aaaggatctc	240
atcgtcatgc	tcaggagacac	tgaygtgaac	aagarggaca	agcaaaaagag	gactgctcta	300
catctggcct	ctgccaatgg	gaattcagaa	gtagtataaac	tctgctgga	cagacgatgt	360
caacttaatg	tccttgacaa	caaaaagagg	acagctctga	yaaaggccgt	acaatgccag	420
gaagatgaat	gtgcgttaat	gttgcctgaa	catggcactg	atccaaatat	tccagatgag	480
tatggaaata	ccactctrca	ctaygctrct	tayaatgaag	ataaattaat	ggccaaagca	540
ctgctcttat	ayggtgctga	tatcgaatca	aaaaacaagg	tatagatcta	ctaattttat	600
cttcaaaaata	ctgaaaatgca	ttcattttta	cattgacgtg	tgtaagggcc	agtcttccgt	660
atlttgaagc	tcaagcataa	cttgaatgaa	aatattttga	aatgacctaa	ttatctaaga	720
ctttattttta	aatattgtta	ttttcaaaga	agcattagag	ggtacagttt	ttttttttta	780
aatgcacttc	tggtaaatat	ttttgttgaa	aacactgaat	ttgtaaaaag	taatacttac	840
tatttttcaa	tttttccctc	ctaggatttt	tttcccctaa	tgaatgtaag	atggcaaaat	900
ttgccttgaa	ataggtttta	catgaaaact	ccaagaaaag	ttaaacatgt	ttcagtgaat	960

agagatcctg ctcctttggc aagttcctaa aaaacagtaa tagatacgag gtgatgcgcc 1020
 tgcagtgcc aaggtttaag atatttctga tctcgtgcc 1059

<210> 299

<211> 329

<212> PRT

<213> Homo sapien

<400> 299

Met	Asp	Ile	Val	Val	Ser	Gly	Ser	His	Pro	Leu	Trp	Val	Asp	Ser	Phe
1				5					10					15	
Leu	His	Leu	Ala	Gly	Ser	Asp	Leu	Leu	Ser	Arg	Ser	Leu	Met	Ala	Glu
			20					25					30		
Glu	Tyr	Thr	Ile	Val	His	Ala	Ser	Phe	Ile	Ser	Cys	Ile	Ser	Ser	Ser
		35					40					45			
Leu	Asp	Gly	Gln	Gly	Glu	Arg	Gln	Glu	Gln	Arg	Gly	His	Phe	Trp	Arg
	50					55					60				
Pro	Gln	Arg	Leu	Leu	Cys	Glu	Asp	Ala	Trp	Glu	Gln	Glu	Val	Gln	Val
65					70					75				80	
Val	Leu	Pro	Leu	Leu	Pro	Leu	Leu	Gln	Gly	Ser	Gly	Lys	Ser	Asn	Val
			85						90					95	
Val	Ala	Trp	Gly	Asp	Tyr	Asp	Asp	Ser	Ala	Phe	Met	Asp	Pro	Arg	Tyr
			100					105					110		
His	Val	His	Gly	Glu	Asp	Leu	Asp	Lys	Leu	His	Arg	Ala	Ala	Trp	Trp
		115					120					125			
Gly	Lys	Val	Pro	Arg	Lys	Asp	Leu	Ile	Val	Met	Leu	Arg	Asp	Thr	Asp
	130					135					140				
Val	Asn	Lys	Arg	Asp	Lys	Gln	Lys	Arg	Thr	Ala	Leu	His	Leu	Ala	Ser
145					150					155					160
Ala	Asn	Gly	Asn	Ser	Glu	Val	Val	Lys	Leu	Val	Leu	Asp	Arg	Arg	Cys
			165						170					175	
Gln	Leu	Asn	Val	Leu	Asp	Asn	Lys	Lys	Arg	Thr	Ala	Leu	Thr	Lys	Ala
			180					185					190		
Val	Gln	Cys	Gln	Glu	Asp	Glu	Cys	Ala	Leu	Met	Leu	Leu	Glu	His	Gly
		195					200				205				
Thr	Asp	Pro	Asn	Ile	Pro	Asp	Glu	Tyr	Gly	Asn	Thr	Thr	Leu	His	Tyr
	210					215					220				
Ala	Val	Tyr	Asn	Glu	Asp	Lys	Leu	Met	Ala	Lys	Ala	Leu	Leu	Leu	Tyr
225					230					235					240
Gly	Ala	Asp	Ile	Glu	Ser	Lys	Asn	Lys	His	Gly	Leu	Thr	Pro	Leu	Leu
			245						250					255	
Leu	Gly	Ile	His	Glu	Gln	Lys	Gln	Gln	Val	Val	Lys	Phe	Leu	Ile	Lys
			260					265					270		
Lys	Lys	Ala	Asn	Leu	Asn	Ala	Leu	Asp	Arg	Tyr	Gly	Arg	Thr	Ala	Leu
			275				280					285			
Ile	Leu	Ala	Val	Cys	Cys	Gly	Ser	Ala	Ser	Ile	Val	Ser	Pro	Leu	Leu
	290					295					300				
Glu	Gln	Asn	Val	Asp	Val	Ser	Ser	Gln	Asp	Leu	Glu	Arg	Arg	Pro	Glu
305					310					315					320
Ser	Met	Leu	Phe	Leu	Val	Ile	Ile	Met							
					325										

<210> 300

<211> 148

<212> PRT

<213> Homo sapien

<220>

<221> VARIANT

<222> (1)...(148)

<223> Xaa = Any Amino Acid

<400> 300

```

Met Thr Xaa Pro Ser Trp Ser Pro Gly Thr Thr Ser Val Glu Lys Ile
 1          5          10          15
Trp Thr Ser Ser Thr Glu Leu Pro Trp Trp Gly Lys Val Pro Arg Lys
      20          25          30
Asp Leu Ile Val Met Leu Arg Asp Thr Asp Val Asn Lys Xaa Asp Lys
      35          40          45
Gln Lys Arg Thr Ala Leu His Leu Ala Ser Ala Asn Gly Asn Ser Glu
      50          55          60
Val Val Lys Leu Xaa Leu Asp Arg Arg Cys Gln Leu Asn Val Leu Asp
      65          70          75          80
Asn Lys Lys Arg Thr Ala Leu Xaa Lys Ala Val Gln Cys Gln Glu Asp
      85          90          95
Glu Cys Ala Leu Met Leu Leu Glu His Gly Thr Asp Pro Asn Ile Pro
      100          105          110
Asp Glu Tyr Gly Asn Thr Thr Leu His Tyr Ala Xaa Tyr Asn Glu Asp
      115          120          125
Lys Leu Met Ala Lys Ala Leu Leu Tyr Gly Ala Asp Ile Glu Ser
      130          135          140
Lys Asn Lys Val
145

```

<210> 301

<211> 1155

<212> DNA

<213> Homo sapien

<400> 301

```

atggttggttg aggttgattc catgccggct gcctcttctg tgaagaagcc atttgggtctc      60
aggagcaaga tgggcaagtg gtgctgcogt tgcttcccct gctgcaggga gagcggcaag      120
agcaactgtg gcacttcttg agaccacgac gactctgcta tgaagacact caggagcaag      180
atgggcaagt ggtgccgcca ctgcttcccc tgctgcaggg ggagtggcaa gagcaactgt      240
ggcgcttctg gagaccacga cgactctgct atgaagacac tcaggaacaa gatgggcaag      300
tggtgctgcc actgcttccc ctgctgcagg gggagcggca agagcaaggt gggcgcttgg      360
ggagactacg atgacagtgc cttcatggag ccaggtacc acgtccgtgg agaagatctg      420
gacaagctcc acagagctgc ctggtggggt aaagtcccca gaaaggatct catcgatcatg      480
ctcagggaca ctgacgtgaa caagaaggac aagcaaaaga ggactgctct acatctggcc      540
tctgccaatg ggaattcaga agtagtaaaa ctctgctggt acagacgatg tcaacttaat      600
gtccttgaca acaaaaagag gacagctctg ataaaggccg tacaatgcca ggaagatgaa      660
tgtgcgttaa tgttgctgga acatggcact gatccaaata ttccagatga gtatggaaat      720
accactctgc actacgctat ctataatgaa gataaattaa tggccaaagc actgctctta      780
tatggtgctg atatcgaatc aaaaaacaag catggcctca caccactgtt acttgggtgta      840
catgagcaaa aacagcaagt cgtgaaatth ttaatcaaga aaaaagcgaa tttaaatgca      900
ctggatagat atggaaggac tgctctcata cttgctgtat gttgtggatc agcaagtata      960
gtcagccttc tacttgagca aaatattgat gtatcttctc aagatctatc tggacagacg      1020
gccagagagt tagctgtttc tagtcatcat catgtaattt gccagttact ttctgactac      1080
aaagaaaaac agatgctaaa aatctcttct gaaaacagca atccagaaaa tgtctcaaga      1140
accagaaata aataa
1155

```

<210> 302

<211> 2000

<212> DNA

<213> Homo sapien

<400> 302

atggtggttg	aggttgattc	catgccggct	gcctcttctg	tgaagaagcc	atttggctctc	60
aggagcaaga	tgggcaagt	gtgctgccgt	tgcttcccct	gctgcaggga	gagcggcaag	120
agcaacgtgg	gcacttctgg	agaccacgac	gactctgcta	tgaagacact	caggagcaag	180
atgggcaagt	ggtgccgcca	ctgcttcccc	tgctgcagg	ggagtggcaa	gagcaacgtg	240
ggcgcttctg	gagaccacga	cgactctgct	atgaagacac	tcaggaacaa	gatgggcaag	300
tggtgctgcc	actgcttccc	ctgctgcagg	gggagcggca	agagcaaggt	gggcgcttgg	360
ggagactacg	atgacagtgc	cttcatggag	cccaggtacc	acgtccgtgg	agaagatctg	420
gacaagctcc	acagagctgc	ctggtgggg	aaagtcccca	gaaaggatct	catcgtcattg	480
ctcagggaca	ctgacgtgaa	caagaaggac	aagcaaaaaga	ggactgctct	acatctggcc	540
tctgccaatg	ggaattcaga	agtagtaaaa	ctcctgctgg	acagacgatg	tcaacttaat	600
gtccttgaca	acaaaaagag	gacagctctg	ataaaggccg	tacaatgcca	ggaagatgaa	660
tggtgcgttaa	tggtgctgga	acatggcact	gatccaaata	ttccagatga	gtatggaaat	720
accactctgc	actacgctat	ctataatgaa	gataaattaa	tggccaaagc	actgctctta	780
tatggtgctg	atatcgaaatc	aaaaaacaag	catggcctca	caccactgtt	acttgggtgta	840
catgagcaaaa	aacagcaagt	cgtgaaattt	ttaatcaaga	aaaaagcgaa	tttaaatgca	900
ctggatagat	atggaaggac	tgctctcata	cttgcgtgat	gttgtggatc	agcaagtata	960
gtcagccttc	tacttgagca	aaatattgat	gtatcttctc	aagatctatc	tggacagacg	1020
gccagagagt	atgctgtttc	tagtcatcat	catgtaattt	gccagttact	ttctgactac	1080
aaagaaaaac	agatgctaaa	aatctcttct	gaaaacagca	atccagaaca	agacttaaaag	1140
ctgacatcag	aggaagagtc	acaaagggtc	aaaggcagtg	aaaatagcca	gccagagaaa	1200
atgtctcaag	aaccagaaat	aaataaggat	ggtgatagag	aggttgaaga	agaaatgaag	1260
aagcatgaaa	gtaataatgt	gggattacta	gaaaacctga	ctaattggtgt	cactgctggc	1320
aatggtgata	atggattaat	tcctcaaagg	aagagcagaa	cacctgaaaa	tcagcaattt	1380
cctgacaacg	aaagtgaaga	gtatcacaga	atttgcgaat	tagtttctga	ctacaaagaa	1440
aaacagatgc	caaaatactc	ttctgaaaac	agcaacccag	aacaagactt	aaagtgcaca	1500
tcagaggaag	agtcacaaaag	gcttgagggc	agtgaataatg	gccagccaga	gctagaaaat	1560
tttatggcta	tcgaagaaat	gaagaagcac	ggaagtactc	atgtcggatt	cccagaaaac	1620
ctgactaatg	gtgccactgc	tggcaatggt	gatgatggat	taattcctcc	aaggaagagc	1680
agaacacctg	aaagccagca	atttccctgac	actgagaatg	aagagtatca	cagtgcagaa	1740
caaaatgata	ctcagaagca	attttgtgaa	gaacagaaca	ctggaatatt	acacgatgag	1800
attctgattc	atgaagaaaa	gcagatagaa	gtggttgaaa	aaatgaattc	tgagctttct	1860
cttagttgta	agaaagaaaa	agacatcttg	catgaaaata	gtacgttgcg	ggaagaaatt	1920
gccatgctaa	gactggagct	agacacaatg	aaacatcaga	gccagctaaa	aaaaaaaaaa	1980
aaaaaaaaaa	aaaaaaaaaa					2000

<210> 303

<211> 2040

<212> DNA

<213> Homo sapien

<400> 303

atggtggttg	aggttgattc	catgccggct	gcctcttctg	tgaagaagcc	atttggctctc	60
aggagcaaga	tgggcaagt	gtgctgccgt	tgcttcccct	gctgcaggga	gagcggcaag	120
agcaacgtgg	gcacttctgg	agaccacgac	gactctgcta	tgaagacact	caggagcaag	180
atgggcaagt	ggtgccgcca	ctgcttcccc	tgctgcagg	ggagtggcaa	gagcaacgtg	240
ggcgcttctg	gagaccacga	cgactctgct	atgaagacac	tcaggaacaa	gatgggcaag	300
tggtgctgcc	actgcttccc	ctgctgcagg	gggagcggca	agagcaaggt	gggcgcttgg	360
ggagactacg	atgacagtgc	cttcatggag	cccaggtacc	acgtccgtgg	agaagatctg	420
gacaagctcc	acagagctgc	ctggtgggg	aaagtcccca	gaaaggatct	catcgtcattg	480
ctcagggaca	ctgacgtgaa	caagaaggac	aagcaaaaaga	ggactgctct	acatctggcc	540
tctgccaatg	ggaattcaga	agtagtaaaa	ctcctgctgg	acagacgatg	tcaacttaat	600
gtccttgaca	acaaaaagag	gacagctctg	ataaaggccg	tacaatgcca	ggaagatgaa	660
tggtgcgttaa	tggtgctgga	acatggcact	gatccaaata	ttccagatga	gtatggaaat	720
accactctgc	actacgctat	ctataatgaa	gataaattaa	tggccaaagc	actgctctta	780
tatggtgctg	atatcgaaatc	aaaaaacaag	catggcctca	caccactgtt	acttgggtgta	840
catgagcaaaa	aacagcaagt	cgtgaaattt	ttaatcaaga	aaaaagcgaa	tttaaatgca	900
ctggatagat	atggaaggac	tgctctcata	cttgcgtgat	gttgtggatc	agcaagtata	960

```

gtcagccttc tacttgagca aaatattgat gtatcttctc aagatctatc tggacagacg 1020
gccagagagt atgctgtttc tagtcatcat catgtaattt gccagttact ttctgactac 1080
aaagaaaaaac agatgctaaa aatctcttct gaaaacagca atccagaaca agacttaaag 1140
ctgacatcag aggaagagtc acaaaggttc aaaggcagtg aaaatagcca gccagagaaa 1200
atgtctcaag aaccagaaat aaataaggat ggtgatagag aggttgaaga agaatgaag 1260
aagcatgaaa gtaataatgt gggattacta gaaaacctga ctaatggtgt cactgctggc 1320
aatggtgata atggattaat tcctcaaagg aagagcagaa cacctgaaaa tcagcaattt 1380
cctgacaacg aaagtgaaga gtatcacaga atttgcgaa tagtttctga ctacaaagaa 1440
aaacagatgc caaaatactc ttctgaaaac agcaaccag aacaagactt aaagctgaca 1500
tcagaaggaa agtcacaaag gcttgagggc agtgaaaatg gccagccaga gaaaagatct 1560
caagaaccag aaataaataa ggatggtgat agagagctag aaaattttat ggctatcgaa 1620
gaaatgaaga agcacggaag tactcatgtc ggattcccag aaaacctgac taatggtgcc 1680
actgctggca atggtgatga tggattaatt cctccaagga agagcagaac acctgaaagc 1740
cagcaatttc ctgacactga gaatgaagag tatcacagt acgaacaaaa tgatactcag 1800
aagcaatttt gtgaagaaca gaacactgga atattacacg atgagattct gattcatgaa 1860
gaaaagcaga tagaagtggg tgaaaaaatg aattctgagc tttctcttag ttgtaagaaa 1920
gaaaagaca tcttgcataa aaatagtagc ttgcgggaag aaattgccat gctaagactg 1980
gagctagaca caatgaaaca tcagagccag ctaaaaaaaa aaaaaaaaaa aaaaaaaaaa 2040

```

<210> 304

<211> 384

<212> PRT

<213> Homo sapien

<400> 304

```

Met Val Val Glu Val Asp Ser Met Pro Ala Ala Ser Ser Val Lys Lys
1      5      10      15
Pro Phe Gly Leu Arg Ser Lys Met Gly Lys Trp Cys Cys Arg Cys Phe
20     25     30
Pro Cys Cys Arg Glu Ser Gly Lys Ser Asn Val Gly Thr Ser Gly Asp
35     40     45
His Asp Asp Ser Ala Met Lys Thr Leu Arg Ser Lys Met Gly Lys Trp
50     55     60
Cys Arg His Cys Phe Pro Cys Cys Arg Gly Ser Gly Lys Ser Asn Val
65     70     75     80
Gly Ala Ser Gly Asp His Asp Asp Ser Ala Met Lys Thr Leu Arg Asn
85     90     95
Lys Met Gly Lys Trp Cys Cys His Cys Phe Pro Cys Cys Arg Gly Ser
100    105    110
Gly Lys Ser Lys Val Gly Ala Trp Gly Asp Tyr Asp Asp Ser Ala Phe
115    120    125
Met Glu Pro Arg Tyr His Val Arg Gly Glu Asp Leu Asp Lys Leu His
130    135    140
Arg Ala Ala Trp Trp Gly Lys Val Pro Arg Lys Asp Leu Ile Val Met
145    150    155    160
Leu Arg Asp Thr Asp Val Asn Lys Lys Asp Lys Gln Lys Arg Thr Ala
165    170    175
Leu His Leu Ala Ser Ala Asn Gly Asn Ser Glu Val Val Lys Leu Leu
180    185    190
Leu Asp Arg Arg Cys Gln Leu Asn Val Leu Asp Asn Lys Lys Arg Thr
195    200    205
Ala Leu Ile Lys Ala Val Gln Cys Gln Glu Asp Glu Cys Ala Leu Met
210    215    220
Leu Leu Glu His Gly Thr Asp Pro Asn Ile Pro Asp Glu Tyr Gly Asn
225    230    235    240
Thr Thr Leu His Tyr Ala Ile Tyr Asn Glu Asp Lys Leu Met Ala Lys
245    250    255
Ala Leu Leu Leu Tyr Gly Ala Asp Ile Glu Ser Lys Asn Lys His Gly

```

```

                260                265                270
Leu Thr Pro Leu Leu Leu Gly Val His Glu Gln Lys Gln Gln Val Val
                275                280                285
Lys Phe Leu Ile Lys Lys Lys Ala Asn Leu Asn Ala Leu Asp Arg Tyr
                290                295                300
Gly Arg Thr Ala Leu Ile Leu Ala Val Cys Cys Gly Ser Ala Ser Ile
305                310                315                320
Val Ser Leu Leu Leu Glu Gln Asn Ile Asp Val Ser Ser Gln Asp Leu
                325                330                335
Ser Gly Gln Thr Ala Arg Glu Tyr Ala Val Ser Ser His His His Val
                340                345                350
Ile Cys Gln Leu Leu Ser Asp Tyr Lys Glu Lys Gln Met Leu Lys Ile
                355                360                365
Ser Ser Glu Asn Ser Asn Pro Glu Asn Val Ser Arg Thr Arg Asn Lys
370                375                380

```

```

<210> 305
<211> 656
<212> PRT
<213> Homo sapien

```

```

<400> 305
Met Val Val Glu Val Asp Ser Met Pro Ala Ala Ser Ser Val Lys Lys
1          5          10          15
Pro Phe Gly Leu Arg Ser Lys Met Gly Lys Trp Cys Cys Arg Cys Phe
20          25          30
Pro Cys Cys Arg Glu Ser Gly Lys Ser Asn Val Gly Thr Ser Gly Asp
35          40          45
His Asp Asp Ser Ala Met Lys Thr Leu Arg Ser Lys Met Gly Lys Trp
50          55          60
Cys Arg His Cys Phe Pro Cys Cys Arg Gly Ser Gly Lys Ser Asn Val
65          70          75          80
Gly Ala Ser Gly Asp His Asp Asp Ser Ala Met Lys Thr Leu Arg Asn
85          90          95
Lys Met Gly Lys Trp Cys Cys His Cys Phe Pro Cys Cys Arg Gly Ser
100         105         110
Gly Lys Ser Lys Val Gly Ala Trp Gly Asp Tyr Asp Asp Ser Ala Phe
115         120         125
Met Glu Pro Arg Tyr His Val Arg Gly Glu Asp Leu Asp Lys Leu His
130         135         140
Arg Ala Ala Trp Trp Gly Lys Val Pro Arg Lys Asp Leu Ile Val Met
145         150         155         160
Leu Arg Asp Thr Asp Val Asn Lys Lys Asp Lys Gln Lys Arg Thr Ala
165         170         175
Leu His Leu Ala Ser Ala Asn Gly Asn Ser Glu Val Val Lys Leu Leu
180         185         190
Leu Asp Arg Arg Cys Gln Leu Asn Val Leu Asp Asn Lys Lys Arg Thr
195         200         205
Ala Leu Ile Lys Ala Val Gln Cys Gln Glu Asp Glu Cys Ala Leu Met
210         215         220
Leu Leu Glu His Gly Thr Asp Pro Asn Ile Pro Asp Glu Tyr Gly Asn
225         230         235         240
Thr Thr Leu His Tyr Ala Ile Tyr Asn Glu Asp Lys Leu Met Ala Lys
245         250         255
Ala Leu Leu Leu Tyr Gly Ala Asp Ile Glu Ser Lys Asn Lys His Gly
260         265         270
Leu Thr Pro Leu Leu Leu Gly Val His Glu Gln Lys Gln Gln Val Val
275         280         285

```

Lys Phe Leu Ile Lys Lys Lys Ala Asn Leu Asn Ala Leu Asp Arg Tyr
 290 295 300
 Gly Arg Thr Ala Leu Ile Leu Ala Val Cys Cys Gly Ser Ala Ser Ile
 305 310 315 320
 Val Ser Leu Leu Leu Glu Gln Asn Ile Asp Val Ser Ser Gln Asp Leu
 325 330 335
 Ser Gly Gln Thr Ala Arg Glu Tyr Ala Val Ser Ser His His Val
 340 345 350
 Ile Cys Gln Leu Leu Ser Asp Tyr Lys Glu Lys Gln Met Leu Lys Ile
 355 360 365
 Ser Ser Glu Asn Ser Asn Pro Glu Gln Asp Leu Lys Leu Thr Ser Glu
 370 375 380
 Glu Glu Ser Gln Arg Phe Lys Gly Ser Glu Asn Ser Gln Pro Glu Lys
 385 390 395 400
 Met Ser Gln Glu Pro Glu Ile Asn Lys Asp Gly Asp Arg Glu Val Glu
 405 410 415
 Glu Glu Met Lys Lys His Glu Ser Asn Asn Val Gly Leu Leu Glu Asn
 420 425 430
 Leu Thr Asn Gly Val Thr Ala Gly Asn Gly Asp Asn Gly Leu Ile Pro
 435 440 445
 Gln Arg Lys Ser Arg Thr Pro Glu Asn Gln Gln Phe Pro Asp Asn Glu
 450 455 460
 Ser Glu Glu Tyr His Arg Ile Cys Glu Leu Val Ser Asp Tyr Lys Glu
 465 470 475 480
 Lys Gln Met Pro Lys Tyr Ser Ser Glu Asn Ser Asn Pro Glu Gln Asp
 485 490 495
 Leu Lys Leu Thr Ser Glu Glu Glu Ser Gln Arg Leu Glu Gly Ser Glu
 500 505 510
 Asn Gly Gln Pro Glu Leu Glu Asn Phe Met Ala Ile Glu Glu Met Lys
 515 520 525
 Lys His Gly Ser Thr His Val Gly Phe Pro Glu Asn Leu Thr Asn Gly
 530 535 540
 Ala Thr Ala Gly Asn Gly Asp Asp Gly Leu Ile Pro Pro Arg Lys Ser
 545 550 555 560
 Arg Thr Pro Glu Ser Gln Gln Phe Pro Asp Thr Glu Asn Glu Glu Tyr
 565 570 575
 His Ser Asp Glu Gln Asn Asp Thr Gln Lys Gln Phe Cys Glu Glu Gln
 580 585 590
 Asn Thr Gly Ile Leu His Asp Glu Ile Leu Ile His Glu Glu Lys Gln
 595 600 605
 Ile Glu Val Val Glu Lys Met Asn Ser Glu Leu Ser Leu Ser Cys Lys
 610 615 620
 Lys Glu Lys Asp Ile Leu His Glu Asn Ser Thr Leu Arg Glu Glu Ile
 625 630 635 640
 Ala Met Leu Arg Leu Glu Leu Asp Thr Met Lys His Gln Ser Gln Leu
 645 650 655

<210> 306

<211> 671

<212> PRT

<213> Homo sapien

<400> 306

Met Val Val Glu Val Asp Ser Met Pro Ala Ala Ser Ser Val Lys Lys
 1 5 10 15
 Pro Phe Gly Leu Arg Ser Lys Met Gly Lys Trp Cys Cys Arg Cys Phe
 20 25 30
 Pro Cys Cys Arg Glu Ser Gly Lys Ser Asn Val Gly Thr Ser Gly Asp

		35						40						45					
His	Asp	Asp	Ser	Ala	Met	Lys	Thr	Leu	Arg	Ser	Lys	Met	Gly	Lys	Trp				
Cys	50	Arg	His	Cys	Phe	Pro	Cys	Arg	Gly	Ser	60	Gly	Lys	Ser	Val				
65					70					75					80				
Gly	Ala	Ser	Gly	Asp	His	Asp	Asp	Ser	Ala	Met	Lys	Thr	Leu	Arg	Asn				
				85					90					95					
Lys	Met	Gly	Lys	Trp	Cys	Cys	His	Cys	Phe	Pro	Cys	Cys	Arg	Gly	Ser				
			100					105					110						
Gly	Lys	Ser	Lys	Val	Gly	Ala	Trp	Gly	Asp	Tyr	Asp	Asp	Ser	Ala	Phe				
			115				120					125							
Met	Glu	Pro	Arg	Tyr	His	Val	Arg	Gly	Glu	Asp	Leu	Asp	Lys	Leu	His				
	130					135					140								
Arg	Ala	Ala	Trp	Trp	Gly	Lys	Val	Pro	Arg	Lys	Asp	Leu	Ile	Val	Met				
145					150					155					160				
Leu	Arg	Asp	Thr	Asp	Val	Asn	Lys	Lys	Asp	Lys	Gln	Lys	Arg	Thr	Ala				
				165					170					175					
Leu	His	Leu	Ala	Ser	Ala	Asn	Gly	Asn	Ser	Glu	Val	Val	Lys	Leu	Leu				
			180					185					190						
Leu	Asp	Arg	Arg	Cys	Gln	Leu	Asn	Val	Leu	Asp	Asn	Lys	Lys	Arg	Thr				
		195					200					205							
Ala	Leu	Ile	Lys	Ala	Val	Gln	Cys	Gln	Glu	Asp	Glu	Cys	Ala	Leu	Met				
	210					215					220								
Leu	Leu	Glu	His	Gly	Thr	Asp	Pro	Asn	Ile	Pro	Asp	Glu	Tyr	Gly	Asn				
225					230					235					240				
Thr	Thr	Leu	His	Tyr	Ala	Ile	Tyr	Asn	Glu	Asp	Lys	Leu	Met	Ala	Lys				
				245					250					255					
Ala	Leu	Leu	Leu	Tyr	Gly	Ala	Asp	Ile	Glu	Ser	Lys	Asn	Lys	His	Gly				
			260				265						270						
Leu	Thr	Pro	Leu	Leu	Leu	Gly	Val	His	Glu	Gln	Lys	Gln	Gln	Val	Val				
		275					280					285							
Lys	Phe	Leu	Ile	Lys	Lys	Lys	Ala	Asn	Leu	Asn	Ala	Leu	Asp	Arg	Tyr				
	290				295						300								
Gly	Arg	Thr	Ala	Leu	Ile	Leu	Ala	Val	Cys	Cys	Gly	Ser	Ala	Ser	Ile				
305					310					315					320				
Val	Ser	Leu	Leu	Leu	Glu	Gln	Asn	Ile	Asp	Val	Ser	Ser	Gln	Asp	Leu				
				325					330					335					
Ser	Gly	Gln	Thr	Ala	Arg	Glu	Tyr	Ala	Val	Ser	Ser	His	His	His	Val				
			340					345					350						
Ile	Cys	Gln	Leu	Leu	Ser	Asp	Tyr	Lys	Glu	Lys	Gln	Met	Leu	Lys	Ile				
		355					360					365							
Ser	Ser	Glu	Asn	Ser	Asn	Pro	Glu	Gln	Asp	Leu	Lys	Leu	Thr	Ser	Glu				
		370				375					380								
Glu	Glu	Ser	Gln	Arg	Phe	Lys	Gly	Ser	Glu	Asn	Ser	Gln	Pro	Glu	Lys				
385					390					395					400				
Met	Ser	Gln	Glu	Pro	Glu	Ile													

Asn	Gly	Gln	Pro	Glu	Lys	Arg	Ser	Gln	Glu	Pro	Glu	Ile	Asn	Lys	Asp
515							520						525		
Gly	Asp	Arg	Glu	Leu	Glu	Asn	Phe	Met	Ala	Ile	Glu	Glu	Met	Lys	Lys
530						535						540			
His	Gly	Ser	Thr	His	Val	Gly	Phe	Pro	Glu	Asn	Leu	Thr	Asn	Gly	Ala
545					550					555					560
Thr	Ala	Gly	Asn	Gly	Asp	Asp	Gly	Leu	Ile	Pro	Pro	Arg	Lys	Ser	Arg
				565					570					575	
Thr	Pro	Glu	Ser	Gln	Gln	Phe	Pro	Asp	Thr	Glu	Asn	Glu	Glu	Tyr	His
			580					585					590		
Ser	Asp	Glu	Gln	Asn	Asp	Thr	Gln	Lys	Gln	Phe	Cys	Glu	Glu	Gln	Asn
		595					600					605			
Thr	Gly	Ile	Leu	His	Asp	Glu	Ile	Leu	Ile	His	Glu	Glu	Lys	Gln	Ile
610						615						620			
Glu	Val	Val	Glu	Lys	Met	Asn	Ser	Glu	Leu	Ser	Leu	Ser	Cys	Lys	Lys
625					630					635					640
Glu	Lys	Asp	Ile	Leu	His	Glu	Asn	Ser	Thr	Leu	Arg	Glu	Glu	Ile	Ala
				645					650					655	
Met	Leu	Arg	Leu	Glu	Leu	Asp	Thr	Met	Lys	His	Gln	Ser	Gln	Leu	
			660					665					670		

<210> 307
 <211> 800
 <212> DNA
 <213> Homo sapien

<400> 307	
atkagcttcc gcttctgaca acactagaga tccctcccct ccctcagggg atggccctcc	60
acttcatttt tggtagataa catctttata ggacaggggt aaaatcccaa tactaacagg	120
agaatgctta ggactctaac aggtttttga gaatgtgttg gtaagggcca ctcaatccaa	180
tttttcttgg tccctcctgt ggtctaggag gacaggcaag ggtgcagatt ttcaagaatg	240
catcagtaag gggcactaaa tccgaccttc ctctgttctc cttgtgtgtc gggaggaaaa	300
ctagtgttct tgttgctgtg tcagttagca caactattcc gatcagcagg gtccaggagc	360
cactgcagggt tcttgggcag ggggagaaac aaaacaaacc aaaaccatgg gcrgttttgt	420
ctttcagatg ggaacactc aggcatacaac aggcacacct ttgaaatgca tcctaagcca	480
atggggacaaa ttgacccac aaaccctgga aaaagagggt gctcattttt ttgacactat	540
ggcttgccc caacattctc tctctgatgg ggaataatgg ccacctgagg gaagtacaga	600
ttacaatact atcctgcagc ttgacctttt ctgtaagagg gaaggcaaat ggagtgaat	660
accttatgtc caagctttct tttcattgaa ggagaataca ctatgcaaag cttgaaattt	720
acatcccaca ggaggacctc tcagcttacc cccatatcct agcctcccta tagctccct	780
tcctattagt gataagcctc	800

<210> 308
 <211> 102
 <212> PRT
 <213> Homo sapien

<220>
 <221> VARIANT
 <222> (1)...(102)
 <223> Xaa = Any Amino Acid

<400> 308	
Met Gly Xaa Phe Val Phe Gln Met Gly Asn Thr Gln Ala Ser Thr Gly	
1 5 10 15	
Ser Pro Leu Lys Cys Ile Leu Ser Gln Trp Asp Lys Phe Asp Pro Gln	
20 25 30	

95

Thr Leu Glu Lys Glu Val Ala His Phe Phe Cys Thr Met Ala Trp Pro
 35 40 45
 Gln His Ser Leu Ser Asp Gly Glu Lys Trp Pro Pro Glu Gly Ser Thr
 50 55 60
 Asp Tyr Asn Thr Ile Leu Gln Leu Asp Leu Phe Cys Lys Arg Glu Gly
 65 70 75 80
 Lys Trp Ser Glu Ile Pro Tyr Val Gln Ala Phe Phe Ser Leu Lys Glu
 85 90 95
 Asn Thr Leu Cys Lys Ala
 100

<210> 309
 <211> 9
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Made in the lab

<400> 309
 Leu Met Ala Glu Glu Tyr Thr Ile Val
 1 5

<210> 310
 <211> 9
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Made in the lab

<400> 310
 Lys Leu Met Ala Lys Ala Leu Leu Leu
 1 5

<210> 311
 <211> 9
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Made in the lab

<400> 311
 Gly Leu Thr Pro Leu Leu Gly Ile
 1 5

<210> 312
 <211> 10
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Made in the lab

<400> 312
 Lys Leu Val Leu Asp Arg Arg Cys Gln Leu
 1 5 10

<210> 313
 <211> 1852
 <212> DNA
 <213> Homo sapiens

<400> 313
 ggcacgagaa ttaaaaccct cagcaaaaaca ggcatagaag ggacatacct taaagtaata 60
 aaaaccacct atgacaagcc cacagccaac ataatactaa atggggaaaa gttagaagca 120
 tttcctctga gaactgcaac aataaatata aggatgctgg attttgtcaa atgccttttc 180
 tgtgtctgtt gagatgctta tgtgactttg cttttaattc tgtttatgtg attatcacat 240
 ttattgactt gcctgtgtta gaccggaaga gctgggggtg ttctcaggag ccaccgtgtg 300
 ctgcggcagc ttcgggataa cttgaggctg catcactggg gaagaaacac aytccctgtcc 360
 gtggcgctga tggctgagga cagagcttca gtgtggcttc tctgcgactg gcttcttcgg 420
 ggagtctctc cttcatagtt catccatatg gctccagagg aaaattatat tattttgtta 480
 tggatgaaga gtattacgtt gtgcagatat actgcagtgt cttcatctct tgatgtgtga 540
 ttgggtaggt tccaccatgt tgccgcagat gacatgattt cagtacctgt gtctggctga 600
 aaagtgtttt tttgtgaatg gatattgtgg tttctggatc tcatcctctg tgggtggaca 660
 gctttctcca ccttgcctga agtgacctgc tgtccagaag tttgatggct gaggagtata 720
 ccacgtgca tgcacttttc atttcctgca tttcttctc cctggatgga cagggggagc 780
 ggcaagagca acgtgggcac ttctggagac cacaacgact cctctgtgaa gacgcttggg 840
 agcaagaggt gcaagtgggt ctgccactgc ttcccctgct gcagggggag cggcaagagc 900
 aacgtggctc cttggggaga ctacgatgac agcgcttca tggatcccag gtaccacgtc 960
 catggagaag atctggacaa gctccacaga gctgcctggg ggggtaaagt ccccagaaag 1020
 gatctcatcg tcatgctcag ggacacggat gtgaacaaga gggacaagca aaagaggact 1080
 gctctacatc tggcctctgc caatgggaat tcagaagtag taaaactcgt gctggacaga 1140
 cgatgtcaac ttaatgtcct tgacaacaaa aagaggacag ctctgacaaa ggccgtacaa 1200
 tgccaggaag atgaatgtgc gttaatgttg ctggaacatg gcaactgatc aaatattcca 1260
 gatgagtagt gaaataccac tctacactat gctgtctaca atgaagataa attaatggcc 1320
 aaagcactgc tcttatacgg tgctgatatc gaatcaaaaa acaagcatgg cctcacacca 1380
 ctgctacttg gtatacatga gcaaaaacag caagtgggtg aattttttaa caagaaaaaa 1440
 gcgaatttaa atgcgctgga tagatatgga agaactgctc tcatacttgc tgtatgttgt 1500
 ggatcagcaa gtatagtcag ccctctactt gagcaaaatg ttgatgtatc ttctcaagat 1560
 ctggaaagac ggccagagag tatgctgttt ctagtcatca tcatgtaatt tgccagttac 1620
 tttctgacta caaagaaaaa cagatgttaa aaatctcttc tgaaaacagc aatccagAAC 1680
 aagacttaaa gctgacatca gaggaagagt cacaaggct taaaggaaat gaaaacagcc 1740
 agccagagct agaagattta tggctattga agaagaatga agaacacgga agtactcatg 1800
 tgggattccc agaaaacctg actaacggtg ccgctgctgg caatggtgat ga 1852

<210> 314
 <211> 879
 <212> DNA
 <213> Homo sapiens

<400> 314
 atgcactctt catttcttgc atttcttcct ccctggatgg acagggggag cggcaagagc 60
 aacgtgggca cttctggaga ccacaacgac tcctctgtga agacgcttgg gagcaagagg 120
 tgcaagtggg gctgccactg cttcccctgc tgcaggggga gcggcaagag caacgtgggtc 180
 gcttggggag actacgatga cagcgcttc atggatcca ggtaccacgt ccatggagaa 240
 gatctggaca agctccacag agctgcctgg tggggtaaag tcccagaaa ggatctcatc 300
 gtcagtctca gggacacgga tgtgaacaag agggacaagc aaaagaggac tgctctacat 360
 ctggcctctg ccaatgggaa ttcagaagta gtaaaactcg tgctggacag acgatgtcaa 420
 cttaatgtcc ttgacaacaa aaagaggaca gctctgacaa aggccgtaca atgccaggaa 480
 gatgaatgtg cgttaatgtt gctggaacat ggcactgatc caaatattcc agatgagat 540
 ggaataacca ctctacacta tgctgtctac aatgaagata aattaatggc caaagcactg 600
 ctcttatacg gtgctgatat cgaatcaaaa aacaagcatg gcctcacacc actgctactt 660
 ggtatacatg agcaaaaaca gcaagtgtgt aaatttttaa tcaagaaaaa agcgaattta 720
 aatgcgctgg atagatatgg aagaactgct ctcatacttg ctgtatgttg tggatcagca 780

```

agtatagtca gccctctact tgagcaaaat gttgatgtat cttctcaaga tctggaaaga 840
cggccagaga gtatgctgtt tctagtcatc atcatgtaa                               879

```

<210> 315

<211> 292

<212> PRT

<213> Homo sapiens

<400> 315

Met His Leu Ser Phe Pro Ala Phe Leu Pro Pro Trp Met Asp Arg Gly
5 10 15

Ser Gly Lys Ser Asn Val Gly Thr Ser Gly Asp His Asn Asp Ser Ser
20 25 30

Val Lys Thr Leu Gly Ser Lys Arg Cys Lys Trp Cys Cys His Cys Phe
35 40 45

Pro Cys Cys Arg Gly Ser Gly Lys Ser Asn Val Val Ala Trp Gly Asp
50 55 60

Tyr Asp Asp Ser Ala Phe Met Asp Pro Arg Tyr His Val His Gly Glu
65 70 75 80

Asp Leu Asp Lys Leu His Arg Ala Ala Trp Trp Gly Lys Val Pro Arg
85 90 95

Lys Asp Leu Ile Val Met Leu Arg Asp Thr Asp Val Asn Lys Arg Asp
100 105 110

Lys Gln Lys Arg Thr Ala Leu His Leu Ala Ser Ala Asn Gly Asn Ser
115 120 125

Glu Val Val Lys Leu Val Leu Asp Arg Arg Cys Gln Leu Asn Val Leu
130 135 140

Asp Asn Lys Lys Arg Thr Ala Leu Thr Lys Ala Val Gln Cys Gln Glu
145 150 155 160

Asp Glu Cys Ala Leu Met Leu Leu Glu His Gly Thr Asp Pro Asn Ile
165 170 175

Pro Asp Glu Tyr Gly Asn Thr Thr Leu His Tyr Ala Val Tyr Asn Glu
180 185 190

Asp Lys Leu Met Ala Lys Ala Leu Leu Leu Tyr Gly Ala Asp Ile Glu
195 200 205

Ser Lys Asn Lys His Gly Leu Thr Pro Leu Leu Leu Gly Ile His Glu
210 215 220

Gln Lys Gln Gln Val Val Lys Phe Leu Ile Lys Lys Lys Ala Asn Leu
225 230 235 240

Asn Ala Leu Asp Arg Tyr Gly Arg Thr Ala Leu Ile Leu Ala Val Cys
245 250 255

Cys Gly Ser Ala Ser Ile Val Ser Pro Leu Leu Glu Gln Asn Val Asp

260 265 270

Val Ser Ser Gln Asp Leu Glu Arg Arg Pro Glu Ser Met Leu Phe Leu
275 280 285

Val Ile Ile Met
290

<210> 316
<211> 584
<212> DNA
<213> Homo sapiens

<400> 316
agttgggcca aattcccctc cccctacagc ttgaagggga cataaccaat agcctgggggt 60
ttttttgtgg tccttttgag atttctttgc ttattttctt ctgggtgggg gtgattagag 120
gaggcttatt actaatagga aggggagcta tagggaggct aggatatggg ggtaagctga 180
gaggctctcc tgtgggatgt aaatttcaag ctttgcatag tgtattctcc ttcaatgaaa 240
agaaagcttg gacataaggt atttcaactc atttgccttc cctcttacag aaaagggtcaa 300
gctgcaggat agtattgtaa tctgtacttc cctcaggtag ccatttttcc ccatcagaga 360
gagaatgttg gggccaagcc atagtgcaga aaaaaaatg agccacctct ttttccaggg 420
tttgtgggtc aaatttgtcc cattggctta ggatgcattt caaagggtgag cctgttgatg 480
cctgagtgtt tcccatctga aagacaaaac tgcccatggt tttggtttgt tttgtttctc 540
ccctgcccc agaactatca aactcctgag ccaacaacta aaaa 584

<210> 317
<211> 829
<212> DNA
<213> Homo sapiens

<400> 317
attagcttcc gcttctgaca acactagaga tccctcccct ccctcagggt atggccctcc 60
acttcatttt tggtagataa catctttata ggacaggggt aaaatcccaa tactaacagg 120
agaatgctta ggactctaac aggtttttga gaatgtgttg gtaagggccca ctcaatccaa 180
tttttcttgg tctccttgt ggtctaggag gacaggcaag ggtgcagatt ttcaagaatg 240
catcagtaag ggccactaaa tccgaccttc ctctgtctct cttgtgtgtc gggaggaaaa 300
ctagtgtttc tgttgctgtg tcagtgcagca caactattcc gatcagcagg gtccaggggac 360
cactgcagggt tcttgggcag ggggagaaac aaaacaaacc aaaaccatgg gcagttttgt 420
ctttcagatg ggaaacactc aggcataaac aggcacacct ttgaaatgca tcctaagcca 480
atgggacaaa tttagccac aaaccctgga aaaagagggt gctcattttt tttagcactat 540
ggcttggccc caacattctc tctctgatgg ggaaaaatgg ccacctgagg gaagtacaga 600
ttacaatact atcctgcagc ttgacctttt ctgtaagagg gaaggcaaat ggagtgaat 660
accttatgtc caagctttct ttctattgaa ggagaataca ctatgcaaag cttgaaattt 720
acatcccaca ggaggacctc tcagettacc cccatatact agcctcccta tagctccct 780
tcctattagt gataagcctc ctctaatac cccacccag aagaaaata 829

<210> 318
<211> 30
<212> PRT
<213> Homo sapien

<400> 318
Thr Ala Ala Ser Asp Asn Phe Gln Leu Ser Gln Gly Gly Gln Gly Phe
1 5 10 15
Ala Ile Pro Ile Gly Gln Ala Met Ala Ile Ala Gly Gln Ile
20 25 30

<210> 319
<211> 41
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR primer

<400> 319

ggcctctgcc aatgggaact cagaagtagt aaaactcctg c 41

<210> 320
<211> 41
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR primer

<400> 320

gcaggagttt tactacttct gagttcccat tggcagaggc c 41

<210> 321
<211> 60
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR primer

<400> 321

ggggaattcc cgctggtgcc gcgcggcagc cctatggtgg ttgaggttga 50
ttccatgccg 60

<210> 322
<211> 42
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR primer

<400> 322

cccgaattct tatttatttc tggttcttga gacattttct gg 42

<210> 323
<211> 1590
<212> DNA
<213> Homo sapiens

```

<400> 323
atgcatcacc atcaccatca cacggccgcg tccgataact tccagctgtc ccaggggtggg 60
cagggattcg ccattccgat cgggcaggcg atggcgatcg cgggccagat caagcttccc 120
accgttcata tcgggcctac cgccttcctc ggcttgggtg ttgtcgacaa caacggcaac 180
ggcgacagag tccaacgcgt ggtcgggagc gctccggcgg caagtctcgg catctccacc 240
ggcgacgtga tcaccgcggt cgacggcgct ccgatcaact cggccaccgc gatggcggac 300
gcgcttaacg ggcatcatcc cgggtgacgtc atctcgggtg cctggcaaac caagtcgggc 360
ggcacgcgta cagggaacgt gacattggcc gagggacccc cggccgaatt cccgctggtg 420
ccgcgcggca gccctatggt ggttgagggt gattccatgc cggctgcttc ttctgtgaag 480
aagccatttg gtctcaggag caagatgggc aagtgggtgct gccgttgett cccctgctgc 540
agggagagcg gcaagagcaa cgtgggcact tctggagacc acgacgactc tgctatgaag 600
acactcagga gcaagatggg caagtgggtc cgccactgct tcccctgctg cagggggagt 660
ggcaagagca acgtgggcgc ttctggagac cacgacgact ctgctatgaa gacactcagg 720
aacaagatgg gcaagtgggt ctgccactgc ttcccctgct gcagggggag cggcaagagc 780
aaggtgggcg cttggggaga ctacgatgac agygccttca tggagcccag gtaccacgtc 840
cgtggagaag atctggacaa gctccacaga gctgcctggt ggggtaaagt cccagaaaag 900
gatctcatcg tcatgctcag ggacactgac gtgaacaaga aggacaagca aaagaggagt 960
gctctacatc tggcctctgc caatgggaat tcagaagtag taaaactcct gctggacaga 1020
cgatgtcaac ttaatgtcct tgacaacaaa aagaggacag ctctgataaa ggccgtacaa 1080
tgccagggaag atgaatgtgc gttaatgttg ctggaacatg gcactgatcc aaatattcca 1140
gatgagtatg gaaataccac tctgcactac gctatctata atgaagataa attaatggcc 1200
aaagcactgc tcttatatgg tgctgatatc gaatcaaaaa acaagcatgg cctcacacca 1260
ctgttacttg gtgtacatga gcaaaaaacag caagtcgtga aatttttaat caagaaaaaa 1320
gcgaatttaa atgcactgga tagatatgga aggactgctc tcatacttgc tgtatgttgt 1380
ggatcagcaa gtatagtcag ccttctactt gagcaaaaata ttgatgtatc ttctcaagat 1440
ctatctggac agacggccag agagtatgct gtttctagtc atcatcatgt aatttgccag 1500
ttactttctg actacaaaga aaaacagatg ctaaaaatct cttctgaaaa cagcaatcca 1560
gaaaatgtct caagaaccag aaataaataa 1590

```

```

<210> 324
<211> 529
<212> PRT
<213> Homo sapiens

```

```

<400> 324
Met His His His His His Thr Ala Ala Ser Asp Asn Phe Gln Leu
      5                      10                      15

Ser Gln Gly Gly Gln Gly Phe Ala Ile Pro Ile Gly Gln Ala Met Ala
      20                      25                      30

Ile Ala Gly Gln Ile Lys Leu Pro Thr Val His Ile Gly Pro Thr Ala
      35                      40                      45

Phe Leu Gly Leu Gly Val Val Asp Asn Asn Gly Asn Gly Ala Arg Val
      50                      55                      60

Gln Arg Val Val Gly Ser Ala Pro Ala Ala Ser Leu Gly Ile Ser Thr
      65                      70                      75                      80

Gly Asp Val Ile Thr Ala Val Asp Gly Ala Pro Ile Asn Ser Ala Thr
      85                      90                      95

Ala Met Ala Asp Ala Leu Asn Gly His His Pro Gly Asp Val Ile Ser
      100                      105                      110

```

101

Val Thr Trp Gln Thr Lys Ser Gly Gly Thr Arg Thr Gly Asn Val Thr
 115 120 125
 Leu Ala Glu Gly Pro Pro Ala Glu Phe Pro Leu Val Pro Arg Gly Ser
 130 135 140
 Pro Met Val Val Glu Val Asp Ser Met Pro Ala Ala Ser Ser Val Lys
 145 150 155 160
 Lys Pro Phe Gly Leu Arg Ser Lys Met Gly Lys Trp Cys Cys Arg Cys
 165 170 175
 Phe Pro Cys Cys Arg Glu Ser Gly Lys Ser Asn Val Gly Thr Ser Gly
 180 185 190
 Asp His Asp Asp Ser Ala Met Lys Thr Leu Arg Ser Lys Met Gly Lys
 195 200 205
 Trp Cys Arg His Cys Phe Pro Cys Cys Arg Gly Ser Gly Lys Ser Asn
 210 215 220
 Val Gly Ala Ser Gly Asp His Asp Asp Ser Ala Met Lys Thr Leu Arg
 225 230 235 240
 Asn Lys Met Gly Lys Trp Cys Cys His Cys Phe Pro Cys Cys Arg Gly
 245 250 255
 Ser Gly Lys Ser Lys Val Gly Ala Trp Gly Asp Tyr Asp Asp Ser Ala
 260 265 270
 Phe Met Glu Pro Arg Tyr His Val Arg Gly Glu Asp Leu Asp Lys Leu
 275 280 285
 His Arg Ala Ala Trp Trp Gly Lys Val Pro Arg Lys Asp Leu Ile Val
 290 295 300
 Met Leu Arg Asp Thr Asp Val Asn Lys Lys Asp Lys Gln Lys Arg Thr
 305 310 315 320
 Ala Leu His Leu Ala Ser Ala Asn Gly Asn Ser Glu Val Val Lys Leu
 325 330 335
 Leu Leu Asp Arg Arg Cys Gln Leu Asn Val Leu Asp Asn Lys Lys Arg
 340 345 350
 Thr Ala Leu Ile Lys Ala Val Gln Cys Gln Glu Asp Glu Cys Ala Leu
 355 360 365
 Met Leu Leu Glu His Gly Thr Asp Pro Asn Ile Pro Asp Glu Tyr Gly
 370 375 380
 Asn Thr Thr Leu His Tyr Ala Ile Tyr Asn Glu Asp Lys Leu Met Ala
 385 390 395 400
 Lys Ala Leu Leu Leu Tyr Gly Ala Asp Ile Glu Ser Lys Asn Lys His
 405 410 415
 Gly Leu Thr Pro Leu Leu Leu Gly Val His Glu Gln Lys Gln Gln Val

102

420	425	430
Val Lys Phe Leu Ile Lys Lys Lys Ala Asn Leu Asn Ala Leu Asp Arg		
435	440	445
Tyr Gly Arg Thr Ala Leu Ile Leu Ala Val Cys Cys Gly Ser Ala Ser		
450	455	460
Ile Val Ser Leu Leu Leu Glu Gln Asn Ile Asp Val Ser Ser Gln Asp		
465	470	475
Leu Ser Gly Gln Thr Ala Arg Glu Tyr Ala Val Ser Ser His His His		
485	490	495
Val Ile Cys Gln Leu Leu Ser Asp Tyr Lys Glu Lys Gln Met Leu Lys		
500	505	510
Ile Ser Ser Glu Asn Ser Asn Pro Glu Asn Val Ser Arg Thr Arg Asn		
515	520	525

Lys

<210> 325

<211> 1155

<212> DNA

<213> Homo sapiens

<400> 325

```

atggtggctg aggtttgttc aatgcccact gcctctactg tgaagaagcc atttgatctc 60
aggagcaaga tgggcaagtg gtgccaccac cgcttcccct gctgcagggg gagcggcaag 120
agcaacatgg gcacttcttg agaccacgac gactccttta tgaagatgct caggagcaag 180
atgggcaagt gttgccgcca ctgcttcccc tgcctgcagg ggagcggcac gagcaacgtg 240
ggcacttctg gagaccatga aaactccttt atgaagatgc tcaggagcaa gatgggcaag 300
tggtgctgtc actgcttccc ctgctgcagg gggagcggca agagcaacgt gggcgcttgg 360
ggagactacg accacagcgc cttcatggag ccgaggtacc acatccgtcg agaagatctg 420
gacaagctcc acagagctgc ctggtggggt aaagtcccca gaaaggatct catcgctcatg 480
ctcagggaca ctgacatgaa caagagggac aaggaaaaga ggactgctct acatttggcc 540
tctgccaatg gaaattcaga agtagtacaa ctctgcttgg acagacgatg tcaacttaat 600
gtccttgaca acaaaaaaag gacagctctg ataaaggcca tacaatgcca ggaagatgaa 660
tgtgtgttaa tgttgcgtga acatggcgct gatcgaaata ttccagatga gtatggaaat 720
accgctctac actatgctat ctacaatgaa gataaattaa tggccaaagc actgctctta 780
tatggtgctg atattgaatc aaaaaacaag gttggcctca caccactttt gcttggcgta 840
catgaacaaa aacagcaagt ggtgaaatth ttaatcaaga aaaaagctaa tttaaatagt 900
cttgatagat atggaaggac tgcctcataa cttgctgtat gttgtggatc agcaagtata 960
gtcaatcttc tacttgagca aaatgttgat gtatcttctc aagatctatc tggacagacg 1020
gccagagagt atgctgtttc tagtcatcat catgtaattt gtgaattact ttctgactat 1080
aaagaaaaac agatgctaaa aatctcttct gaaaacagca atccagaaaa tgtctcaaga 1140
accagaaata aataa

```

<210> 326

<211> 384

<212> PRT

<213> Homo sapiens

<400> 326

Met	Val	Ala	Glu	Val	Cys	Ser	Met	Pro	Thr	Ala	Ser	Thr	Val	Lys	Lys
				5						10				15	

103

Pro Phe Asp Leu Arg Ser Lys Met Gly Lys Trp Cys His His Arg Phe
 20 25 30
 Pro Cys Cys Arg Gly Ser Gly Lys Ser Asn Met Gly Thr Ser Gly Asp
 35 40 45
 His Asp Asp Ser Phe Met Lys Met Leu Arg Ser Lys Met Gly Lys Cys
 50 55 60
 Cys Arg His Cys Phe Pro Cys Cys Arg Gly Ser Gly Thr Ser Asn Val
 65 70 75 80
 Gly Thr Ser Gly Asp His Glu Asn Ser Phe Met Lys Met Leu Arg Ser
 85 90 95
 Lys Met Gly Lys Trp Cys Cys His Cys Phe Pro Cys Cys Arg Gly Ser
 100 105 110
 Gly Lys Ser Asn Val Gly Ala Trp Gly Asp Tyr Asp His Ser Ala Phe
 115 120 125
 Met Glu Pro Arg Tyr His Ile Arg Arg Glu Asp Leu Asp Lys Leu His
 130 135 140
 Arg Ala Ala Trp Trp Gly Lys Val Pro Arg Lys Asp Leu Ile Val Met
 145 150 155 160
 Leu Arg Asp Thr Asp Met Asn Lys Arg Asp Lys Glu Lys Arg Thr Ala
 165 170 175
 Leu His Leu Ala Ser Ala Asn Gly Asn Ser Glu Val Val Gln Leu Leu
 180 185 190
 Leu Asp Arg Arg Cys Gln Leu Asn Val Leu Asp Asn Lys Lys Arg Thr
 195 200 205
 Ala Leu Ile Lys Ala Ile Gln Cys Gln Glu Asp Glu Cys Val Leu Met
 210 215 220
 Leu Leu Glu His Gly Ala Asp Arg Asn Ile Pro Asp Glu Tyr Gly Asn
 225 230 235 240
 Thr Ala Leu His Tyr Ala Ile Tyr Asn Glu Asp Lys Leu Met Ala Lys
 245 250 255
 Ala Leu Leu Leu Tyr Gly Ala Asp Ile Glu Ser Lys Asn Lys Val Gly
 260 265 270
 Leu Thr Pro Leu Leu Leu Gly Val His Glu Gln Lys Gln Gln Val Val
 275 280 285
 Lys Phe Leu Ile Lys Lys Lys Ala Asn Leu Asn Val Leu Asp Arg Tyr
 290 295 300
 Gly Arg Thr Ala Leu Ile Leu Ala Val Cys Cys Gly Ser Ala Ser Ile
 305 310 315 320

```
<210> 327
<211> 634
<212> DNA
<213> Homo sapiens
```

<400> 327							
gactgctcta	catctggcct	ctgccaatgg	aaattcagaa	gtagtaaaac	tctgtctgga	60	
cagacgatgt	caacttaata	tccttgacaa	caaaaagagg	acagctctga	caaaggccgt	120	
acaatgccag	gaagatgaat	gtgcgttaat	gttgctggaa	catggcactg	atccgaatat	180	
tccagatgag	tatggaatac	ccgctctaca	ctatgctatc	tacaatgaag	ataaaattaat	240	
ggccaaagca	ctgtctttat	acgggtgctg	tatcgaatca	aaaaacaagc	atggcctcac	300	
accactgtta	cttggtgtac	atgagcaaaa	acagcaagtg	gtgaaatttt	taatcaagaa	360	
aaaagcaaat	ttaaatgcac	tggatagata	tggaagaact	gctctcatat	ttgctgtatg	420	
ttgtggatcg	gcaagtatag	tcagccttct	acttgagcaa	aacattgatg	tatcttctca	480	
agatctatct	ggacagacgg	ccagagagta	tgctgtttct	agtcgtcata	atgtaatttg	540	
ccagttactt	tctgactaca	aagaaaaacta	gatactaaaa	gtctcttctg	aaaacagcaa	600	
tccaggaaat	gtctcaagaa	ccagaaataa	ataa			63	

```
<210> 328
<211> 1155
<212> DNA
<213> Homo sapiens
```

<400>	328						
atggttggttg	aggttgattc	catgccggct	gcctcttctg	tgaagaagcc	atttggtctc	60	
aggagcaaga	tgggcaagtg	gtgctgccgt	tgcttccct	gctgcaggga	gagcggcaag	120	
agcaacgttg	gcactcttgc	agaccacgac	gactctgcta	tgaagacact	caggagcaag	180	
atgggcaagt	ggtgcgcgca	ctgcttcccc	tgtgcaggg	ggagtggcaa	gagcaacgtg	240	
ggcgcttctg	gagaccacga	cgactctgct	atgaagacac	tacggaacaa	gatgggcaag	300	
tggtgctgcc	actgcttccc	ctgctgcagg	gggagcagca	agagcaaggt	gggcgcttgg	360	
ggagactacg	atgacagtgc	cttcatggag	cccaggtacc	acgtccgtgg	agaagatctg	420	
gacaagctcc	acagagctgc	ctggtggggg	aaagtcctcca	gaaaggatct	catcgtcatg	480	
ctcagggcaca	ctgacgtgaa	caagcaggac	aagcaaaaga	ggactgctct	acatctggcc	540	
tctgccaatg	ggaattcaga	agtagtaaaa	ctctcgctgg	acagacgatg	tcaacttaat	600	
gtccttgaca	acaaaaagag	gacagctctg	ataaaggccg	tacaatgcca	ggaagatgaa	660	
tgtgctgttaa	tgttgctgga	acatggcact	gatccaaata	ttccagatga	gtatggaaat	720	
accactctgc	actacgctat	ctataatgaa	gataaattaa	tggccaaagc	actgctctta	780	
tatggtgctg	atatcgaatt	aaaaaacgaag	catggcctca	caccactgtt	acttgggtga	840	
catgagcaaa	aacagcaagt	cgtaaaaattt	ttaattaaaga	aaaagcgaa	tttaaatgca	900	
ctggatagat	atggaaggac	tgctctcata	cttgctgtat	gttggtggatc	agcaagtata	960	
gtcagccttc	tacttgagca	aaatattgat	gtatcttctc	aagatctatc	tggaacagacg	1020	
gccagagagt	atgctgtttc	tagtcatcat	catgtaattt	gccagttact	ttctgactac	1080	
aaagaaaaac	agatgctaaa	aatctcttct	gaaaacagca	atccagaaaa	tgtctcaaga	1140	
accagaaata	aataa					1155	

<210> 329

<211> 1155
 <212> DNA
 <213> Homo sapiens

<400> 329
 atggtggtg aggtttgttc aatgcccgt gctctgctg tgaagaagcc atttgatctc 60
 aggagcaaga tgggcaagt gtgccaccac cgcttcccct gctgcagggg gagcggcaag 120
 agcaacatgg gcacttcttg agaccacgac gactccttta tgaagacgct caggagcaag 180
 atgggcaagt gttgccacca ctgcttcccc tgctgcaggg ggagcggcac gagcaatgtg 240
 ggcacttctg gagaccatga caactccttt atgaagacac tcaggagcaa gatgggcaag 300
 tgggtgctgc actgcttccc ctgctgcagg gggagcggca agagcaacgt gggcacttgg 360
 ggagactacg acgacagcgc cttcatggag ccgaggtacc acgtccgtcg agaagatctg 420
 gacaagctcc acagagctgc ctggtggggg aaagtcccca gaaaggatct catcgctcatg 480
 ctcagggaca ctgacatgaa caagagggac aagcaaaaga ggactgctct acatttggcc 540
 tctgccaatg gaaattcaga agtagtacia ctctgcttg acagacgatg tcaacttaac 600
 gtcccttgaca acaaaaaaag gacagctctg ataaaggccg tacaatgcca ggaagatgaa 660
 tgtgtgttaa tgttctgga acatggcgct gatggaaata ttcaagatga gtatggaaat 720
 accgctctac actatgctat ctacaatgaa gataaattaa tggccaaagc actgctctta 780
 tatggtgctg atattgaatc aaaaaacaag tgtggcctca caccactttt gcttggcgta 840
 catgaacaaa aacagcaagt ggtgaaattt ttaatcaaga aaaaagctaa tttaaatgca 900
 cttgatagat atggaagaac tgccctcata cttgctgtat gttgtggatc agcaagtata 960
 gtcaatcttc tacttgagca aaatgttgat gtatcttctc aagatctatc tggacagacg 1020
 gccagagagt atgctgtttc tagtcatcat catgtaattt gtgaattact ttctgactat 1080
 aaagaaaaac agatgctaaa aatctcttct gaaaacagca atccagaaaa tgtctcaaga 1140
 accagaaata aataa 1155

<210> 330
 <211> 1155
 <212> DNA
 <213> Homo sapiens

<400> 330
 atggtggctg aggtttgttc aatgcccact gctctactg tgaagaagcc atttgatctc 60
 aggagcaaga tgggcaagt gtgccaccac cgcttcccct gctgcagggg gagcggcaag 120
 agcaacatgg gcacttcttg agaccacgac gactccttta tgaagatgct caggagcaag 180
 atgggcaagt gttgccacca ctgcttcccc tgctgcaggg ggagcggcac gagcaacgtg 240
 ggcacttctg gagaccatga aaactccttt atgaagatgc tcaggagcaa gatgggcaag 300
 tgggtgctgc actgcttccc ctgctgcagg gggagcggca agagcaacgt gggcgcttgg 360
 ggagactacg accacagcgc cttcatggag ccgaggtacc acatccgtcg agaagatctg 420
 gacaagctcc acagagctgc ctggtggggg aaagtcccca gaaaggatct catcgctcatg 480
 ctcagggaca ctgacatgaa caagagggac aaggaaaaga ggactgctct acatttggcc 540
 tctgccaatg gaaattcaga agtagtacia ctctgcttg acagacgatg tcaacttaac 600
 gtcccttgaca acaaaaaaag gacagctctg ataaaggcca tacaatgcca ggaagatgaa 660
 tgtgtgttaa tgttctgga acatggcgct gatcgaaata ttccagatga gtatggaaat 720
 accgctctac actatgctat ctacaatgaa gataaattaa tggccaaagc actgctctta 780
 tatggtgctg atattgaatc aaaaaacaag tgtggcctca caccactttt gcttggcgta 840
 catgaacaaa aacagcaagt ggtgaaattt ttaatcaaga aaaaagctaa tttaaatgta 900
 cttgatagat atggaagaac tgccctcata cttgctgtat gttgtggatc agcaagtata 960
 gtcaatcttc tacttgagca aaatgttgat gtatcttctc aagatctatc tggacagacg 1020
 gccagagagt atgctgtttc tagtcatcat catgtaattt gtgaattact ttctgactat 1080
 aaagaaaaac agatgctaaa aatctcttct gaaaacagca atccagaaaa tgtctcaaga 1140
 accagaaata aataa 1155

<210> 331
 <211> 210
 <212> PRT
 <213> Homo sapiens

106

<400> 331

Thr Ala Leu His Leu Ala Ser Ala Asn Gly Asn Ser Glu Val Val Lys
 5 10 15
 Leu Leu Leu Asp Arg Arg Cys Gln Leu Asn Ile Leu Asp Asn Lys Lys
 20 25 30
 Arg Thr Ala Leu Thr Lys Ala Val Gln Cys Gln Glu Asp Glu Cys Ala
 35 40 45
 Leu Met Leu Leu Glu His Gly Thr Asp Pro Asn Ile Pro Asp Glu Tyr
 50 55 60
 Gly Asn Thr Ala Leu His Tyr Ala Ile Tyr Asn Glu Asp Lys Leu Met
 65 70 75 80
 Ala Lys Ala Leu Leu Leu Tyr Gly Ala Asp Ile Glu Ser Lys Asn Lys
 85 90 95
 His Gly Leu Thr Pro Leu Leu Leu Gly Val His Glu Gln Lys Gln Gln
 100 105 110
 Val Val Lys Phe Leu Ile Lys Lys Lys Ala Asn Leu Asn Ala Leu Asp
 115 120 125
 Arg Tyr Gly Arg Thr Ala Leu Ile Leu Ala Val Cys Cys Gly Ser Ala
 130 135 140
 Ser Ile Val Ser Leu Leu Leu Glu Gln Asn Ile Asp Val Ser Ser Gln
 145 150 155 160
 Asp Leu Ser Gly Gln Thr Ala Arg Glu Tyr Ala Val Ser Ser Arg His
 165 170 175
 Asn Val Ile Cys Gln Leu Leu Ser Asp Tyr Lys Glu Lys Gln Ile Leu
 180 185 190
 Lys Val Ser Ser Glu Asn Ser Asn Pro Gly Asn Val Ser Arg Thr Arg
 195 200 205
 Asn Lys
 210

<210> 332

<211> 384

<212> PRT

<213> Homo sapiens

<400> 332

Met Val Ala Glu Val Cys Ser Met Pro Thr Ala Ser Thr Val Lys Lys
 5 10 15
 Pro Phe Asp Leu Arg Ser Lys Met Gly Lys Trp Cys His His Arg Phe
 20 25 30
 Pro Cys Cys Arg Gly Ser Gly Lys Ser Asn Met Gly Thr Ser Gly Asp
 35 40 45

107

His Asp Asp Ser Phe Met Lys Met Leu Arg Ser Lys Met Gly Lys Cys
 50 55 60
 Cys Arg His Cys Phe Pro Cys Cys Arg Gly Ser Gly Thr Ser Asn Val
 65 70 75 80
 Gly Thr Ser Gly Asp His Glu Asn Ser Phe Met Lys Met Leu Arg Ser
 85 90 95
 Lys Met Gly Lys Trp Cys Cys His Cys Phe Pro Cys Cys Arg Gly Ser
 100 105 110
 Gly Lys Ser Asn Val Gly Ala Trp Gly Asp Tyr Asp His Ser Ala Phe
 115 120 125
 Met Glu Pro Arg Tyr His Ile Arg Arg Glu Asp Leu Asp Lys Leu His
 130 135 140
 Arg Ala Ala Trp Trp Gly Lys Val Pro Arg Lys Asp Leu Ile Val Met
 145 150 155 160
 Leu Arg Asp Thr Asp Met Asn Lys Arg Asp Lys Glu Lys Arg Thr Ala
 165 170 175
 Leu His Leu Ala Ser Ala Asn Gly Asn Ser Glu Val Val Gln Leu Leu
 180 185 190
 Leu Asp Arg Arg Cys Gln Leu Asn Val Leu Asp Asn Lys Lys Arg Thr
 195 200 205
 Ala Leu Ile Lys Ala Ile Gln Cys Gln Glu Asp Glu Cys Val Leu Met
 210 215 220
 Leu Leu Glu His Gly Ala Asp Arg Asn Ile Pro Asp Glu Tyr Gly Asn
 225 230 235 240
 Thr Ala Leu His Tyr Ala Ile Tyr Asn Glu Asp Lys Leu Met Ala Lys
 245 250 255
 Ala Leu Leu Leu Tyr Gly Ala Asp Ile Glu Ser Lys Asn Lys Cys Gly
 260 265 270
 Leu Thr Pro Leu Leu Leu Gly Val His Glu Gln Lys Gln Gln Val Val
 275 280 285
 Lys Phe Leu Ile Lys Lys Lys Ala Asn Leu Asn Val Leu Asp Arg Tyr
 290 295 300
 Gly Arg Thr Ala Leu Ile Leu Ala Val Cys Cys Gly Ser Ala Ser Ile
 305 310 315 320
 Val Asn Leu Leu Leu Glu Gln Asn Val Asp Val Ser Ser Gln Asp Leu
 325 330 335
 Ser Gly Gln Thr Ala Arg Glu Tyr Ala Val Ser Ser His His His Val
 340 345 350

108

Ile Cys Glu Leu Leu Ser Asp Tyr Lys Glu Lys Gln Met Leu Lys Ile
 355 360 365

Ser Ser Glu Asn Ser Asn Pro Glu Asn Val Ser Arg Thr Arg Asn Lys
 370 375 380

<210> 333

<211> 384

<212> PRT

<213> Homo sapiens

<400> 333

Met Val Ala Glu Val Cys Ser Met Pro Ala Ala Ser Ala Val Lys Lys
 5 10 15

Pro Phe Asp Leu Arg Ser Lys Met Gly Lys Trp Cys His His Arg Phe
 20 25 30

Pro Cys Cys Arg Gly Ser Gly Lys Ser Asn Met Gly Thr Ser Gly Asp
 35 40 45

His Asp Asp Ser Phe Met Lys Thr Leu Arg Ser Lys Met Gly Lys Cys
 50 55 60

Cys His His Cys Phe Pro Cys Cys Arg Gly Ser Gly Thr Ser Asn Val
 65 70 75 80

Gly Thr Ser Gly Asp His Asp Asn Ser Phe Met Lys Thr Leu Arg Ser
 85 90 95

Lys Met Gly Lys Trp Cys Cys His Cys Phe Pro Cys Cys Arg Gly Ser
 100 105 110

Gly Lys Ser Asn Val Gly Thr Trp Gly Asp Tyr Asp Asp Ser Ala Phe
 115 120 125

Met Glu Pro Arg Tyr His Val Arg Arg Glu Asp Leu Asp Lys Leu His
 130 135 140

Arg Ala Ala Trp Trp Gly Lys Val Pro Arg Lys Asp Leu Ile Val Met
 145 150 155 160

Leu Arg Asp Thr Asp Met Asn Lys Arg Asp Lys Gln Lys Arg Thr Ala
 165 170 175

Leu His Leu Ala Ser Ala Asn Gly Asn Ser Glu Val Val Gln Leu Leu
 180 185 190

Leu Asp Arg Arg Cys Gln Leu Asn Val Leu Asp Asn Lys Lys Arg Thr
 195 200 205

Ala Leu Ile Lys Ala Val Gln Cys Gln Glu Asp Glu Cys Val Leu Met
 210 215 220

Leu Leu Glu His Gly Ala Asp Gly Asn Ile Gln Asp Glu Tyr Gly Asn
 225 230 235 240

```
<210> 334
<211> 384
<212> PRT
<213> Homo sapiens
```

```

<400> 334
Met Val Val Glu Val Asp Ser Met Pro Ala Ala Ser Ser Val Lys Lys
          5                      10                      15

Pro Phe Gly Leu Arg Ser Lys Met Gly Lys Trp Cys Cys Arg Cys Phe
          20                      25                      30

Pro Cys Cys Arg Glu Ser Gly Lys Ser Asn Val Gly Thr Ser Gly Asp
          35                      40                      45

His Asp Asp Ser Ala Met Lys Thr Leu Arg Ser Lys Met Gly Lys Trp
          50                      55                      60

Cys Arg His Cys Phe Pro Cys Cys Arg Gly Ser Gly Lys Ser Asn Val
          65                      70                      75                      80

Gly Ala Ser Gly Asp His Asp Asp Ser Ala Met Lys Thr Leu Arg Asn
          85                      90                      95

Lys Met Gly Lys Trp Cys Cys His Cys Phe Pro Cys Cys Arg Gly Ser
          100                     105                     110

Ser Lys Ser Lys Val Gly Ala Trp Gly Asp Tyr Asp Asp Ser Ala Phe
          115                     120                     125

```


110

Met	Glu	Pro	Arg	Tyr	His	Val	Arg	Gly	Glu	Asp	Leu	Asp	Lys	Leu	His
130						135					140				
Arg	Ala	Ala	Trp	Trp	Gly	Lys	Val	Pro	Arg	Lys	Asp	Leu	Ile	Val	Met
145					150					155					160
Leu	Arg	Asp	Thr	Asp	Val	Asn	Lys	Gln	Asp	Lys	Gln	Lys	Arg	Thr	Ala
				165					170					175	
Leu	His	Leu	Ala	Ser	Ala	Asn	Gly	Asn	Ser	Glu	Val	Val	Lys	Leu	Leu
			180					185					190		
Leu	Asp	Arg	Arg	Cys	Gln	Leu	Asn	Val	Leu	Asp	Asn	Lys	Lys	Arg	Thr
		195					200					205			
Ala	Leu	Ile	Lys	Ala	Val	Gln	Cys	Gln	Glu	Asp	Glu	Cys	Ala	Leu	Met
		210				215					220				
Leu	Leu	Glu	His	Gly	Thr	Asp	Pro	Asn	Ile	Pro	Asp	Glu	Tyr	Gly	Asn
225					230					235					240
Thr	Thr	Leu	His	Tyr	Ala	Ile	Tyr	Asn	Glu	Asp	Lys	Leu	Met	Ala	Lys
				245					250					255	
Ala	Leu	Leu	Leu	Tyr	Gly	Ala	Asp	Ile	Glu	Ser	Lys	Asn	Lys	His	Gly
			260					265					270		
Leu	Thr	Pro	Leu	Leu	Leu	Gly	Val	His	Glu	Gln	Lys	Gln	Gln	Val	Val
		275					280					285			
Lys	Phe	Leu	Ile	Lys	Lys	Lys	Ala	Asn	Leu	Asn	Ala	Leu	Asp	Arg	Tyr
	290					295					300				
Gly	Arg	Thr	Ala	Leu	Ile	Leu	Ala	Val	Cys	Cys	Gly	Ser	Ala	Ser	Ile
305					310					315					320
Val	Ser	Leu	Leu	Leu	Glu	Gln	Asn	Ile	Asp	Val	Ser	Ser	Gln	Asp	Leu
				325					330					335	
Ser	Gly	Gln	Thr	Ala	Arg	Glu	Tyr	Ala	Val	Ser	Ser	His	His	His	Val
			340					345					350		
Ile	Cys	Gln	Leu	Leu	Ser	Asp	Tyr	Lys	Glu	Lys	Gln	Met	Leu	Lys	Ile
		355					360					365			
Ser	Ser	Glu	Asn	Ser	Asn	Pro	Glu	Asn	Val	Ser	Arg	Thr	Arg	Asn	Lys
		370				375					380				

THIS PAGE BLANK (USPTO)